

## Effect of Drying Temperature on the Quality of Dry Surimi Powder from *Pangasius*

### Abstract

Demands for fish protein including dried fish protein to develop functional food are gradually growing in the world. Surimi, the concentrated myofibrillar protein extracted from fish flesh by washing minced meat, separated from bones, skin, and guts with added cryoprotectants such as sugar or alcohol (most commonly used cryoprotectant in the surimi industry is 1:1 mixture of sucrose and sorbitol at a concentration of 8%), finally stored in frozen condition in block form, is used as a raw material for preparation of number of value-added products. The dried form of surimi can be prepared from frozen surimi blocks by adopting different drying technologies and it offers many advantages such as ease of handling, lower distribution costs and more convenient storage. The present work is aimed to study the effect of drying temperature on the quality of dry surimi powder, prepared from *Pangasius* meat. A significantly higher ( $p < 0.05$ ) value of ash ( $1.83 \pm 0.47\%$ ) was recorded in surimi powder dried at  $60^{\circ}\text{C}$  than at  $50^{\circ}\text{C}$  and  $70^{\circ}\text{C}$ . The moisture content significantly ( $p < 0.05$ ) reduced at  $60^{\circ}\text{C}$  ( $9.05 \pm 0.22\%$ ) and  $70^{\circ}\text{C}$  ( $9.55 \pm 0.51\%$ ) as compared to  $50^{\circ}\text{C}$ . The quality parameters such as TVB-N, PV, pH and TPC were all found to be well within the recommended level of acceptability except for the surimi powder dried at  $50^{\circ}\text{C}$ , wherein the TVBN ( $36.24 \pm 1.26$  mg/100g) crossed the limit of acceptability. Drying temperature was found to affect the colour of the surimi powder with the optimum acceptable colour score achieved at  $60^{\circ}\text{C}$  ( $6.38 \pm 0.52$ ). At higher temperature of  $70^{\circ}\text{C}$  darkening was observed with consequent decrease in the colour scores ( $5.75 \pm 0.46$ ). Therefore, the optimal temperature for drying of *Pangasius* surimi into its powdered form was achieved during its exposure at temperature of  $60^{\circ}\text{C}$ .

**Key words:** *Pangasianodon hypophthalmus*, dry surimi powder, drying temperature, biochemical quality, microbial quality, sensory evaluation.

### Introduction

Demands for fish protein ingredients including dried fish protein to develop functional food or ready-to-eat products are gradually growing in the world (Thorkelsson *et al.*, 2009). The white flesh and low-fat content fish are considered the most suitable species for developing fish protein ingredients (Hultin *et al.*, 2005; Park and Lin, 2005), while, there are other fish protein sources that can be used for producing protein ingredients such as dark muscles/underutilized/low value fish species and fish by-products for human consumption (Arason *et al.*, 2009).

Minced meat, the most important ingredient for production of surimi and surimi-based products, is the flesh separated from the skin, bones, scales and fins of the fish. Surimi is a product of Japanese origin, derived from a traditional Japanese way of using and preserving fresh fish, refers to concentrated myofibrillar protein extracted from fish flesh by washing minced meat that has been separated from bones, skin, and guts (Santana *et al.*, 2012a). During washing with cold water, fat and other water-soluble contents are removed, whereas insoluble myofibrillar protein is isolated. After being mixed with cryoprotectant, this protein is called surimi (Okada, 1992). The addition of cryoprotectants is required to retain the functional properties of the myofibrillar proteins. The most commonly used

cryoprotectant in the surimi industry is a 1:1 mixture of sucrose and sorbitol at a concentration of 8%. Surimi generally comes in a block form and is stored in frozen condition. Surimi possess some important functional properties such as gel forming ability and water holding capacity (WHC) due to its content of myofibrillar protein that plays the most critical role during meat processing like stabilization of comminuted and restructured meat products (Zhou *et al.*, 2006). The physicochemical state of myofibrillar proteins affects the functionality of meat system and plays a direct role in determining the quality and value of processed meat (Li and Wick, 2001). Surimi is used as a raw material for preparation of number of value-added products such as fish sausage, cakes, cutlets, patties, balls, pastes, texturized products, etc. Today, surimi is a popular food item not only in Japan but also in many other countries due to its unique textural properties and high nutritional value (Park and Lin, 2005).

Surimi can be produced from both marine and fresh-water fish, including both white-muscled and dark-muscled fish, such as Alaska Pollock, blue whiting, croaker, lizardfish, sardine, tilapia and bigeye snapper (Nopianti *et al.*, 2010). Commonly, certain species are used due to their easy capture and low price. The use of alternative species in order to obtain surimi of good gel-forming ability is one of the aims of the fishing industry. As, freshwater fish are excellent sources of high-quality protein due to presence of well-balanced highly digestible essential amino acids (Karmas and Lauber, 1987), the surimi making ability of many freshwater species could be upgraded by manipulating processing techniques (Onibala *et al.*, 1997).

*Pangasianodon hypophthalmus* (Ali *et al.*, 2012), also known as *Pangasius sutchi* has been widely cultured and propagated among fish farmers and breeders especially in West Bengal and Andhra Pradesh. According to DADF, GOI, 2017, it is called as “GAME CHANGER” because of its domestication through extensive culture and large-scale production and become an important fishery due to its remarkable growth rate (attains almost 1 kg in 90 days) in Indian environment. *Pangasius* meat has high nutritive qualities and excellent sensory properties such as tender flesh, sweet taste, absence of fishy odour and spines, delicate flavour and firm texture when cooked; these are the attributes that favour consumer preference for sutchi catfish.

Recent research indicates that surimi could be converted to a dried form, surimi powder that can be kept without frozen storage. The powdered surimi offers many advantages in commerce, such as ease of handling, lower distribution costs, more convenient storage and usefulness in dry mix application (Green and Lanier, 1985). Surimi powder can be prepared

from frozen surimi blocks by adopting different drying technologies. The main purpose of drying technologies developed in food industries is to prolong the shelf life of a food product. Available drying methods for making surimi powder include freeze drying, spray drying, oven drying, solar drying, and mechanical drying where the heating temperature of evaporation can be lowered by lowering pressure using a vacuum to prevent the protein from heat denaturation.

Surimi powder can be turned into wet surimi by rehydrating it with four times its weight of water, so that wet rehydrated surimi powder would have water content similar to that of a frozen surimi block (Santana *et al.*, 2012b). Another advantage of surimi powder is its usefulness in dry mixtures (Green and Lanier, 1985) that could help industries to modify the formulation of surimi-derived products, resulting in more homogenous blends and easier protein standardization. With the advancement of food technology, processing of surimi in dried form is flourishing so that it can be used in dry mixing. Thus, the purpose of the study is to establish and standardize the hypothesis that the functional properties of surimi powder from *Pangasius* is acceptable as a commercial preparation. Thus, the objective of the present study is to study the effect of drying temperature on the quality of dry surimi.

## **MATERIAL AND METHOD**

### **Raw material**

Live pangas also known as Sutchi Catfish (*Pangasianodon hypophthalmus*) were bought from the market by simple random sampling (Garia fish market, Kolkata) and used for the present study. The fishes were transferred to the laboratory within an hour from the market in iced condition and processed immediately.

### **Washing procedure for mince and production of surimi**

Surimi production was carried out according to the method of Rawdkuen *et al.* (2009) as illustrated in Figure 1. The fish mince was washed with cold water (40°C) using a mince to washing medium ratio of 1:4 (w/v) to remove sarcoplasmic proteins, blood, pigment, fat and other low molecular weight components. The mixture was continuously stirred for 10 minutes in a cold room (40°C). The washed mince was then filtered through four layers of cheese-cloth and subsequently dewatered by using a hydraulic pressing machine. Washing was performed four times. The last washing step was carried out using 0.5% NaCl solution with mince to NaCl solution ratio of 1:4 (w/v). Finally, the meat was subjected to pressing and the final moisture content of the product was maintained about 79% level referred to as dewatered minced meat (DWM).

### **Production of surimi powder**

Surimi samples were dried in three different drying temperature viz, 50°C, 60°C and 70°C to prepare dry surimi powder (Figure 2), following the oven drying method Huda *et al.* (2012) with slight modifications.

The surimi powder was then subjected for proximate composition, biochemical, microbiological and sensory analysis to determine the suitable most drying temperature. Moisture content determination of the surimi powder was done by the method described by AOAC (1995). Total nitrogen was estimated by Kjeldahl method (AOAC, 1995). Crude protein value was calculated by multiplying the total nitrogen value by a factor of 6.25. Estimation of total lipid was done by the method described by Bligh and Dyer (1959). The ash content was measured by the method of AOAC (1995). The surimi powder was then subjected in triplicate for Total Plate Count (TPC), physicochemical and sensory analyses. TPC was determined by spread plating appropriate dilutions on Total Plate Count Agar (Hi-media) (Nath *et al.*, 2014) and results expressed as log cfu/g. The physicochemical indices like TVBN, PV and pH of surimi powder were measured by the method described by AOAC (1995), by Folchet *et al.* (1957) and by Ozyurt *et al.* (2009) (using pH meter Five Easy™ FE20, made by Mettler Todelo AG, Switzerland) respectively. Sensory evaluation was performed by a sensory panel composed of 15 experienced members based on 7-point hedonic scale as described by Siah and Tahir (2011).

All of the data were checked for normal distributions with normality plots prior to analysis of variance (ANOVA), to determine significant differences among means at  $\alpha = 0.05$  level, using statistical tools of Microsoft Office Excel (2007) and R software (Version 2.14.1).

### **RESULTS AND DISCUSSION**

In the present study, surimi powder was prepared from surimi of Sutchi Catfish (*Pangasianodon hypophthalmus*). The quality of surimi prepared from fish vary depending on species and on all those factors, which affect the composition of the fish like seasonal variation, feeding habit, pH of the habitat water, adaptation, temperature, lipid content, sex and spawning. In the present study, Sutchi Catfish (*Pangasianodon hypophthalmus*) was chosen due to easy availability and low price.

Surimi samples were dried in three different drying temperature viz, 50°C, 60°C and 70°C to prepare dry surimi powder following the oven drying method Huda *et al.* (2012) with modifications. The drying process was continued overnight until the moisture content was reduced below 15% or less. The dried surimi was then milled into powder using conventional

blender and sieved through a 30mm screen mesh. The powder thus obtained was packed in air tight container and kept under refrigeration (40°C). Proximate composition, biochemical and microbiological analysis were done to determine the suitable drying temperature. Sensory assessment was also conducted to select the suitable temperature for drying.

Proximate composition of the surimi powder varies upon different fish and different incorporation rates of sugars (sucrose, sorbitol, trehalose etc.), which were added with surimi to protect its functional properties against freezing and drying. In the present study, surimi was prepared by adding 4% sucrose, 4% sorbitol and 0.3% sodium tripolyphosphate. The final proximate composition for surimi powder obtained at 50°C was 12.98±1.31%, 57.33±2.82%, 10.13±0.67%, 1.15±0.08% and 18.11±0.39% for moisture, protein, lipid, ash and carbohydrate respectively (Table 1). A significantly higher ( $p<0.05$ ) value of ash (1.83±0.47%) was recorded in case of surimi powder dried at 60°C than at 50°C and 70°C. Drying temperature of 60°C and 70°C was also found to significantly reduce the moisture content ( $p<0.05$ ) as compared to 50°C with 9.05±0.22% and 9.55±0.51% recorded for 60°C and 70°C respectively. The drying temperatures within the range of 50°C to 70°C, however do not seem to have any influence on the composition of protein, fat and carbohydrate ( $p>0.05$ ).

At 60°C and 70°C the protein content observed was 60.86±1.61% and 61.04±1.31% respectively while the moisture content was 10.53±0.23% and 10.65±0.85% respectively. The protein contents of tilapia and trout surimi powder were reported to be 57.8% and 64.8% respectively (Huda *et al.*, 2001a) which corroborates to the results of the present findings. Similarly, a study conducted by Ramirez *et al.* (1999) postulated that freeze-dried tilapia surimi powder contains 62% protein, 4.6% moisture, 2.9% fat, 1.6% ash and 8% carbohydrate when 8% sucrose was used as cryoprotectant during surimi preparation which corroborates with the results of the present investigation. The high content of carbohydrate in surimi powder was observed due to the addition of cryoprotectants during surimi preparation. Huda *et al.* (2001) found that incorporation of 8% cryoprotectant during tilapia and trout surimi preparation resulted in a final carbohydrate content of 33.5% and 30.7% respectively, which is found to be higher than the carbohydrate content (17.66±0.59% at temperature of 60°C) observed in the present study.

EIC (1995) recommended that TVBN value of 35 mg/100g in meat is considered as limit of acceptability, above which fishery products are considered unfit for human consumption. Lakshmanan (2000) recommended a level of PV in seafoods is 10 – 20 meq O<sub>2</sub>/kg of fat as a limit of acceptability. The acceptable upper limit for the pH of fish is 6.8 – 7 (Ludorff and

Meyer, 1973). International Commission on Microbiological Specification for Foods recommended the maximum acceptable count for freshwater fish is  $10^7$  log cfu/g. The quality parameters such as TVB-N, PV, pH and TPC (Table 2) were all found to be well within the recommended level of acceptability as indicated earlier except for the surimi powder prepared at the drying temperature of  $50^{\circ}\text{C}$ , wherein the TVBN crossed the limit of acceptability reaching a value of  $36.24 \pm 1.26$  mg/100g.

Based on the TVBN values, the temperature of  $50^{\circ}\text{C}$  was inferred to be not suitable for the drying of surimi. However, considering the values of TVBN, PV, pH and TPC it was difficult to select among the drying temperatures of  $60^{\circ}$  and  $70^{\circ}\text{C}$  as the values for all these parameters were within acceptable limits for both the temperatures. Therefore, the final selection of drying temperature was done based on sensory scores of the surimi powders (Table 3).

Drying temperature was found to affect the colour of the surimi powder with the optimum acceptable colour score achieved at  $60^{\circ}\text{C}$  ( $6.38 \pm 0.52$ ). At higher temperature of  $70^{\circ}\text{C}$  darkening was observed with consequent decrease in the colour scores ( $5.75 \pm 0.46$ ). Therefore, considering the overall acceptability score the drying temperature of  $60^{\circ}\text{C}$  ( $6.25 \pm 0.24$ ) was adjudged suitable for preparation of pangus surimi powder. Thus, we can conclude from the following study that the best temperature for drying of *Pangasius* surimi into its powdered form was achieved during its exposure at temperature of  $60^{\circ}\text{C}$  assessing its sensory characteristics, though having no change in its protein, fat and ash values.

## REFERENCES

- Ali H, Haque M M and Belton B (2012) Striped catfish (*Pangasianodon hypophthalmus*, Sauvage, 1878) aquaculture in Bangladesh: an overview Authors. *Aquacult. Res.* 44(6), 950–965.

- AOAC (Association of Official Analytical Chemistry) (1995) In: *Official Methods of Analysis*, 16th edn., Arlington, VA.
- Arason A, Karlsdottir M, Valsdottir T, Slizyte R, Rustad T, Falch E, Eysturskard J and Jakobsen G (2009) Maximum resource utilisation-value added fish by-products. Nordic innovation centre, Project nr: 04275, pp 35–46.
- Bligh E G and Dyer W J (1959) A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37(8), 911-917.
- EIC (1995) Manual of Analytical Methods for Fish and Fishery Products. Quality Development Center, Ministry of Commerce, Govt. of India, Madras, India.
- Folch J, Less M and Sloane G W S (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226, 497–509.
- Green D and Lanier T C (1985) Fish as the 'soybean of the sea'. In: *Proceedings of the International Symposium on Engineered Seafood Including Surimi*, pp. 42-52 (Eds. Martin R E and Collete R L) National Fisheries Institute, Washington.
- Huda N, Abdullah A and Babji A S (2001) Functional properties of surimi powder from three Malaysian marine fish. *IJFST*. 36, 401-406.
- Huda N, Abdullah R, Santana P and Yang T A (2012) Effects of Different Dryoprotectants on Functional Properties of Threadfin Bream Surimi Powder. *JFAS*, 7(3), 215-223.
- Hultin H O, Kristinsson H G, Lanier T C and Park J W (2005) Process for recovery of functional proteins by pH shifts. In: *Surimi and surimi sea food*, pp. 107–139 (Eds. Park J W) Taylor and Francis Group, Boca Raton.
- Karmas E and Lauber E (1987) Novel products from underutilized fish using combined processing technology. *J. Food Sci.* 52,7–9.
- Lakshmann P T (2000) Fish Spoilage and Quality Assessment. In: *Quality Assessment in Seafood Processing*, pp 26-40 (Eds. Iyer T S G, Kandoran M K, Thomas M and Mathew P T) CIFT and SOFT, India.
- Li C T and Wick M (2001) Improvement of the physicochemical properties of pale soft and exudative (PSE) pork meat products with an extract from mechanically deboned turkey meat (MDTM). *Meat Sci.* 58,189-195.
- Ludorff W and Meyer V (1973) In: *Fische und fischerzeugnisse*, Vol. 6, 2<sup>nd</sup>edn.,pp. 294, Verlag Paul Parey, Berlin und Hamburg.
- Nath S, Chowdhury S, Dora K C (2014) Effect of lactic acid bacteria application on shelf life and safety of fish fillet at 6±1°C. *IJAR*. 2(4), 201-207.

- Nopianti R, Huda N and Ismail N (2010) Loss of functional properties of proteins during frozen storage and improvement of gel-forming properties of surimi. *As. J. Food Ag-Ind.* 3(06), 535-547.
- Okada M (1992) History of surimi technology in Japan. In: *Surimi Technology*, pp. 3–21 (Eds. Lanier T C and Lee C M) (Eds). Marcel Dekker Inc., New York.
- Onibala H, Takayama T, Shindo J, Hayashi S and Miki H (1997) Influence of freshness on occurrence of setting and disintegration in heat-induced gels from tilapia. *Fisheries Science.* 63:276–280.
- Özyurt G, Kuley E, Özkütük S, Özogul F (2009) Sensory, microbiological and chemical assessment of the freshness of red mullet (*Mullus barbatus*) and goldband goatfish (*Upeneusmoluccensis*) during storage in ice. *Food Chem.* 114(2), 505-10.
- Park J W and Lin T M J (2005) Surimi: manufacturing and evaluation. In: *Surimi and Surimi Seafood*, 2nd edn, pp. 33-98 (Eds. Park J W) CRC Press, Boca Raton.
- Santana P, Huda N and Yang T A (2012a) Gel Characteristics of Surimi Powder Added with Hydrocolloids. *UMT 11th International Annual Symposium on Sustainability Science and Management*, 9th -11th July, Terengganu, Malaysia.
- Santana P, Huda N and Yang T A (2012b) Technology for production of surimi powder and potential of applications. *IFRJ*, 19(4),1313-1323.
- Siah W M and Mohd Tahir S (2011) Effect of different packaging materials on the shelf life of modified atmosphere packaged red tilapia (*Oreochromis mossambica*) fillets. *IFRJ.* 18(3), 59-63.
- Thorkelsson G, Slizyte R, Gildberg A and Kristinsson H G (2009) Fish proteins and peptides. Processing methods, quality and functionality. In: *Marine functional foods*, pp. 115–133 (Eds. Luten J B) Wageningen University Press, Wageningen.
- Zhou A, Benjakul S, Pan K, Gong J and Liu X (2006) Cryoprotective effects of trehalose and sodium lactate of tilapia (*Sarotherodonnilotica*) surimi during frozen storage. *Food Chem.* 96, 96-103.

**Table 1. Proximate composition of surimi powder at different drying temperature**



Drying Temperature Parameters	50°C	60°C	70°C
Ash (%)	1.15±0.08 <sup>a</sup>	1.83±0.47 <sup>b</sup>	1.25±0.10 <sup>a</sup>
Moisture (%)	12.98±1.31 <sup>a</sup>	9.05±0.22 <sup>b</sup>	9.55±0.51 <sup>b</sup>
Fat (%)	10.13±0.67 <sup>a</sup>	10.53±0.23 <sup>a</sup>	10.65±0.85 <sup>a</sup>
Protein (%)	57.33±2.82 <sup>a</sup>	60.86±1.61 <sup>a</sup>	61.04±1.31 <sup>a</sup>
Carbohydrate (%)	18.11±0.39 <sup>a</sup>	17.66±0.59 <sup>a</sup>	17.61±0.69 <sup>a</sup>

\*Results are mean of 5 determinations with s.d.

#Values of mean within the same row with different superscripts vary significantly ( $p < 0.05$ )

**Table 2. Biochemical and microbiological analysis of surimi powder at different drying temperature**

Drying Temperature Parameters	50°C	60°C	70°C
pH	6.60±0.02 <sup>a</sup>	6.70±0.04 <sup>b</sup>	6.80±0.06 <sup>c</sup>
TVBN (mg/100g)	36.24±1.26 <sup>a</sup>	27.40±0.35 <sup>b</sup>	14.68±0.33 <sup>c</sup>
PV (meq. Active oxygen/kg)	2.97±0.32 <sup>a</sup>	3.07±0.24 <sup>a</sup>	3.07±0.31 <sup>a</sup>
TPC (logcfu/g)	3.90±0.10 <sup>a</sup>	3.84±0.08 <sup>a</sup>	3.86±0.29 <sup>a</sup>

\*Results are mean of 5 determinations with s.d.

#Values of mean within the same row with different superscripts vary significantly ( $p < 0.05$ )

**Table 3. Sensory analysis of surimi powder at different drying temperature**

Drying Temperature Parameters	Drying Temperature		
	50°C	60°C	70°C
Colour	5.88±0.83	6.38±0.52	5.75±0.46
Flavour	6.13±0.64	6.13±0.35	5.63±0.52
Texture	5.75±0.46	6.25±0.46	5.88±0.35
Overall Acceptability	5.92±0.30	6.25±0.24	5.75±0.30

\*Results are mean of 5 determinations with s.d.

#Values of mean within the same row with different superscripts vary significantly ( $p < 0.05$ )

