

## Original Research Article

# Extraction, Physicochemical Characteristics and Fatty Acids Profile of Kernel Oil from *Mangifera indica* L. Cultivated in Sudan

### ABSTRACT

**Aims:** This study was aimed to investigate the physicochemical properties and fatty acids composition of *Mangifera indica* L. seed kernel oil; in addition to investigating the effect of solvent type and extraction duration on extracts properties.

**Study design:** Extraction of *Mangifera indica* L. seed kernel oil in different trials under the same conditions using two different solvents for different time of extraction, and determining their physicochemical properties and fatty acids constituents.

**Place and Duration of Study:** This study was conducted at the Department of Applied and Industrial chemistry International University of Africa (IUA), Khartoum, Sudan, between July and November 2019.

**Methodology:** The oil from *Mangifera indica* L. seed kernel was extracted using n-hexane and petroleum ether in a soxhlet apparatus for 4 and 7 h. the physicochemical properties of the extracted oils were determined using standard official methods. Fatty acid profile of n-hexane extract was identified by gas chromatography/mass spectrometer (GC/MS) after methylation.

**Results:** n-Hexane exhibits better extraction efficiency ( $11.40 \pm 0.66\%$  for 7 h) than petroleum ether ( $10.80 \pm 0.44\%$  for 7 h). The density and refractive index of the oil were  $0.89 \pm 0.01 \text{ g/cm}^3$  and  $1.46 \pm 0.01$  at  $28^\circ\text{C}$  respectively. The physicochemical properties of n-Hexane and petroleum ether extracts were acid value ( $3.35 \pm 0.54$  and  $2.52 \pm 0.13 \text{ mg KOH/g oil}$ ), peroxide value ( $4.32 \pm 0.65$  and  $5.11 \pm 1.03 \text{ meq O}_2/\text{kg}$ ), saponification value ( $201.05 \pm 0.95$  and  $198.66 \pm 1.04 \text{ mg KOH/g oil}$ ), ester value ( $197.59 \pm 0.67$  and  $192.54 \pm 0.20 \text{ mg KOH/g oil}$ ) respectively. The statistical analysis of obtained data revealed no significant difference, at 95% confidence interval, between the standard deviation and the mean of two data sets of physicochemical properties of *Mangifera indica* L. seed kernel oils extracted with the two solvents used. GC/MS analysis revealed a total of 18 fatty acids were identified in which the majors are stearic acid (39.79%), oleic acid (36.77%), palmitic acid (10.34%), linoleic acid (6.02%) and eicosanoic acid (3.83%).

**Conclusion:** The results suggest that mango seed kernel contains stable oil which can be potentially extracted by n-hexane; however, the solvent type has no significant effect on the physicochemical properties of the extracted oil and has the potential usefulness to be used in soap industry.

**Keywords:** *Mangifera indica* L., kernel oil, physicochemical properties, fatty acid, solvent extraction.

19 **1. INTRODUCTION**

20

21 *Mangifera indica* L., commonly called mango, belongs to the family *Anacardiaceae* [1]. The  
22 mango trees can reach a height of more than 35 - 40 m, with a radius of 10 m. Its leaves are  
23 evergreen, flat, 15 - 35 cm long and 6 - 16 cm wide. Mango fruits ripen after 3 - 6 months of  
24 flowering. Ripe fruits have different sizes and colors depending on the variety [2]. Mango  
25 trees grow in the tropics and subtropics of Asia and Africa. India produces 44.14% of the  
26 world's mango production [3,4]. *Mangifera indica* L. extracts of bark, leaves, stems and  
27 unripe fruits have been conventionally used as antibiotics and in treatment of typhoid fever,  
28 dysentery, diarrhea, sore throat disease and digestive disorder [5,6]. Moreover mango seed  
29 oil contains a high level of antioxidants and free of charge radical scavenging chemical  
30 substances [7]. Mango kernel oil is rich in unsaturated fatty acids and phenolic compounds,  
31 making it used as nutritious oil and in the cosmetics industry [8]. Previous studies on the  
32 kernel of *Mangifera indica* varieties revealed high levels of saponification value ranges  
33 between 143.6 to 207 mg KOH/g oil [1,2,9]. The major fatty acids detected in  
34 *Mangifera indica* are stearic acid, oleic acid, linoleic and arachidonic acid [1,3,8]. Mango  
35 handling creates huge quantity of waste, where the peeling process and disposes of seeds  
36 bring about 45 % of the weight of the fruit as waste, Kernels take-up about 17 - 22% of the  
37 fruit [10]. Removal of this waste material may cause environmental risks which might further  
38 increase when exposed to climatic factors. Recent research has tended to utilize waste as  
39 by-products for further use and extraction of useful parts from them [11]. Since the main  
40 components of mango seeds are starch, fat and protein [10], the mango seed kernel has  
41 been successfully used in the production of starch [4] and biodegradable plastic  
42 polyhydroxyalkanoate as an alternative to glucose [12]. However, consumers consider the  
43 mango kernel as waste, so it is disposed of. Therefore, this study was aimed to investigate  
44 the physicochemical properties and fatty acids composition of *Mangifera indica* L. seed  
45 kernel oil; in addition to investigate the effect of solvent type and duration time on extracts  
46 properties.

47

48 **2. MATERIAL AND METHODS**

49

50 **2.1 Sample Collection and Preparation**

51 *Mangifera indica* L. fruits of *Totapuri* mangoes cultivar were harvested from Abu-Jubaiha  
52 city, South Kordofan State, Sudan. The pulp was separated mechanically from the seeds;  
53 and then seeds were manually cracked to obtain the kernels. The kernels were ground using  
54 a kitchen blender and pass through 2000 microns sieve. The kernel powder was then stored  
55 in a plastic container in a cool and dry place for further work.

56 **2.2 Extraction of Oil**

57 Oil was extracted from the kernel using two different solvents (n-hexane and petroleum  
58 ether) for different times of extraction (4 and 7 h) in a soxhlet apparatus, the extraction  
59 procedure was conducted in triplicate for each solvent. 140 g of kernel powder was  
60 encapsulated in gauze of canvas and inserted into the soxhlet extractor each time and the oil  
61 was extracted using the mentioned solvents for duration of 4 h and 7 h. At the end of the  
62 period, the solvent was recovered by rotary evaporator and residual oil was oven dried at 75  
63 °C for one hour. The extracted oil was then allowed to cool to a room temperature in a  
64 desiccator before analyses. The percentage extraction yield of oil was calculated using  
65 equation (2.1).

66 Percentage extraction yield of oil =  $\frac{\text{mass of oil}}{\text{mass of sample}} \times 100\%$  (2.1)

### 67 2.3 Physicochemical Characteristics of the Oil

68 The density and refractive index were determined according to the procedures described by  
69 (ASTM International) [13,14].

70 Peroxide value was measured by titration according to the American Oil Chemist' Society  
71 AOCS official method [15], the sample was dissolved in acetic acid/isooctane solution and  
72 excess amount of potassium iodide was added, the liberated iodine was titrated against  
73 standard sodium thiosulphate solution. The PV was expressed in meq O<sub>2</sub>/kg.

74 Saponification value was determined according to AOCS official method [16], two grams of  
75 the oil sample was treated with a known excess amount of alcoholic KOH, and the mixture  
76 was heated on a water bath for two minutes then the unreacted KOH was titrated with  
77 standardized hydrochloric acid using phenolphthalein as indicator. The SV was expressed in  
78 mg KOH/ g of oil using equation (2.2).

$$79 \text{ Saponification value} = \frac{(X - Y) \times N \times 56.1}{W} \quad (2.2)$$

80 Where: X = blank titrate value (ml); Y = Sample titrate value (ml); N = normality of HCl; 56.1  
81 = the molecular weight of KOH; W = weight of sample (g).

82 Acid value determined using the procedures described by (AOAC) [17]. In a typical  
83 procedure, 2.0 g of sample was dissolved in aqueous ethanol solution (1:1) and the mixture  
84 was titrated against standard KOH solution using phenolphthalein as indicator. The acid  
85 value was calculated mathematically using equation (2.3).

$$86 \text{ Acid value (mg KOH g}^{-1}\text{)} = \frac{V \times N \times 56.1}{W} \quad (2.3)$$

87 Where: V is the volume (ml) of standard KOH; N = normality of KOH; W = weight of oil used  
88 (g); the number 56.1 is the molecular weight of KOH.

89 Ester value was obtained by subtracting the acid value from the saponification value [17].  
90 Ester value represents the number of milligrams of potassium hydroxide required to saponify  
91 the esters present in one gram of the oil.

### 92 2.4 Determination of Fatty acids by GC-MS Analysis

93 Fatty acid profile of *Mangifera indica* L. seed kernel oil was identified by gas  
94 chromatography/mass spectrometer. Two grams of the sample were treated with 7 ml  
95 alcoholic sodium hydroxide solution and left to stand overnight then extracted with n-hexane.  
96 5 µl from the n-hexane extract was diluted with 5 ml of diethyl ether. The solution was filtered  
97 through a syringe filter 0.45 µm and dried with 1g of anhydrous sodium sulphate as a drying  
98 agent. 1µl of the diluted sample was injected in the GC/MS instrument. GC/MS analysis was  
99 performed with GC-QP2010-Ultra Shimadzu, coupled with Shimadzu TQ8040 plus mass  
100 spectroscopy detector. Capillary column (Rtx-5ms - 30 m × 0.25 mm × 0.25 µm). The  
101 sample was injected by using a split mode, helium as the carrier gas passed with flow rate  
102 1.61 ml/min, the temperature program was started from 60 °C to 300 °C with a rate of 10  
103 °C/min, the injection port temperature was 300 °C, the ion source temperature was 200 °C  
104 and the interface temperature was 250 °C. The sample was analyzed by using scan mode in  
105 the range of m/z 40-500 charge to ratio and the total run time was 27 minutes. Identification  
106 of components was achieved by comparing the spectral data obtained with those available in  
107 the National Institute of Standards and Technology (NIST) libraries.

## 108 2.5 Statistical Analysis

109 Oil extractions and all analyses were performed in triplicates using dry sample and the  
110 results were expressed as means  $\pm$  standard deviation. The standard deviations and the  
111 means of the two data sets of the physicochemical properties are compared using F test,  
112 equation (2.4), and Student's *t* test, equation (2.5), respectively [18]. Multiple comparisons of  
113 means were done by the LSD (least significance difference) test. 95 % confident interval was  
114 considered significant.

$$115 F_{calculated} = \frac{s_1^2}{s_2^2} \quad (2.4)$$

$$116 t_{calculated} = \frac{|\bar{x}_1 - \bar{x}_2|}{s_{pooled}} \sqrt{\frac{n_1 n_2}{n_1 + n_2}} \quad (2.5)$$

## 117 3. RESULTS AND DISCUSSION

### 118 3.1 Optimization of Solvent Used for Extraction

121 Petroleum ether and n-Hexane were used to extract oil from the mango seed kernel in  
122 different trials under the same conditions. The extraction yield, as shown in Table 1,  
123 increased as the time of extraction increases from 4 to 7 hours; for n-hexane the yield  
124 percentage was  $5.46 \pm 0.49$  % and  $11.40 \pm 0.66$  % respectively and for petroleum ether it  
125 was  $4.61 \pm 0.75$  % and  $10.80 \pm 0.44$  % respectively. Similar reports of Nwaokobia et al. [2]  
126 and Kemal et al. [19], declared that the yield has been shown to be time and particle size  
127 dependent. n-Hexane solvent gives the best yield with duration time of extraction 7 h this  
128 result is in agreement with that presented by Sikdar et al.[3], the ether extract is less than  
129 25.57% reported in a previous study [20].

130  
131  
132

**Table 1. Effect of Solvent and duration time on extraction of *Mangifera indica* L. kernel oil**

	n-Hexane		Petroleum ether	
Solvent volume (ml)	250	250	250	250
Sample used (g)	140	140	140	140
Duration time (h)	4	7	4	7
Extraction yield (%) <sup>*</sup>	$5.46 \pm 0.49$	$11.40 \pm 0.66$	$4.61 \pm 0.75$	$10.80 \pm 0.44$

133 <sup>\*</sup> Values are means of triplicate  $\pm$  standard deviations.

134  
135

### 136 3.2 Physicochemical Properties

136 The obtained results presented in Table 2 showed that there is no significant difference in  
137 density and refractive index of *Mangifera indica* L. kernel oil extracted by n-hexane and  
138 petroleum ether. The density of mango kernel oil was between  $0.89 \pm 0.01$  g/cm<sup>3</sup>; this value  
139 is fall within the range reported in previous studies [9,21]. The refractive index was found to  
140 be  $1.46 \pm 0.01$  at 28 °C for both n-hexane and petroleum ether extracts. This value is agreed  
141 with that obtained by Kemal et al. [1], Nzikuo et al. [21] and Nwaokobia et al. [2] which lies  
142 within the range of some butter and edible oils like cocoa butter (1.455 to 1.458), cotton seed  
143 oil (1.458 to 1.466) and shea butter (1.463 to 1.468) [1].

144 Peroxide value is one of the most widely used testing for oxidative rancidity in oils; it is a  
145 very useful parameter for appreciating the first stages of oxidative deterioration. The results  
146 showed that the peroxide values of *Mangifera indica* L. kernel oil ( $4.32 \pm 0.65$  to  $5.11 \pm 1.03$   
147 meq O<sub>2</sub>/kg oil) are lower than the allowed value for crude vegetable oils.

148 Basically, the acid value is used to quantify the amount of acid (free fatty acids, acid  
149 phosphates or amino acids) present in a sample. For oils, it is a measure of the free fatty  
150 acid content. From Table 2 below it is shown that both n-hexane and petroleum ether extract  
151 have low acid values,  $3.35 \pm 0.54$  and  $2.52 \pm 0.13$  mg KOH/g oil respectively. These values  
152 are less than the Codex standard value for virgin vegetable oils ( $4.0$  mg KOH g<sup>-1</sup> Oil) [22].  
153 The acid value of both extracts agreed with that obtained by Kemal et al. for Ethiopian  
154 *Mangifera indica* seed kernels ( $2.39$  mg KOH/g) [19].

155 The saponification values ( $201.05 \pm 0.95$  mg KOH/g for n-hexane extract and  $198.66 \pm 1.04$   
156  $95$  mg KOH/g for petroleum ether extract) are significantly same. Hence, the saponification  
157 value of mango oil is not dependent on the extraction solvent used. A high saponification  
158 value may suggest use of the oil in the soap industry. Therefore, mango oil has a very high  
159 chance of being used for the manufacturing of soap. Both saponification values of the  
160 mango oil falls within the literature range [2,21,23].

161 The ester value was high in hexane extract  $197.59 \pm 0.67$  mg KOH/g oil than petroleum  
162 ether extracts  $192.54 \pm 0.20$  mg KOH/g oil. Both ester values fall within the literature range  
163 of ester values [2,21].

164 The results of statistical analysis of data were presented in Table 3. The results revealed  
165 that the values of  $F_{\text{calculated}}$  for seven properties are less than  $F_{\text{table}}$  ( $= 19.0$ ) [18], this indicated  
166 that the standard deviations of the two data sets are not significantly different from each  
167 other at 95 % confident interval. The comparison between the means of the two data sets  
168 was performed by student's t-test, equation (2.5) the values of  $t_{\text{calculated}}$  are obviously less  
169 than the critical value for  $t_{\text{table}}$  ( $= 2.776$ ) for 95% confidence and 4 degrees of freedom [18].  
170 Therefore, there is more than a 5% chance that the two sets of results lie within experimental  
171 error of each other. It was concluded that the results are not significantly different at the  
172 chosen confidence level (95%).

173  
174  
175

**Table 2: Physicochemical properties of *Mangifera indica* seed kernels oil**

property	Hexane Extract	Petroleum ether Extract
Density (g/cm <sup>3</sup> )	$0.89 \pm 0.01$	$0.89 \pm 0.01$
Refractive index	$1.46 \pm 0.01$	$1.46 \pm 0.01$
Peroxide value (meq O <sub>2</sub> /kg)	$4.32 \pm 0.65$	$5.11 \pm 1.03$
Acid value (mg KOH/g)	$3.35 \pm 0.54$	$2.52 \pm 0.13$
Saponification value (mg KOH/g)	$201.05 \pm 0.95$	$198.66 \pm 1.04$
Ester value (mg KOH/g)	$197.59 \pm 0.67$	$192.54 \pm 0.20$

176 \*Values are means of triplicate  $\pm$  standard deviations. ( $n = 3$  and  $P = .05$ )

177  
178  
179  
180  
181  
182

183 **Table 3: Calculated F and student's t values**  
 184

	Yield% (4h)	Yield% (7h)	D	RI	AV	PV	SP
F <sub>calculated</sub>	2.35	2.26	1.47	2.05	18.47	1.53	1.20
t <sub>calculated</sub>	1.65	0.95	0.23	0.28	2.51	1.40	2.43

185 \*Abbreviations: D = density, RI = refractive index, AV = acid value, PV = peroxide value and SP =  
 186 Saponification value. Confidence interval 95%, n<sub>1</sub> = 3 and n<sub>2</sub> = 3.  
 187

188 **3.3 GC/MS Analysis**


189 Fatty acids profile of *Mangifera indica* L. kernel oil was determined using GC/MS the  
 190 obtained results were shown in Table 3 and the chromatogram of Fig. 1.

191 The GC-MS data revealed the presence of 18 fatty acids. The major identified fatty acids  
 192 were stearic acid (39.79 %), oleic acid (36.77 %), palmitic acid (10.34 %), linoelaidic acid  
 193 (6.02 %) and eicosanoic acid (3.83 %). These results were compared to the results obtained  
 194 by Sikdar et al. [3], where it found that their stearic acid and oleic acid (43.32 % and 42.25 %  
 195 respectively) were higher than our obtained results for the same acids. About 55.98 % of the  
 196 fatty acid contents of *Mangifera indica* L. kernel oil are saturated and the unsaturated fatty  
 197 acids represent approximately about 43.2 % of the total fatty acids.  
 198

199 **Table 3: Main fatty acids content of *Mangifera indica* L. kernel oil**  
 200

Lipid numbers	Common (IUPAC) name	Formula	Ret. Time	Area %
<b>Saturated fatty acids</b>				
C16:0	Palmitic acid (hexadecanoic acid)	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	15.284	10.34
C17:0	Margaric acid (Heptadecanoic acid)	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	16.258	0.21
C18:0	Stearic acid (Octadecanoic Acid)	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	17.234	39.77
C20:0	Arachidic acid (Eicosanoic acid)	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	18.943	3.83
C22:0	Behenic acid (Docosanoic acid)	C <sub>22</sub> H <sub>44</sub> O <sub>2</sub>	20.560	0.81
C24:0	Lignoceric acid (Tetracosanoic acid)	C <sub>24</sub> H <sub>48</sub> O <sub>2</sub>	22.061	1.02
<b>Monounsaturated fatty acids</b>				
C18:1n-9	Oleic acid ((Z)-octadec-9-enoic acid)	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	17.011	36.77
C20:1n-11	Eicosenoic acid ((Z)-icos-11-enoic acid)	C <sub>20</sub> H <sub>38</sub> O <sub>2</sub>	18.741	0.41
<b>Polyunsaturated fatty acid</b>				
C18:2n-9,12	Linoelaidic acid ((9E,12E)-octadeca-9,12-dienoic acid)	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	16.934	6.02

201



202  
203

Fig. 1. GC Chromatogram of fatty acids of *Mangifera indica* oil

204  
205  
206

#### 4. CONCLUSION

207 In this study oil was effectively extracted from *Mangifera indica* L. seed kernel (which is  
208 generally generated as waste), using n-hexane and petroleum ether as extracting solvents.  
209 The extraction yield was found to be time dependent; n-hexane gave a higher yield than  
210 petroleum ether. However, the solvent type has no significant effect on physicochemical  
211 characteristics of the extracted oils. The results showed relatively low acid and peroxide  
212 values and high saponification and ester values. This indicates good stability of the oil and  
213 gives it potential usefulness in soap industry. The GC-MS analysis showed that *Mangifera*  
214 *indica* L. seed kernel oil has got 18 fatty acids, the predominates of them are stearic acid  
215 (39.79%), oleic acid (36.77%), palmitic acid (10.34%), linoelaidic acid (6.02%) and  
216 eicosanoic acid (3.83%).

217  
218  
219  
220  
221

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

222  
223  
224  
225

#### REFERENCES

- 226 1. Kassi ABB, Soro Y, Koffi END, Sorho S. Physicochemical study of kernel oils from  
227 ten varieties of *Mangifera indica* (Anacardiaceae) cultivated in Cote d'Ivoire. Afr. J.  
228 Food Sci. 2019;13(7):135-42.  
229 Available: <https://academicjournals.org/journal/AJFS/article-abstract/C5F418561462>  
230 2. Nwaokobia K, Idibie C, Ogboru R, Okolie P. Effects of extraction solvents on the  
231 yield and physicochemical properties of *Mangifera indica* L. seed oil. The  
232 Pharmaceutical and Chemical Journal. 2018;5(2):136-43.

- 233 Available: <http://tpcj.org/archive/volume-5-issue-2-2018>  
234  
235 3. Sikdar DC, Siri H, Vinay S, Varsha S, Rao R. Solvent extraction of mango  
236 (*Mangifera indica* L.) seed kernel oil and its characterization. International Journal of  
237 Technical Research and Applications. 2017;5(4):43-47.  
238 Available: <https://www.ijtra.com/ijtra-issue174.php>  
239  
240 4. Sonthalia M, Sikdar DC. Production of starch from mango (*Mangifera Indica* L.) seed  
241 kernel and its characterization. International Journal of Technical Research and  
242 Applications. 2015;3(3):346-49.  
243 Available: <https://www.ijtra.com/ijtra-issue153.php>  
244  
245 5. Kamble VA, Somkuwar DO, Wankhade SJ. Chemical composition, antioxidant and  
246 antisalmonella activity of *Mangifera indica* L. flower and seed kernel extracts. Int.  
247 Curr. Pharm. J. 2016;5(10):82-93.  
248 Available: <http://www.icpjonline.com/archives/Archive-5-10.htm>  
249  
250 6. Okwunodulu FU, Chukwuemeka-Okorie HO, Okorie FC. Biological synthesis of  
251 cobalt nanoparticles from *Mangifera indica* leaf extract and application by detection  
252 of manganese (II) ions present in industrial wastewater. *Chem. Sci. Int. J.*  
253 2019;27(1):1-8.  
254  
255 7. Khammuang S, Sarnthima R. Antioxidant and antibacterial activities of selected  
256 varieties of Thai mango seed extract. *Pak. J. Pharm. Sci.* 2011;24(1):37-42.  
257 Available: [http://www.pjps.pk/?page\\_id=258](http://www.pjps.pk/?page_id=258)  
258  
259 8. Kittiphoom S, Sutasinee S. Mango seed kernel oil and its physicochemical  
260 properties. International Food Research Journal. 2013;20(3):1145-49.  
261 Available: <http://www.ifrj.upm.edu.my/volume-20-2013.html>  
262  
263 9. Sadiq AS, Elijah IR, Aroke UO. Extraction and assessment of physicochemical  
264 properties of rosigold mango (*Mangifera indica*) seed kernel oil for biosesin  
265 production. AZOJETE. 2017;13(5):643-54. Accessed 15 July 2019.  
266 Available: [www.azojete.com.ng](http://www.azojete.com.ng)  
267  
268 10. Kittiphoom S. Utilization of mango seed. *Int. Food Res. J.* 2012;19(4):1325-35.  
269 Available: <http://www.ifrj.upm.edu.my/volume-19-2012.html>  
270  
271 11. Yatnatti S, Vijayalakshmi D. Study of Soup Mix Incorporated with Starch Extract  
272 from Mango "*Mangifera indica*" Seed Kernels. *Curr. Res. Nutr Food Sci Jour.*  
273 2018;6(3):816-25.  
274 Available: <http://www.foodandnutritionjournal.org/toc/?vol=6&no=3>  
275  
276 12. Nasir-Naeem KO, Shittu KO, Kabiru AY. Production and characterization of  
277 polyhydroxyalkanoate (PHA) using mango seed kernel as an alternative to glucose.  
278 *Br. Biotechnol. J.* 2016;13(3):1-11.  
279  
280 13. ASTM D4052-18a. Standard test method for density, relative density, and API  
281 gravity of liquids by digital density meter. West Conshohocken PA: ASTM  
282 International; 2011.  
283  
284 14. ASTM D1747-09(2019). Standard test method for refractive index of viscous  
285 materials. West Conshohocken PA: ASTM International; 2019.  
286  
287 15. American Oil Chemist' Society AOCS. Official method Cd 8b-90. Peroxide value,  
288 acetic acid, isooctane method. 7th ed. USA: Official Methods and Recommended  
289 Practices of the AOCS; 2017.  
290  
291 16. American Oil Chemist' Society AOCS. Official method Cd 3-25. Saponification value  
292 of fats and oils. 7<sup>th</sup> ed. USA: Official Methods and Recommended Practices of the  
293 AOCS; 2017.  
294  
295 17. AOAC. Official methods of analysis. 15th ed. Washington: Association of Official  
296 Analytical Chemists; 1990.  
297  
298 18. Harris DC. Quantitative chemical analysis. 8th ed. New York: Freeman and  
299 Company; 2010.  
300  
301 19. Kemal M, Jabasingh SA, Yimam A, Kumar JA. Extraction and characterization of oil  
302 from Ethiopian *Mangifera indica* seed kernels. *J. Sci. Ind. Res.* 2018;77(2):131-35.



- 286 Available: <http://nopr.niscair.res.in/handle/123456789/43537>  
287 20. Olajumoke OE. Extraction and characterization of vegetable oil from mango seed,  
288 *Mangifera indica*. IOSR-JAC. 2013;5(3):06-08.  
289 Available: <http://www.iosrjournals.org/iosr-jac/pages/v5i3.html>  
290 21. Nzikou JM, Kimbonguila A, Matos L, Loumouamou B, Pambou-Tobi NPG, Ndangui  
291 CB, et al. Extraction and characteristics of seed kernel oil from mango (*Mangifera*  
292 *indica*). Res. J. Environ. Earth Sci. 2010;2(1):31-3.  
293 Available: <https://maxwellsci.com/jp/issue.php?jid=RJEES&no=28>  
294 22. Alimentarius C. Codex standard for edible fats and oils not covered by individual  
295 standards. Codex stan. 1999;19:1981.  
296 23. Fahimdanesh M, Bahrami ME. Evaluation of physicochemical properties of Iranian  
297 mango seed kernel oil. In International Conference on Nutrition and Food Sciences.  
298 2013;53(9):44-49.  
299

UNDER PEER REVIEW