

Effect of Tillage and Water Management on Aggregate Stability of a Gleyic Cambisols

Abstract

The objective of this study was to determine the effect of tillage and water management on the physical stability of irrigated lowland rice field, in Kwalkwalawa, Sokoto State. The experiment was carried out in a farmer's field, near the Usmanu Danfodiyo University Teaching and Research Farm, Kwalkwalawa, Sokoto State. The coordinates of the area were taken using global positioning system (GPS) model Garmin etrex 20.0, which shows that the area is located on (N13°05.963'E005°12.650' and at 252m asl). The soils of the study area were classified as Aeric Endoaquepts at subgroup level in the USDA Soil Taxonomy System which correlated with Gleyic Cambisols in the World Reference Base. The treatments consisted of factorial combination of two tillage systems (conventional tillage (CT) and reduced tillage (RT), three irrigation water managements (Alternate one, two and three days irrigation intervals, which were carried out from one week after transplanting to hard dough stage) and three rice varieties (FARO 44, 60 and 61) all laid in a split plot design and replicated three times. After harvest, disturbed soil samples were collected with the aid of soil augar, prepared and passed through 5 mm sieve for aggregate stability determination. Result reveals that a consistent trend in aggregate size fraction was observed between the two tillage systems in both years, were a significant decrease in values of aggregate fractions of both the CT and RT in 2019 compared to in 2018. Aggregate size fraction of 5-2 mm had a significantly high value of RT compared to CT, 2-0.25 mm fractions were at par (0.41) while a greater value of aggregate fractions for CT in both 0.25-0.005 mm and <0.005 mm were observed compared to RT. Alternate days to irrigation and sampling depths increase with a corresponding in all the aggregate size classes in the two years of this study. A significant difference in MWD and GMD between the two tillage, water management and depth was noticed in both years of the trial where RT and alternate one day had high value while MWD and GMD increase with increasing depth.

Keywords: Tillage and Aggregate Stability.

1.0 INTRODUCTION

Soil aggregation is defined as the process whereby individual soil particles are joined into compound particles, clusters or aggregates (Sullivan, 2004). Aggregation is an important indicator for the soils' overall quality (Boix-fayos *et al.*, 2001). It has potential benefits on soil moisture status, nutrients dynamics, tilth maintenance and erosion reduction (Kemper and Rosenau, 1986). The term aggregate stability is used by soil physicists to refer to the ability of the bonds of the aggregation to resist stress upon exposure (Rohoskova and Valla, 2004). In

determining the stability of soil aggregates, soil physicists generally subject samples of aggregates to artificially induced forces designed to simulate phenomena which are likely to occur in the field (Hillel, 1980). Resistance of soil solids to the mechanical abrasion arising from the two forces of destruction i.e water and air for long has been used to measure stability of aggregates. The technique for aggregate stability analysis is described by Kamper and Rosenau (1986). Dry sieving by White (1993) is the common method employed to simulate aggregate resistance to wind erosion.

Numerous factors affect stability of soils (Kay, 1997), most of which can be broadly grouped into two: endogenous and exogenous factors. The endogenous factors are those that are due to inherent soil properties. These factors include soil characteristics such as texture, clay mineralogy and nature of exchangeable cations, and quantity and quality of humus fractions (Lal and Shukla, 2004). The exogenous factors that affect soil aggregate stability include weather, biological processes, land use and management. Soil management or land use has been reported to influence soil aggregate stability (Six *et al.*, 2000 and Castro *et al.*, 2002).

Pinheiro *et al.*, (2004) found that soil surface aggregates of >2mm fractions formed (50%) soil under no-tillage compared with animal traction (35%) and conventional tillage (30%). He also observed that soil organic carbon (SOC) was higher under no-tillage than conventional tillage. Gajic *et al.* (2006) reported that conversion of forest to continuous cropping by conventional cultivation significantly decreased the stability of soil aggregates in the plough horizon. Also, Adesodun and Odejimi, (2010) reported that mean weight diameter (MWD) was significantly higher in un-cultivated land, whereas addition of compost to the cultivated land improved stability of the soil. It has been reported that soil organic matter and clay content are the main soil properties affecting soil aggregation, as they binds individual particles (Fryrear *et*

al., 1994 and Oliveira *et al.*, 2004). Any practices that reduce organic matter and disturbed clay particles reduce soil aggregation.

This work was developed with the objectives of evaluating the aggregate size fractions, MWD and GMD between the two tillage operations, water managements and depths.

2.0 MATERIALS and METHODS

2.1 Experimental Location

The experiment was conducted in a farmer's field, near the Usmanu Danfodiyo University Teaching and Research Farm, Kwankwalawa, Sokoto State. The coordinates of the area were taken using global positioning system (GPS) model Garmin etrex 20.0, which shows that the area is located on latitude 13⁰05`N and longitude 05⁰12`E. The soils of the study area were classified as Aeric Endoaquepts at subgroup level in the USDA Soil Taxonomy System (USDA, 2014) which correlated with Gleyic Cambisols in the World Reference Base (FAO, 2015). The area experiences a long dry season from October to May and a short rainy season from June to September. The dry season consists of a cold dry spell (Harmattan) roughly from November to February, followed by a hot dry spell from March to May. The rainfall is erratic, small in quantity and uneven distribution with peak in August and temperature fluctuates roughly between 40⁰C maximum and 15⁰C minimum (Noma, 2005).

2.2 Treatments and Experimental Design

The treatments consisted of factorial combination of two tillage systems (Conventional tillage (CT); which involves cutting, inverting, puddling and leveling the field plots and reduced tillage (RT); which involves puddling and leveling of the plots all with local hoes, shovels and rakes), three water managements (Alternate one day, two days and three days irrigation intervals, which were carried out from one week after transplanting to hard dough stage) and three rice varieties; (FARO 44, 60 and 61).

The treatments were laid in a split plot design replicated three times. Tillage system was allocated to the main plots, while water managements and varieties were allocated to the sub-plots. Field observations and measurements were made for the two consecutive seasons using the same experimental design and field layout. After each seasons harvest, disturbed soil samples were collected with the aid of soil auger, prepared and passed through a 5mm sieve, while the aggregate stability was determined by dry sieving methods as described by Van Bavel (1950) as modified by Kemper and Rosenau (1986). Two hundred grams of bulk soil from the 5mm sieve was weighed and transferred into nest of sieves consisting of diameters 2.0 mm, 0.25 mm and 0.005 mm. The nest of sieves was placed on a mechanical shaker and shaken for 2 minutes after which the weight of soil retained in each sieve was determined.

Aggregate size distribution was calculated as the proportion of soil retained in each sieve as:

$$\text{Aggregate size distribution} = \frac{\text{Weight of soil retained in a sieve}}{\text{Total weight of soil taken}}$$

While the mean weight diameter (MWD) and geometric mean diameter (GMD) was calculated as follows:

$$\text{MWD}_{\text{dry}} = \sum_{i=1}^n w_i x_i$$

100 x_i = The mean diameter of the class (mm)

101 w_i = the proportion of each size class with respect to the total sample

102
$$\text{GMD}_{\text{dry}} = \exp\left[\frac{\sum_{i=1}^n \log x_i}{\sum_{i=1}^n w_i}\right]$$

103 w_i = weight of aggregate of each size class (g)

104 $\log x_i$ = logarithm of the mean diameter of the size classes.

105 **2.3 Data Analysis**

106 Data generated were subjected to analysis of variance (ANOVA) using SAS 9.0 software,
107 (SAS, 2002). Significant means were compared using Duncan multiple range test (DMRT) at
108 $P < 0.05$ (Gomez and Gomez, 1984).

109 **3.0 RESULTS and DISCUSSION**

110 **3.1 Effects of Tillage, Water Management and Depths on Aggregate Size Fractions in** 111 **2018**

112 From the result (Table 1), analysis of variance revealed a significant effect of tillage on
113 aggregate size fractions in all the separates, with CT having a significantly higher values, except
114 in 5-2 mm fractions, where RT have significant higher value compared to CT.

115 The results also shows a significant difference in all the aggregate size fractions with alternating
116 days of irrigations; with alternating one day irrigation having more of the aggregates size
117 fractions within all the separates while alternating three days irrigation having the least value
118 across all the separates.

119 However, sampling depth shows significant effects as they increased with depth. The observed
120 differences in the different aggregate separates between the two tillage and water management

may be as result of the degree of soil disturbances and the variation in timing of the water application among the treatment combination, this made the CT with alternate one day irrigation prone to ease of removal and being dislodged. Oades, (1984) observed that macro and micro aggregates depends on organic matter for stability against destructive forces cause by quick wetting. Aggregates were more disrupted in CT compare to RT according to Mrabet, (2002). The decline in the size of the aggregates due to tillage could be credited to mechanical disruption of macro aggregates which may have exposed soil organic matter previously protected against oxidation. This is similar to the findings of Elijah (2016).

Significant interactions between the tillage and depth were noted among all the aggregate size fractions across all the separates as depicted in Figure 1

Table 1. Effects of tillage, water management and Depths on Aggregate size fractions in 2018

Treatment	5-2 mm	2-0.25 mm	0.25-0.005 mm	<0.005 mm
Tillage (T)				
CT	0.21b	0.42a	0.51a	0.058a
RT	0.22a	0.41b	0.48b	0.048b
SE±	0.0018	0.0044	0.0021	0.0020
Water management (W)				
Alternate one day (W1)	0.22a	0.42a	0.50a	0.059a
Alternate two days (W2)	0.21b	0.42a	0.49b	0.053b
Alternate three days (W3)	0.21b	0.41b	0.48c	0.048c
SE±	0.0022	0.0054	0.0024	0.0024
Depth				
0-10 cm	0.18b	0.39b	0.46b	0.024b
10-20 cm	0.24a	0.45a	0.52a	0.083
SE±	0.0018	0.0044	0.0021	0.0020
Interaction				
T × W	NS	NS	NS	NS
T × D	**	**	**	**
W × D	NS	NS	NS	NS
T × W × D	NS	NS	NS	NS

Means followed by the same letter(s) within the same column are not significant at 0.05 level of probability, RT=Reduced tillage, CT= Conventional tillage, SE±=Standard error




Fig.1: Interactions between tillage and depths on aggregate size fractions in 2018 (SE± represented in bar)

3.2 Effects of Tillage, Water Management and Depths on Aggregate Size Fractions in 2019

There was a significantly decreased in values of aggregates fractions of both the CT and RT in 2019. Table 2 reveals that aggregate size fraction of 5-2 mm had a significantly high value of RT compared to CT, 2-0.25 mm fraction were at par (0.41) while a similar trend of greater values of the remaining aggregate fractions for CT in both 0.25-0.005 mm and <0.005 mm were observed compared to RT.

Considering the alternating irrigation with the variation in all the aggregate size classes, one day alternate irrigation had the highest mean values among all the aggregate size classes among the separates while three days alternate irrigation is the least with the value (0.045) of alternate two days irrigation for <0.005 mm fraction being similar with (0.047) of alternate three days irrigation for the same aggregate.

Similar trend was also observed in the sampling depths as in 2018, with a significant effect as they increased with depth. Significant interaction between the tillage and depth was observed among all the aggregate fractions in 2019 except on <0.005 mm aggregate fraction which shows no significant interaction. This consistent trend in the variation of all the aggregates may be as a result of the degree of disruption of all the aggregates fractions both in CT and RT, with CT been more pronounced in the second year due to continuous disturbance of those soil without subsequent addition of organic matter that will boost the bond strength of those soils. Enjugu (2014) and Elijah (2016) also observed a steady increase in aggregate size fraction over a two year tillage research in Samaru.

Table 2. Effects of tillage, water management and Depths on Aggregate size fractions in 2019

Treatment	5-2 mm	2-0.25 mm	0.25-0.005 mm	<0.005 mm
Tillage (T)				
CT	0.20b	0.41	0.50a	0.055a
RT	0.21a	0.41	0.47b	0.043b
SE \pm	0.0021	0.0050	0.0026	0.0025
Water management (W)				
Alternate one day (W1)	0.21a	0.42a	0.49a	0.055a
Alternate two days (W2)	0.21a	0.41b	0.49a	0.045b
Alternate three days (W3)	0.20b	0.40c	0.48b	0.047ab
SE \pm	0.0026	0.0062	0.0032	0.0034
Depth				
0-10 cm	0.18b	0.38b	0.45b	0.022b
10-20 cm	0.24a	0.44a	0.52a	0.077a
SE \pm	0.0021	0.0050	0.0026	0.0025
Interaction				
T \times W	NS	NS	NS	NS
T \times D	**	**	**	NS
W \times D	NS	NS	NS	NS
T \times W \times D	NS	NS	NS	NS

Means followed by the same letter(s) within the same column are not significant at 0.05 level of probability, RT=Reduced tillage, CT= Conventional tillage, SE \pm =Standard error




Fig. 2: Interactions between tillage and depths on aggregate size fractions in 2019 (SE± represented in bar)

3.3 Effects of Tillage, Water Management and Depths on MWD and GMD of Aggregate in 2018 and 2019

The mean weight diameter (MWD) and geometric mean diameter (GMD) were significantly affected by tillage (Table 3) in the two years of this study. In 2018, a significant difference was noticed in tillage with RT shows significantly higher value for MWD and GMD. The alternating days to irrigation also shows a significant increase in values of both the MWD and GMD, where alternate one day irrigation records the highest value while alternate three days irrigation had the least.

Significant increase in MWD and GMD across the two depth increase with increasing depth in 2018, and there was significant interaction between tillage and depth among the MWD and GMD of the aggregate in 2018 (Fig. 3)

A similar trend in MWD and GMD were also observed in 2019 (Table 3). This trend shows a gradual decrease in the value of both MWD and GMD for the two tillage systems across the two years. This shows the decrease of the stability of the soil to various tillage operations. Unger, (1991) reported that an increase in MWD and GMD is its ability to withstand erosion. The MWD and GMD increased as the depth of sampling increases indicating that the deeper the soil, the less vulnerable the soil will be exposed to erosion. Oguike and Mbagwu, (2009) observed that tillage with traditional hoeing and clean weeding together with reduced organic matter content, as in the case of this trial field may explain the low value of MWD and GMD observed under CT compared to the RT tillage practices. Fuentes *et al.* (2011) and Enjugu (2014) also reported that soil with conventional tillage plus reduced organic matter had a low MWD and GMD compared to soil with no-tillage plus residue, which indicates that despite the incorporation of residues, there was a negative effect on soil stability with tillage.

219 The variation of MWD and GMD and soil depth could be probably due to reduced soil
220 disturbance, redistribution of residues and homogenization effect of plowing (Tan *et al.*, 2004)

221 Table 3 Effects of tillage, water management and Depth on Mean Weight Diameter and
222 Geometric Mean Diameter in 2018 and 2019

Treatment	2018		2019	
	MWD	GMD	MWD	GMD
Tillage (T)				
CT	1.09b	0.80	1.08b	0.79b
RT	1.15	0.82	1.14a	0.81a
SE±	0.0011	0.0083	0.0011	0.0027
Water management (W)				
Alternate one day (W1)	1.13a	0.82a	1.12a	0.81a
Alternate two days (W2)	1.12b	0.81b	1.11b	0.80b
Alternate three days (W3)	1.11c	0.80c	1.11b	0.80b
SE±	0.0013	0.0010	0.0013	0.0033
Depth				
0-10 cm	1.09b	0.78b	1.08b	0.77b
10-20 cm	1.15a	0.84a	1.15a	0.83a
SE±	0.0011	0.0083	0.0011	0.0027
Interaction				
T × W	NS	NS	NS	NS
T × D	**	**	**	**
W × D	NS	NS	NS	NS
T × W × D	NS	NS	NS	NS

223 Means followed by the same letter(s) within the same column are not significant at 0.05 level of
224 probability, RT=Reduced tillage, CT= Conventional tillage, SE±=Standard error
225

226

227

228




Fig. 3: Interactions between tillage and depths on MWD and GMD in both 2018 and 2019 (SE \pm represented in bar).

4.0 CONCLUSION

The use of tillage implements interferes with the stability of the soil aggregate, where reduced tillage have higher MWD and GMD compared to conventional tillage.

238

239

REFERENCES

- 240 Adesodun, J.K. and Odejimi, O.E. (2010). Carbon-nitrogen sequestration potentials and
241 structural stability of a tropical Alfisol as influenced by pig-composted manure.
242 *International Agrophysics*, **24**:333-338.
- 243 Boix-fayos, C., Calvo-cases, A., Imeson, A.C. and Soriano-sato, M.D. (2001). Influence of soil
244 properties on the aggregation of some Mediterranean soils and the use of aggregate size
245 and stability as land degradation indicators. *Catena*, **44**: 47-67.
- 246 Castro Filho, C., Lourenco, A ., Guimaraes, M.F., Fonseca, I.C.B. (2002). Aggregate stability
247 under different soil management systems in a Red Latosol in the sate of Paraná, Brazil.
248 *Soil and Tillage Research*, **65**:45-51
- 249 Elijah, O.O. (2016). Contribution of conservation agriculture to improvement of soil quality and
250 maize (*Zea mays* L.) yield in Samaru, Northern Guinea Savanna of Nigeria. An
251 unpublished M.Sc thesis, submitted to the Department of Soil Science, Ahmadu Bello
252 University, Zaria.
- 253 Enjugu, A.M. (2014). Effect of tillage practices on soil carbon stock, hydro-physical properties
254 and yield of maize in a Northern Guinea Savanna alfisol. An unpublished M.Sc thesis,
255 submitted to the Department of Soil Science, Ahmadu Bello University, Zaria.
- 256 FAO (2015). World Reference Base for Soil Resources 2015. International Soil Classification
257 System for Naming Soil and Creating Legends for Soil Maps.
- 258 Fryrear, D.W., Krammes , C.A., Williamson, C.M. and Zobeck, T.M. (1994). Computing the
259 wind erodible fraction of soils. *Journal of Soil and Water Conservation*, **49**:183-188.
- 260 Fuent, M., Claudia H., Jorge, E., Fernando, D.L., Armando, g., Luc, d., Nele, V., and Bram,
261 G. (2011). Conservation agriculture increased organic carbon in the top soil macro-
262 aggregates and reduced soil CO₂ emmisions. *Plant soil*.
- 263 Gajic, B., Dugalic, G. and Djurovic, N. (2006). Comparison of soil organic matter content,
264 aggregate composition and water stability of Gleyic Fluvisol from adjacent forest and
265 cultivated areas. *Agronomy Research*, **4(2)**:499-508.
- 266 Gomez, K.A. and Gomez, A.A. (1984) *Statistical Procedures for Agricultural Research*, Wiley
267 Inter-science, New York, NY. 704 pp
- 268 Hillel, D. (1980). *Fundamental of Soil Physics*. Academic Press, Inc. New York. 109pp.
- 269 Kay, B.D. (1997). Rates of change in soil structure under difficult cropping systems. *Advance*
270 *Soil Science*, **12**:1-52.
- 271 Kember, W.D. and Rosenau, R.C. (1986). Aggregate stability and size distribution. In: *Methods*
272 *of Soil Analysis*. Part 1, 2nd Ed. A. Klute (Ed.). Agronomy monograph. ASA and SSSA,
273 Madison, WI, 425-442pp.
- 274 Lal, R. and Shukla, M.K. (2004). *Principles of Soil Physics*. Marcel Dekker, Inc. 18, 377pp.

- Mrabet, R. (2002). Stratification of soil aggregation and organic matter under conservation tillage systems in Africa. *Soil and Tillage Research* **66**:119-128.
- Noma, S.S. (2005). Properties, genesis and classification of soils of Sokoto-Rima floodplains at Sokoto, Nigeria. An unpublished Ph.D thesis, submitted to the Department of Soil Science and Agricultural Engineering, Usmanu Danfodiyo University, Sokoto.
- Oades, J.M., (1984). Soil organic matter band structural stability: mechanisms and implications for management. *Plant soil*, **76**:319-337.
- Oguike P.C. and Mbagwu J.S.C (2009) Variations in Some Physical Properties and Organic Matter Content of Soils of Coastal Plain Sand under Different Land Use Types. *World Journal of Agricultural Sciences* **5** (1): 63-69, 2009 ISSN 1817-3047.
- Oliveira, G.C., Dias. Junior., M.S., Resckn, D.V.S. and Curi, N.(2004). Chemical and physical-water characterization of a Red Latosol after twenty years of soil management and cultivation. *Brazilian Journal of Soil Science*, **28**:327-336
- Pinheiro, E.F.M., Pereira M.G. and Anjos. L.H.C. (2004). Aggregate distribution and soil organic matter under different tillage systems for vegetable crops in a Red Latosol from Brazil. *Soil and Tillage Research*, **77**:79–84.
- Rohoskova, M. and Valla, M. (2004). Comparison of two methods for aggregate stability measurement-a review. *Plant, Soil Environment*, **50**:379-382.
- SAS. (2002) Proprietary Software Version 9.0. SAS Institute Inc, Cary, NC, USA.
- Six, J., Elliot, E.T., and Paustian. K. (2000). Soil macro aggregate turnover and micro aggregate formation: a mechanism for C sequestration under no tillage agriculture. *Soil and Biochemistry*, **32**:2099-2103.
- Sullivan, P. (2004). Sustainable Soil Management. ATTRA (National Sustainable Agricultural Information Service). Publication. #1P027/33, ATTRA, Fayetteville, USA. Access 3/4/2018.
- Tan, Z. X., Lal, R., Izaurrealde, R. C and Post, W. M. (2004). Biochemically protected soil organic carbon at the North Applachian experiment water shed. *Soil Science Society of American journal* **9**: 1-10.
- Unger, P.W. (1991). Overwinter changes in physical properties of no-tillage soil. *Soil Science Society of America Journal*, **55**:778-782pp.
- USDA- United States Department of Agriculture. (2014). Soil survey staff: Keys to soil taxonomy (12th ed...) Washington DC; USDA Natural Resources Conservation Service.
- Van Bavel, C.H.M. (1950). Mean weight diameter of soil aggregates as a statistical index of aggregation. *Soil Science of American Proceedings*, **14**:20-23.
- White, W.M. (1993). Dry aggregate distribution. In M.R. Carter (Ed.) “*Soil Sampling and Methods of Analysis*”. Lewis Publishers, Boca Raton, FL, 659-662pp.