

**KNOWLEDGE, PRACTICES AND PERCEPTION OF MALARIA AND ITS HOME
MANAGEMENT USING ARTEMICININ-BASED COMBINED THERAPY (ACT)
AMONG MOTHERS OF UNDER-FIVE**

ABSTRACT

Malaria is a serious public health problem, yet preventable and treatable. The disease is one of the world's highest rates of all cause of mortality for children under five, and about one in six children die before their fifth birthday. Hence, mothers of under-five and caregiver have a pivotal role to play in tackling this issue by improving their knowledge and skills concerning the treatment, prevention and control using appropriate approach. This study was carried out to assess knowledge, practices and perception of malaria and its home management using Artemisinin-based Combined Therapy (ACT) in Yemetu community of Ibadan North Local Government. The study was a descriptive cross-sectional survey involving the use of Expanded Programme on Immunization (EPI) to facilitate the sampling and interview of respondents. This included recruiting all the mothers of under-five in Yemetu community who gave consent for the study. Four hundred (400) mothers of under-five in Yemetu community consented to participate in the study and were selected. A validated semi-structured questionnaire interviewed and self-administered questionnaire was used for data collection and respondents were assessed on a 62-points knowledge scale, 5-points practice scale and 17-points perception scales. Knowledge score ≤ 21 were rated poor, scores $\geq 22 \leq 42$ fair and scores ≥ 43 were considered good. Practice score ≤ 3 was recorded poor practice while scores ≥ 3 good practice. Perception scores ≤ 9 were considered unfavorable perception and scores ≥ 9 were considered favorable. Descriptive statistics and chi square tests were used to analyse data at 95% level of significance.

Respondents' mean age was 29.9 ± 7.0 years and majority of them (91.1%) of them were Yoruba. Majority (91.0%) were married and (91.1%) were Yorubas. Only (23.0%) correctly identified plasmodium as a cause of malaria. The correctly mentioned signs and symptoms of simple malaria were; cold (89.3%), body ache (91.3%) and fever (88.5%). The fairly corrected home management practice steps include; Exposure of baby to fresh air, administration of paracetamol, and then provision of coartem(2.6%) and bathing the baby, use of paracetamol and administration of coartem (1.3%). Negative perception shown by the respondents include: Only (15.0%) believed that malaria is a disease of the poor and preference of herbal medicine to

34 medical medicine for treating children at home when they have malaria episode because it is
35 cheaper (19.8%). Overall, (2.8%) had poor knowledge, majority (87.3%) had fair knowledge and
36 9.3% had good knowledge. There are several gaps in the respondents' knowledge relating to
37 malaria and its management in under-five. Therefore, there is need for peer education/training
38 approach in this regard to upgrade mothers' knowledge and skills concerning the treatment,
39 prevention and control of malaria.

40

41 **Keywords:** Under-five, Home management of malaria, Artemisinin Combination Therapy
42 (ACT)

43 **1.0 INTRODUCTION**

44 Malaria remains one of the world's greatest childhood killers and is a substantial obstacle to
45 social and economic development in the tropics. It is a major cause of morbidity and mortality
46 especially among the vulnerable groups to which children, especially aged less than 5 years
47 belong. (Idro, Otieno, White, Kahindi, Fegan, Ogutu, Mithwani, Maitland, Neville, and Newton
48 C.R., 2007). It was observed that malaria accounts for 25 per cent of infant mortality and
49 30 percent of childhood mortality in Nigeria thereby imposing great burden on the country in
50 terms of pains and trauma suffered by its victims as well as loss in outputs and cost of treatments
51 (WHO, 2000).

52 The parasite responsible for these deaths—*Plasmodium falciparum*—is transmitted to people
53 when they are bitten (usually at night) by an infected mosquito. In the human body, the parasites
54 reproduce in the liver before invading red blood cells. Here, they multiply again before bursting
55 out and infecting more red blood cells as well as causing a high fever and sometimes damaging
56 vital organs. The transmission cycle is completed when a mosquito bites an infected person and
57 ingests parasites with its blood meal.

58 To reduce the global burden of malaria, this cycle needs to be broken. This can be done in
59 several ways. First, mosquitoes can be controlled with insecticides. Second, individuals can
60 avoid mosquito bites by sleeping under insecticide-treated nets. Finally, antimalarial drugs can
61 be used to reduce the illness and death caused by the malaria parasite and can lessen the
62 likelihood that a mosquito will pick up the parasite when it bites a person. (WHO, 2005).
63 Regardless of the fact that it is one of the oldest recorded diseases, malaria remains one of the
64 world's most deadly infectious diseases. It is arguably, the greatest menace to modern society in
65 terms of morbidity and mortality. Though preventable, treatable and curable, there is no known

66 immunity. Several centuries after its discovery, malaria still remains a devastating human
67 infection, resulting in 300-500 million clinical cases and three million deaths every year. (WHO,
68 2005).

69 It is also believed to contribute up to 11 percent maternal mortality, 25 percent infant mortality,
70 and 30 percent under-five mortality. It is estimated that about 132 billion Naira lost to malaria
71 annually in the form of treatment costs, prevention and loss of work time in Nigeria (FMOH and
72 NMCP, 2009)

73 Nigeria is known for high prevalence of malaria and it is a leading cause of morbidity and
74 mortality in the country. Available records show that at least 50 per cent of the population of
75 Nigeria suffers from at least one episode of malaria each year and this accounts for over 45 per
76 cent of all outpatient visits. (Ojurongbe, Ogunbamigbe, Fagbenro- Beyioku, Fendel, Kremsner,
77 and Kun 2007).

78 Malaria is known to have a negative impact on performance and learning in children according
79 to Holding and Snow (2001). It also aggravates anaemia and malnutrition in children and
80 pregnant women. (Murphy and Breman 2001).

81 There are strategies being promoted for the management of malaria as a result of the emergence
82 of chloroquine resistance aimed at preventing the occurrence of malaria. The World Health
83 Organization (WHO) currently recommends Artemisinin-based Combination Therapies (ACTs)
84 for malaria control. The use of insecticide-treated nets is also now being strongly promoted.
85 (WHO, 2006). Arigbabuwo, (2010) in his study also opined that prevention is better than cure,
86 advising that people should learn to maintain personal and environmental hygiene.

87 Mothers have a crucial role to play in recognition, treatment and prevention of malaria in under-
88 five. The knowledge, perception and skills and practices relating to malaria among mothers of
89 under-five in urban settings are yet to be well investigated. This study therefore focuses on
90 knowledge, practices and perception of malaria and its home management using Artemisinin
91 Combination Therapy (ACT) among mothers of under-five in Yemetu one of the communities in
92 Ibadan metropolis.

93 **2.0 Materials and Method**

94 The study was a descriptive cross-sectional survey involving the use of Expanded Programme on
95 Immunization (EPI) to facilitate the sampling and interview of respondents. This included

96 recruitment of all the mothers of under-five in Yemetu community who gave consent for the
97 study. Yemetu is a community located in Ibadan North Local Government Area (LGA) in ward
98 3. of Oyo State, Ibadan. South Western Nigeria. With ward number (3). It has a total population
99 of 65,949. Four hundred (400) mothers of under-five in Yemetu community however, consented
100 to participate in the study and were selected. The community had three (3) health facilities: A
101 government owned secondary health facility called Adeoyo Maternity Teaching Hospital, and
102 two private health facilities namely Kola Daisi Foundation Center (for primary and community
103 health) and Vine Branch Medical Clinic which are accessible to the people in the community.
104 The total number of under-five children is 13,190. (National Population Commission, 2006.) The
105 community is heterogeneous consisting of people from different part of the country. The
106 Yorubas however constituted the majority and their major occupation is trading.

107

108 **2.1 Target Population**

109 The populations for this study were mothers of under-five children residing in Yemetu
110 Community of Ibadan North LGA in Oyo State, South West of Nigeria who merited the
111 inclusion criteria for the study.

112

113 **2.2 Sampling Procedure**

114 The Expanded Programme on Immunization (EPI) sampling technique was used to facilitate the
115 sampling and interview of the respondents. The investigator started data collection by moving to
116 the center of Yemetu community and spinned a bottle. The spinned bottle was allowed to turn
117 round and round unhindered and allowed to come to rest. Interview started from the part of the
118 community to which the mouth of the bottle was pointing. Every third house in the direction was
119 selected and visited and one eligible respondent was selected by balloting for interview if more
120 than one eligible respondent was met in a house. In a house where there was one mother, such a
121 mother was purposively selected for interview if she consented to participate in the study.

122 After reaching the end of the community, the investigator and the research assistants moved back
123 to the center of the community and started recruitment and interview in another direction. This
124 way a total of 422 eligible mothers of under-five who consented to be involved in the study were
125 interviewed.

126 **2.3 Ethical Consideration**

127 All interviews was conducted in compliance with the ethics of the health promotion and
128 education profession. Copies of the research proposal were submitted to Oyo state Ethical
129 Review Committee for approval before the study commenced. This was done in order to ensure
130 that the study was conducted in accordance with ethical principle covering studies involving
131 human objects. The research assistants were well trained to obtain informed consent for
132 respondents before interview. Respondent were informed on the purpose of the study and were
133 given option to participate through written or verbal consent or withdraw from participating.
134 Informations provided by the respondent were treated with confidentiality. Registration number
135 was assigned to each questionnaire, no identifiers such as names, address or phone numbers was
136 required on the questionnaire.

137 **2.4 Data collection procedure:**

138 A semi-structured questionnaire was used for data collection. The semi-structured interviewer
139 and self-administered questionnaire was divided into six sections labeled sections A, B, C, D,
140 and E consisting of open ended and close ended questions. It was developed from literature
141 review and adapted questions from related past studies in Nigeria. The structured questionnaire
142 comprised of open and close ended questions which were used to elicit information on menace of
143 malaria disease, home management and practices and perception. The Yoruba version of the
144 questionnaire was produced after necessary modification to the English version had been done. It
145 consisted of 28 questions divided into five sections. The questionnaire was validated by the
146 researchers and experts in Public Health they included, pediatricians, statisticians working on
147 malaria control in university of Ibadan. It was pretested in Ekotedo community in Ibadan North
148 LGA a similar community in Ibadan North LGA. The data were then subjected to descriptive
149 statistics which was basically frequencies and charts. Reliability coefficient obtained was
150 determined using the Cronbach's Alpha technique. Any coefficient > 0.5 is said to be reliable. In
151 the study, reliability coefficient score which is also called chronbach Alpha was calculated to be
152 of 0.733.

153 **3.0 RESULTS**

154 **3.1 Socio-demographic demographic characteristics of respondents**

155 Table 1 presents the socio-demographic characteristics of the respondents. Respondents within
156 the age group 20-29 years constituted the highest (41.8%) followed by those aged 30-39 (40%).

157 Respondents' aged less than 20 years were the least (6.0%). The mean age of the respondents
158 was 29.9 ± 7.0 years.

159 Most of the respondents (90.8%) were married. Respondents with secondary school (56.5%)
160 topped the list of highest level of education. While those with tertiary accounted for (23.8%) of
161 the respondents.

162 Over half of the respondents were traders (57.8%). While artisans constituted (31.3%).
163 Respondents in monogamous constituted the majority (71.0%). (See table 1 for details)

164

165

166 **Table 1: Socio-demographic characteristics of the respondents**

167

Socio-demographic characteristics	Frequency	Percent (%)
Age in years: (n=397)*		
Less than 20 years	24	6.0
20-29 years	166	41.8
30-39 years	161	40.6
40-49 years	46	11.6
Marital status: (n=393)		
Single	29	7.3
Married	364	91.0
Religion: (n=400)		
Christianity	207	51.8
Islam	190	47.5
Traditional	3	0.75
Ethnic group: (n=395)		
Yoruba	360	91.1
Igbo	23	5.8
Hausa	12	3.0
Highest level of education: (n=400)		
Primary	79	19.8
Secondary	226	56.5
Tertiary	95	23.8
Type of tertiary education: (n=75)		
University	24	32.0
Polytechnic	39	52.0
Diploma/nursing	12	16.0
Occupation : (n=396)		
Trading	229	57.8
Civil servant	41	10.4
Artisan	124	31.3

Unemployed	2	0.4
Family type: (n=387)		
Polygyny	112	29.0
Monogamous	275	71.0
Children aged less than five: (n=397)		
One	262	66.0
Two	129	32.5
Three	6	1.5

168

169 *Mean age: 29.9± 7.0

170

171

172

173

174

175 **Table 2: Pattern of use and storage of antimalarial drugs and related medicine for treating**
 176 **under-fives**

Pattern of use of antimalarial	No	%
Ever used an antimalarial to treat under-five (N=398)		
Yes	357	89.7
No	41	10.3
Types of malarial medicine used(N=399)		
Coartem *	268	67.2
Artesunate*	65	16.3
Chloroquine -	28	7.0
Paracetamol -	18	4.5
Ampiclox -	13	3.3
Alabukun-	7	1.8
Pain relieving medicine normally used for treating underfive at home in case of malaria(N=399)		
Paracetamol*	322	80.7
Novagen-	47	11.8
Ibuprofen -	22	5.5
Alabukun -	8	2.0
Places where antimalarial medicines are kept at home(N=399)		
Cool dry place*	287	72.1
Inside nylon +	60	15.1

Inside wardrobe+	32	8.0
In the kitchen -	19	4.8

177 **Malaria treatment seeking pathways for under-five during episodes of malaria preceding**
178 **study.**

Where sought treatment	No	%
Hospital*	156	39.1
Patent Medicine Vendors (PMV) ±	120	30.1
Health centre*	50	12.5
Private clinic*	37	9.3
Primary Health Care (PHC)*	30	7.5
Community Medicine Distributors (CMDs)+	6	1.5

179 **Respondents step by step home management of malaria involving under-five child**

Steps taking at home	No	%
Use paracetamol for the baby+	177	46.7
Bath the baby±	52	13.7
Bath the baby and use PCM for the baby+	55	14.5
Use agbo for the baby±	44	11.6
Mop the body with cloth soaked in cold water±	11	2.9
Bath the baby, use paracetamol for him/her and take him/her to the hospital*	25	6.6
Expose to fresh air, give paracetamol and give coartem*	10	2.6
Bath for the baby, use paracetamol and give the baby coartem*	5	1.3

180
181 **3.2 Respondents' practices related to home management of malaria**
182 Respondents pattern of use and storage of antimalarial drugs and related medicine for treating
183 under-fives is highlighted in table 2. Majority (89.7%) of the respondents had used an
184 antimalarial drug to treat their under-five child (ren) while only 10.3% had never used
185 antimalarial drugs. The antimalarial drug normally used included; coartem (34.2%), artesunate
186 (29.0%), amalar (24.3%), while (10.5%) listed fansidar. Respondents were further asked about
187 type of malarial medicine used. Respondents that used coartem (67.2%) topped the list.
188 Respondents that used paracetamol for pain relieve in treating under-five at home had the highest
189 proportion of (80.7%) A majority (72.1%) kept their related malaria medicine in a cool dry
190 places. (See table 2 for details).

191 Table 2 highlighted malaria treatment seeking pathways for under-five during episodes of
192 malaria preceding the study. The highest proportion (39.1%) sought for treatment in a hospital,
193 followed by Patent Medicine Vendors (PMVs) (30.1%). The other listed places are contained in
194 the table under reference. Respondents' step by step home management of malaria involving
195 under-five children is specified in table 2. The fairly correct steps mentioned were as follows:
196 Exposure of baby to fresh air, administration of paracetamol, and then provision of

197 coartem(2.6%) and bathing the baby, use of paracetamol and administration of coartem (1.3%)
 198 (See table 2 for detail). Categorization of overall respondents' practice score was assessed using
 199 a 5-point scale. Respondents with good practice (4-5points) constituted 94.0%, while the
 200 proportion of respondents with poor practice (0-3) accounted for 6.0%.

201 **Table 3: Respondents' knowledge of causes of malaria**

Causes of malaria [^]	True (%)	False (%)	Don't know (%)	Total
Mosquito	392(98.0)	8(2.0)	0(0%)	400
Too much sun	221(55.8)	175(44.2)	4(1.0)	396
Change of weather	97(24.3)	281(70.3)	22(5.5)	400
Plasmodium	92(23.0)*	253(63.3)	55(13.8)	400
Taking too much palm oil	141(35.3)	230(57.5)	29(7.3)	400
Overwork/too much work	185(46.3)	199(49.8)	16(4.0)	400
Witchcraft	76(19.0)	292(73.2)	31(7.8)	399

202

203

204

205

206

207

208 **Table 4: Respondents' knowledge of factors or condition that can make mosquito breed or**
 209 **multiply**

Factors/conditions	True (%)	False (%)	Don't know (%)
Blocked gutters/drains with water	388(97.0)*	11(2.8)	1(0.3)
Improper refuse disposal	385(96.3)*	15(3.8)	0(0)
Stagnant water	340(85.0)*	58(4.5)	2(0.5)
Empty containers or vessels (e.g. bottles, cans, plastics etc.)	217(54.3)*	162(40.5)	21(5.3)
Engine oil in a container that is not covered+	120(30.0)	250(62.5)	30(7.5)
Stagnant water containing spent engine oil+	122(30.5)	246(61.5)	32(8.0)

210

211

212 **Table 5: Respondents' knowledge of signs and symptoms of simple malaria**

Signs and symptoms of simple malaria	Responses		
	Correct (%)	Wrong (%)	Don't know (%)
Inflammation of the skin	169(42.3)	175(43.8)	56(14.0)
Fever	354(88.5)*	38(9.5)	8(2.0)
Nausea+	347(86.8)	45(11.3)	8(2.0)
Diarrhoea	247(61.8)	136(34.0)	17(4.3)
Vomiting	320(80.0)*	71(17.8)	9(2.3)
Cold	357(89.3)*	36(9.0)	7(1.8)
Tiredness	358(89.5)*	34(8.5)	8(2.0)
Catarrh	365 (91.3)*	34(8.5)	1(.3)
Body ache	365(91.3)*	32(8.0)	3(.8)
Itching +	264(66.0)	125(31.3)	11(2.8)
Fatigue	310(77.5)*	75(18.8)	15(3.8)
Sore throat+	230(57.5)	148(37.0)	22(5.5)

213

214 **Table 6: Respondents' knowledge of signs and symptoms of severe malaria**

Symptoms of severe malaria^	Responses			Total
	Correct (%)	Wrong (%)	Don't know (%)	
Fever	361(90.5)*	25(6.3)	13(3.3)	399(100%)
Chills	352(88.0)*	31(7.8)	17(4.3)	400(100%)
Organs dysfunction	154(38.5)*	126(31.5)	120(30.0)	400(100%)
Abnormal bleeding	104(26.0)	167(41.8)	129(32.3)	400(100%)
Clinical jaundice	140(35.0)*	129(32.3)	131(32.8)	400(100%)
Febrile convulsion	175(43.8)*	107(26.8)	118(29.5)	400(100%)
Respiratory distress	130(32.5)*	119(29.8)	151(37.8)	400(100%)
Impaired consciousness	121(30.3)*	118(29.5)	161(40.3)	400(100%)

215

216 **Table 7: Respondents knowledge of preventive measures against malaria**

Preventive measures	True (%)	False (%)	Don't know (%)	Total
Using insecticide treated net	367(91.8)*	31(7.8)	2(.5)	400(100%)
Eating balanced diet	218(54.5)+	171(42.8)	9(2.3)	389(100%)
Clearing of residential environment of grasses/overgrown weeds	322(80.5)*	77(19.3)	1(.3)	400(100%)
Clearing blocked gutters	311(77.8)*	87(21.8)	2(.5)	400(100%)
Bathing daily	136(34.0)+	248(62.0)	15(3.8)	400(100%)
Use of insecticide	292(73.0)*	97(24.3)	11(2.8)	400(100%)
Use of antimalarial drug(SP) by pregnant women	199(49.8)*	175(43.8)	26(6.5)	400(100%)
Having enough sleep	87(21.8)+	283(70.8)	30(7.5)	400(100%)
Not eating too much palm oil	98(24.5)+	272(68.0)	30(7.5)	400(100%)
Not working in the sun for a long time	90(22.5)+	273(68.3)	37(9.3)	400(100%)

217

218

219

220 **Table 8: Respondents' knowledge of malaria related treatment actions involving under-five children treatment**

221

Treatment steps actions^	True (%)	False (%)	Don't know (%)	Total
Tepid sponging	363(92.2)*	25(6.3)	10(2.5)	398
Use of paracetamol	384(96.2)*	14(3.5)	1(0.3)	399
Use of coartem	363(91.0)*	22(5.5)	14(3.5)	399
Use of agbo	267(66.9)	125(31.3)*	7(1.8)	399
Use of chloroquine	256(64.2)	132(33.1)*	11(2.8)	399
Going to a health care facility for treatment	365(91.5)*	27(6.8)	7(1.8)	399

222


223

224 **Table 9: Respondents' knowledge of coartem and paracetamol dosage regimen for children**
 225 **aged 0-5years**

Ages(years)	Coartem dosage regimen [^]	Right	Wrong	Don't know	Total
6months-3years	1 tablet twice daily(3days)*	248(62.2)	93(23.3)	58(14.5)	399
	2 tablet twice daily(3days)	55(13.9)	280(70.7)	61(15.4)	396
	1 tablet thrice daily(3days)	33(8.3)	300(75.8)	63(15.9)	396
3-5years	1 tablet twice daily(3days)	100(25.3)	246(62.1)	50(12.6)	396
	2 tablet twice daily(3days)*	200(50.1)	150(37.6)	49(12.2)	399
	3 tablet thrice daily(3days)	55(13.9)	290(73.2)	51(12.9)	396
Paracetamol dosage regimen					
6months-3years	1/2 tablet twice daily(3days)*	256(64.0)	93(23.3)	51(12.8)	400
	1 tablet twice daily(3days)	74(18.6)	270(68.0)	53(13.4)	397
	1/2 tablet thrice daily(3days)	38(9.6)	308(77.6)	51(12.8)	397
3-5years	1 tablet twice daily(3days)	190(47.9)	169(42.6)	38(9.6)	397
	1 tablet thrice daily(3days)*	143(35.8)	219(54.8)	38(9.5)	400
	1 tablet once daily(3days)	48(12.1)	306(77.1)	42(10.6)	397

226

227 The knowledge of treatment/dosage regimen for coartem and paracetamol for children aged 0-5
 228 years is summarized in table 9. More than half of the respondents (62.2%) were knowledgeable
 229 about the correct treatment/dosage of coartem for children ages 6 months – 3 years. About half
 230 (50.0%) were conversant with the correct treatment/dosage regimen for coartem for children
 231 aged 3-5 years. Majority (64.0%) knew the correct treatment/dosage regimen of paracetamol for
 232 children aged 6 months – 3years while only (35.8%) were knowledgeable of the correct
 233 treatment/dosage regimen of paracetamol for children aged 3–5 years. (See table 9 for details).



234

235 Figure: 1 Categorization of respondents' knowledge scores relating to general knowledge of
 236 malaria and home management of malaria.

237 **Table 10: Respondents perception relating to vulnerability to seriousness and of treatment**
 238 **of malaria.**

Perception	Agree	Undecided	Disagree
Perception relating to vulnerability			
My child is not prone to malarial so no need of taking preventive measures	11(2.8)±	4(1.0)	385(96.3)+
I believe malaria is a disease of the poor, our child cannot get it because we are not poor	60(15.0) ±	3(0.8)	337(84.3)+
I make sure my child stays away from people or other children having malaria to avoid getting it.	55(13.8) ±	9(2.3)	336(84.0)+
I am of the opinion that a child that is well fed cannot have Malaria	50(12.5) ±	12(3.0)	338(84.5)+
Perception relating to seriousness			
I do not believe malaria is a serious disease for children	12(3.0) ±	4(1.0)	384(96.0)+
Malaria cannot lead to death of children aged less than five years	7(1.8) ±	9(2.3)	384(96.0)+
Perception relating to treatment			
Malaria infection is caused by witches and wizards, so telling me about using drugs to treat it is a waste of time	6(1.5) ±	28(7.0)	366(91.5)+
I believed malaria infection will disappear on its own without treatment/medicine	11(2.8) ±	12(3.0)	377(94.3)+

239

240

241 **Table 11: Respondents' perception relating to home management of malaria**

Perception	Agree	Undecided	Disagree
I believe chloroquine alone is enough to treat my child of any kind of malaria at home	20(5.0) ±	52(13.0)	328(82.0)+
I am of the opinion that coartem should be used at home only when the child's malaria is serious	27(6.8) ±	68(17.0)	305(76.3)+
Malaria infection in a child is best treated at home with chloroquine than Arthemisinin-based Combined Therapy (e.g coartem, artesunate etc)	24(6.0) ±	70(17.5)	306(76.5) +

The first dosage of malaria drug is enough to treat children when they have malaria	9(2.3) ±	47(11.8)	344(86.0) +
I prefer herbal medicine to medical medicine for treating my child at home when he/she has malaria because it is cheaper for treating under-five with malaria	79(19.8) ±	68(17.0)	253(63.3) +
Traditional medicine used at home is more effective for treating malaria in children aged less than five years	82(20.5) ±	67(16.8)	251(62.8) +
It is better to wait for a day or two to see whether an under-five has malaria before treating him/her at home with malaria medicine	105(26.3) ±	24(6.0)	271(67.8) +
Every mother should keep medicine at home for the home management of malaria when the need arises	301(75.3) ±	9(2.3)	90(22.5) +
It is wrong for a mother to treat her under-five children at home in case of malaria	133(33.3) ±	8(2.0)	259(64.8) _

242

243 3.3 Perception relating to malaria

244 Respondents in this study had good perception of home management of malaria. Majority of the
 245 respondent believed that ACT is best used in treating under-five children at home when they
 246 have malaria. A similar study was carried out by Ajayi, and Falade (2006); Salako, Brieger,
 247 Afolabi et al (2001) where respondents use chloroquine, and sulphadoxime/pyrimethamine (SP)
 248 at home for the treatment of malaria.

249

250 4.0 Conclusion

251 This study revealed that the level of awareness and knowledge of malaria among respondents
 252 was fair. However, there are several gaps in the respondents' knowledge relating to the disease
 253 and its management in under-five. The respondents had a poor knowledge of the cause and fair
 254 knowledge of the factors that could promote breeding of malaria. Advocacy, training and public
 255 enlightenment are necessary to address the situation.

256 5.0 Recommendations

257 The recommendations based on the findings of this study are as follow:

- 258 1. Sustained public enlightenment interventions relating to malaria targeted at mothers of under-
 259 five are needed. These intervention should be aimed at improving their knowledge and their
 260 malaria prevention and control skills.
- 261 2. Artemisinin Combination Therapy (ACT) is the new strategy for managing malaria. Training
 262 is needed to improve their knowledge and skill relating to the approach.

263 3. Formal health care facilities are commonly used by the residents for the management of
264 malaria in under-five. The capacity of health workers should be enhanced to help upgrade
265 mothers' knowledge and skills relating to correct treatment regimen for managing malaria.

266 4. Training on home management of malaria should be organized for respondent. A peer
267 education approach should be used in this regard to upgrade mothers' knowledge and skills
268 concerning the treatment, prevention and control of malaria.

269 6.0 REFERENCES

270 Ajayi IO. (2006): Development and Assessment of a treatment guideline to improve home
271 management of malaria in children in Ona-Ara local Government, Oyo state, Nigeria.
272 [Ph.D. dissertation]. Ibadan, Nigeria: University of Ibadan;

273 Arigbabuwo Adeleye (2010). Malaria: Killer at large. Vanguard Newspaper special report

274 Eckert E (Ed.); Scaling up home-based management of malaria – from research to
275 implementation. World Health Organisation, Geneva, 2004:2

276 Falade C.O., Ajayi, I.O., Yusuf, O.B., Pagnoni,F.(2014) 'High acceptance of artemisinin based
277 combination therapy for the home management of malaria in rural communities in
278 southwest Nigeria' *Malaria World Journal*;Vol. 5: 5

279 Falade C O, Ogundiran M O, Bolaji M O (2006): The influence of cultural perception of
280 causation, complication, and severity of childhood malaria on determinants and
281 treatment preventive pathways. *International Quarterly of Community Health
282 Education, a Journal of Policy and Applied Research* 2006. 24.4: 347 – 363.

283 FMOH (2009). Federal Republic of Nigeria training manual for management of malaria in
284 Nigeria Participants' Manual Federal Ministry of Health National Malaria and Vector
285 Control Division, Abuja- Nigeria

286 Holding P.A., Snow R.W., (2001). "Impact of Plasmodium falciparum malaria on performance
287 and learning: review of the evidence". *Am. J. Trop. Med. Hyg.* 64 (1-2 Suppl): 68–75.

288 Idro, R, Otieno G, White S, Kahindi A, Fegan G, Ogutu B, Mithwani S, Maitland K, Neville
289 BG, Newton CR. (2007). "Decorticate, decerebrate and opisthotonic posturing and
290 seizures in Kenyan children with cerebral malaria". *Malaria Journal* 4 (57). *Online
291 biomed Central.* 9

292 Murphy S.C., and Breman J.G. (2001). Gaps in the childhood malaria burden in Africa: cerebral
293 malaria, neurological sequelae, anemia, respiratory distress, hypoglycemia, and
294 complications of pregnancy. *Am. J. Tropical Med. and Hyg.* 64(1-2 Suppl):57-67

295 Ojurongbe O, Ogungbamigbe TO, Fagbenro- Beyioku AF, Fendel R, Kremsner PG, Kun JF
296 2007. Rapid detection of Pfcrt and Pfmdr1 mutations in Plasmodium falciparum isolates
297 by FRET and in vivo response to chloroquine among children from Osogbo, *Nigeria*.
298 *Malar J* 6: 41. 3

299 Salako, L.A., Brieger, W.R., Afolabi B.M. (2001). Treatment of childhood fever and other illness
300 in the three rural Nigerian communities. *J Trop pediatr*.

301 WHO (2000) The Abuja declaration and the plan of action. Geneva: World Health Organization.
302 WHO/CDS/RBM/2000.17. Available: [http://](http://www.rbm.who.int/docs/abuja_declaration.pdf)
303 www.rbm.who.int/docs/abuja_declaration.pdf. Accessed 10 March 2007.

304 WHO 2006 Calls for an immediate Halt to provision of single drug artemisinin Malaria Pills
305 WHO recommendations. 2005 , RBM Department; WHO Geneva