

Prevalence of childhood obesity among primary school students at Taif governorate, Saudi Arabia

Abstract

○ **Background:** Obesity in children is rapidly emerging as a global epidemic that will have profound public health consequences as overweight children become overweight adults particularly if obesity is present in adolescence.

○ **Objectives:** To determine the prevalence of overweight and obesity among school children between 6-12 years (males and females) at Taif primary schools and determine the associated dietary risk factors for overweight and obesity.

○ **Methods:** A cross sectional analytic study was implemented included a representative sample of primary school students (boys and girls) enrolled in governmental schools in Taif, Saudi Arabia, 2014-2015. Data were collected by using valid self-administered questionnaire, with family help. It including socio-demographic characteristics and detailed dietary history. Body mass index-for-age percentiles of CDC were utilized in this study to diagnose obesity.

○ **Results:** Among schoolboys, the prevalence of overweight was 10.7%, while prevalence of obesity was 7.6%; the problem seemed to be worse in girls, where the prevalence of overweight and obesity were 16.8% and 18.2%, respectively. Multivariate logistic regression analysis revealed that girls were almost double risk for developing overweight/obesity compared to boys ((aOR=1.93; 95%CI: 1.03-3.60, $p<0.05$). Habit of eating cheeps potatoes at school at abnormal rate was associated with almost three-folded risk for obesity or overweight compared to eating cheeps potatoes within normal (aOR=3.14; 95%CI: 1.17-8.41). School children who had the habit of taking fast foods at least three times/week were more likely to develop obesity or overweight compared to those who did not take fast foods (aOR=1.98; 95%CI:1.08-6.30).

○ **Conclusions:** Overweight and obesity among primary school children in Taif city, Saudi Arabia is a public health problem affecting more than one-third of girls and almost one fifth of boys. It confirms the significant association between some unhealthy dietary habits and increase prevalence of childhood obesity.

Keywords: Obesity, Saudi Arabia, childhood, dietary habits

INTRODUCTION

Obesity is one of the most common disorders encountered in clinical practice and has major public health implications. Unfortunately, it is also one of the most difficult and frustrating disorders to manage successfully.¹

Obesity is defined as the presence of excess adipose tissue.¹ It is a complex condition, with serious social and psychological dimensions^{2, 3} that affect virtually all age and socioeconomic groups and threatens to overwhelm both developed and developing countries.⁴

Mortality rises exponentially with increasing bodyweight. The risk of coronary heart disease is doubled if the body mass index (BMI) is >25 and nearly quadrupled if the index is >29 .^{5,6} The risk of developing diabetes increases with increasing weight and people with a body mass index >35 have a 40 fold higher risk of developing the disease than non-obese people.^{5,7} Osteoarthritis and respiratory diseases, particularly sleep apnoea are more common in obese people.⁵

Obesity was significantly associated with an increase in both systolic and diastolic blood pressure,^{8, 9} stroke, and certain forms of cancer.⁴ The prevalence of overweight and obesity has increased in the last few years.¹⁰ Between 1980 and 1995, the prevalence of obesity in Britain doubled from 8-15%.⁵ In 1995, there were an estimated 200 million obese adults worldwide and another 18 million children under-5-years classified as overweight. By the year 2000, the number of obese adults had increased to over 300 million.⁴ Children and adolescents are also involved in those changes.

In the United States of America (USA), the percentage of children and adolescents who are overweight has more than doubled in the past 30 years.¹¹ Although obesity in children is rarely associated with morbidity or mortality, it is rapidly emerging as a global epidemic that will have profound public health consequences as overweight children become overweight adults particularly if obesity is present in adolescence.¹² For example, the risk of developing adult obesity in children aged >9 years who are obese is up to 80% at age 35 years.¹³ Unfortunately, it is evident that obesity is a common health problem among Saudis.¹⁴

Overweight and obesity in the adult Saudi population were reported in different studies with a range of males overweight 26-34%, obesity 12-23% and females overweight 24-29%, obesity 19-41%.¹⁴⁻¹⁹ This high prevalence of overweight and obesity is a cause of concern, as obesity is associated with several complications that increase both morbidity and mortality.

Rationale:

- 1- Up to our knowledge, there is no previous study to assess prevalence of childhood obesity among primary school students in Taif.
- 2- Childhood obesity is a worldwide as well a Saudi health problem and need a lot of actions to face it.
- 3- It is a preventable problem through simple procedure such as parental education and counseling.

Aim:

To assess the magnitude of the problem of childhood obesity among primary school students in Taif governorate, 2014-2015.

Objectives:

- To determine the prevalence of overweight and obesity among school children between 6-12 years at Taif primary schools.
- To compare the prevalence between male and female students.
- To determine the associated dietary risk factors for overweight and obesity.

2. LITERATURE REVIEW

While reviewing the literatures about childhood obesity prevalence, a lot of articles worldwide were cited. In Saudi Arabia, there were 6 articles published in different local and international journals.

2.1 Local studies

-Overweight and obesity among Saudi Arabian children and adolescent between 1994 and 2000 by Abalkhil and he observed that the combined prevalence of overweight and obesity in the country was estimated to be around 27.5% among boys between 6 and 18 years of age in 1996, and 28 % among girls between 12 and 19 years in 1999, in comparison between 1994 and 2000, there is total increase in BMI for boys 10-12 % (up to age 16) followed by gradual decline with age , and for girls 2.4-4.9 % (up to age 15) followed by gradual increase in BMI reaching 9 % at age 20.²⁰

-Another study conducted by Al Herbish to identify the etiology of childhood obesity. The researcher found that the nutritional factor is the most important factor representing 88.5 % of cases.²¹

-El-Hazmi and Warsy studied the prevalence of obesity and overweight in 1-18 year old Saudi children. The result revealed that the overall prevalence of overweight was 10.7 % for boys and 12.7% for girls, while obesity was 6% in boys and 6.74% in girls. The highest frequency was in the eastern region, while the lowest was in southern region.²²

-Obesity among Saudi male adolescents in Riyadh was studied by Al-Rukban. He found that the prevalence of overweight was 13.8 % and obesity was 20.5 %.²³

- khaled, et al. conducted a study to describe the prevalence of childhood overweight and obesity in rural high- and low-altitude populations of southwestern Saudi Arabia and to identify specific at-risk groups within these populations and they found that the overall prevalence of overweight and obesity is 10%. Among risk factors of overweight and obesity, high altitude was a significant and independent factor.²⁴

-Recently, Al-Shehri carried out a cross-sectional study to determine the prevalence of overweight and obesity among school children between 6-12 years as well as to identify the associated unhealthy dietary habits for overweight and obesity. It included a representative sample of primary schoolboys in Al-Iskan sector in Makkah AlMukarramah. The sample size was distributed among the three primary school in Al-Iskan sector and determined as a percentage proportionally related to the total number of the student in each school. Self-administered questionnaire (with family help) including socio-demographic characteristic and detailed dietary history, was used for data collection. Trained staff collected

anthropometric measurements of weight and height. The study included 258 male primary school children. Their age ranged between 6 and 12 years. The prevalence of overweight was 12.4% and that of obesity was 20.2%. Among studied socio-demographic variables associated with overweight and obesity, birth weight above normal was significantly associated with them ($p < 0.05$) while among studied dietary factors, eating fried potatoes and chocolate at a rate more than normal was significantly associated with overweight and obesity among male school children. The habit of eating fast foods frequently (at least three times/week) was accompanied with higher prevalence of obesity (30.4%) compared to only 8% among school children who did not eat fast foods. This association between BMI and habit of eating fast foods was statistically significant ($p < 0.05$). He concluded that the study provided alarming evidence-based data on the considerable prevalence of childhood overweight and obesity among primary school children in Holy Makkah, Saudi Arabia.²⁵

2.2 Regional studies

-In gulf countries, 3 related studies conducted in Kuwait, UAE, and Iraq were cited. The Iraqi one reported the prevalence and possible risk factors of childhood obesity and revealed that the prevalence of overweight was 6 % and of obesity was 1.3%.²⁶

-In UAE, A national study was carried out by Al-Haaddad who conclude that the frequency of obesity among UAE youth was two to three times greater than the recently published international standard.²⁷

-In Kuwait, prevalence of obesity among adolescents (10-14 years) in Kuwait was studied by Sorkhou. He found that the overall prevalence of overweight among adolescent Kuwaiti children was 30.7% and for obesity was 14.6 %.²⁸

-In Egypt, Badawi et al conducted a study to estimate the prevalence of overweight and obesity among primary school children, aged from 6 to 12 years and to estimate risk factors of obesity and overweight, defined by body mass index (BMI). A cross-sectional study was carried out at Port Said city during the second term of school year 2010/2011. Eight hundred and fifty-two students participated in this study. Prevalence of overweight and obesity was 17.7% and 13.5% respectively. The rate of obesity was the highest at the age of 7–8 years (grade 2) and decreased with an increase in age, while overweight increased with an increase in age to be the highest at the age of 9–10 (grade 4) and 10–11 (grade 5). Socioeconomic class, faulty dietary habits, sedentary life, low level of physical activity and positive family history of overweight and/or obesity were significantly associated with student's BMI.

2.3 International studies

-In USA, changes in state-specific childhood obesity and overweight prevalence in the United States from 2003 to 2007 was studied by Singh et al. The results revealed that in 2007, the prevalence of obesity was 16.4% and overweight was 31.6% and there is 10% increase in prevalence between 2003 and 2007.³⁰

-In Japan, prevalence of childhood obesity from 1978 to 2007 was studied by Yoshinaga et al. The result showed gradual decrease in the prevalence of obesity since the early 2000s.³¹

-In Tanzania, Muhihi et al reported an overall prevalence of child obesity of 5.2% and was higher among girls (6.3%) compared to boys (3.8%). Most obese children were from households with fewer children ($p=0.019$) and residing in urban areas ($p=0.002$).

Controlling for other variables, age above 10 years (aOR=3.3, 95% CI=1.5-7.2), female sex (aOR=2.6, 95% CI=1.4-4.9), urban residence (aOR=2.5, 95% CI=1.2-5.3) and having money to spend at school (aOR=2.6, 95% CI=1.4-4.8) were significantly associated with child obesity.³²

3. METHODOLOGY

3.1 The study area:

Taif city is located in the Mecca Province of Saudi Arabia at the West of Saudi Arabia in an elevation of 1700 meters on the slopes of the Al-Sarawat Mountains. It has a population of 987,914 (2010 census). Taif city includes 605 primary governmental schools (day shift), 310 for boys and 295 for girls. The number of primary schoolchildren is 66294 (31754 boys and 34,540 girls)

3.2 -Study design:

A cross sectional analytic study

3.3 Study population:

Primary school students (boys and girls) enrolled in governmental schools in Taif, Saudi Arabia, 2014-2015, who were present at the time of the study and were willing to participate in it.

3.4 Population selection criteria:

Inclusion criteria:

- Students enrolled in governmental primary schools in Taif, Saudi Arabia, 2014-2015
- Boys and girls

Exclusion criteria:

- Primary school students who were not present at the time of conducting the study or not willing to participate.
- Those aged over 12 years.

3.5 Sample size:

The sample size was calculated by Raosoft online calculator.³³ It was 247 students (based on 20.2% expected prevalence).²⁵ The worst accepted prevalence was 22% and confidence interval of 95%. This sample will be increased to 275 students to compensate for non-response (132 boys and 143 girls).

3.6 Sampling Technique:

The total number of students in the 237 schools is 66294 students and the sample size is 275 (0.37%). Two female and two male primary schools were randomly selected in Taif city. All students from all levels (one to six) in the selected schools were eligible for inclusion in the study. The sample size was distributed among the four randomly selected primary schools and determined as a percentage proportionally related to the total number of the student in each school. Systematic random sampling technique was applied to select the sample size from each school. A sufficient sample size was recruited from these schools.

3.7 Data collection:

Data were collected by using self-administered questionnaire, with family help. It has been previously validated and proved to be reliable in a study conducted in Makkah by Al-Shehri.²⁵ It including socio-demographic characteristic and detailed dietary history. Trained staff collected anthropometric measurements of weight and height. Height was measured without shoes to the nearest “0.5cm” and weight to the nearest “100g with the subject in light clothes and without shoes. A single scale was used for weighing all the students. This scale was calibrated daily, and zero was assured before weighing any student.

Body mass index which is the weight in kilogram divided by the height in meters squared (kg/m^2), was calculated for all the study participants.

3.8 Outcome variables:

Under weight; when BMI < 5th percentile, Overweight; when BMI > or = 85th percentile and < 95th percentile and Obesity; when BMI > or = 95th percentile. Body mass index-for-age percentiles of CDC were utilized in this regard.³⁴

3.9 Preparatory phase

Intensive literature review has been made using internet midline search and journals for both national and international related studies were obtained.

3.10 Pilot study

Pilot study was conducted before data collection and no modifications were needed on the pilot testing results. Randomly selected 12 female and 12 male students in two schools (2 from each level) were selected for the pilot testing. Data of the pilot study were included in the actual study since there was no significant variations from the main survey results.

3.11 Data entry and analysis:

Statistical Package for Social Sciences (SPSS) software version 20.0 was used for data entry and analysis. Descriptive statistics (e.g. number, percentage) and analytic statistics using Chi Square tests (χ^2) to test for the association and/or the difference between two categorical variables was applied. P-value equal or less than 0.05 was considered statistically significant. Multivariate regression analysis was used to adjust for confounding factors for obesity and overweight.

3.12 Ethical consideration

- Permission from Joint Program of Family Medicine in Taif was obtained.
- Permission of the education director in Taif was obtained.
- All information were kept confidential and results will be submitted to the Ministry of Education in Taif as feedback.
- We tried to follow the obese boys and girls and refer them to primary care centers.

4. RESULTS:

The study included 275 students enrolled in Taif governmental primary schools (143 girls and 132 males). Table 1 presents their socio-demographics. Their age ranged between 6 and 12 years with a mean of 9.44 years and standard deviation of ± 1.89 years. Birth weight was normal among majority of them (87.3%) while it was above normal among only 3.6% of them. Majority of them (96.4%) live with both parents. Most of them (81.1%) had family size ranged of 8 persons or less. More than two-thirds of them (69.1%) had two brothers or less whereas slightly more than half of them (52.4%) had two sisters or less. Almost a fifth of them (21.8%) were first birth order. More than one third of fathers (36.7%) and mothers (37.4%) were at least university graduated.

Table (1): Socio-demographic characteristics of the primary schoolchildren, Taif (n=275)

| Socio-demographic characteristics | Number | percentage |
|-----------------------------------|--------|------------|
| Age in years | | |
| 6-9 | 140 | 50.9 |
| >9-12 | 135 | 49.1 |
| Gender | | |
| Males | 132 | 48.0 |
| Females | 143 | 52.0 |
| School grade | | |
| 1 st | 32 | 11.6 |
| 2 nd | 48 | 17.5 |
| 3 rd | 33 | 12.0 |
| 4 th | 54 | 19.6 |
| 5 th | 46 | 16.7 |
| 6 th | 62 | 22.5 |
| Birth weight | | |
| Less than normal | 25 | 9.1 |
| Normal | 240 | 87.3 |
| More than normal | 10 | 3.6 |
| Living status | | |
| With both parents | 265 | 96.4 |
| Either father or mother | 10 | 3.6 |

| | | |
|---------------------------|-----|------|
| Family size | | |
| ≤8 | 223 | 81.1 |
| >8 | 52 | 18.9 |
| Number of brothers | | |
| ≤2 | 190 | 69.1 |
| >2 | 85 | 30.9 |
| Number of sisters | | |
| ≤2 | 144 | 52.4 |
| >2 | 131 | 47.6 |
| Birth order | | |
| First | 60 | 21.8 |
| 2-5 | 167 | 60.7 |
| >5 | 48 | 17.5 |

Table (2): Socio-demographic characteristics of the primary schoolchildren, Taif (n=275) (Cont.)

| Socio-demographic characteristics | Number | percentage |
|--|---------------|-------------------|
| Father`s educational level | | |
| Illiterate | 9 | 3.3 |
| primary | 18 | 6.5 |
| Intermediate | 31 | 11.3 |
| Secondary | 116 | 42.2 |
| University | 84 | 30.5 |
| Postgraduate | 17 | 6.2 |
| Mother`s educational level | | |
| Illiterate | 17 | 6.2 |
| primary | 28 | 10.2 |
| Intermediate | 29 | 10.5 |
| Secondary | 98 | 35.7 |
| University | 96 | 34.9 |
| Postgraduate | 7 | 2.5 |

Prevalence of overweight and obesity:

As shown in figure (1), overweight and obesity were reported among 16.8% and 18.2% of girls, respectively compared to 10.6% and 7.6% of boys, respectively. The

difference was statistically significant, $p < 0.001$ (Table 2). Among girls, the highest rate of obesity was observed in grade 3 (40%) whereas the highest rate of overweight was observed in grade 6 (35.3%). Among boys, the highest rate of obesity and overweight were reported in grade 3 (25% for each). The difference between boys and girls in this regard was statistically significant, $p < 0.001$. Figure 2




Figure 1: Distribution of Body Mass Index among primary school children in Taif according to gender.




Figure 2: Prevalence of overweight and obesity among primary school children in Taif according to grades and gender.

Factors associated with overweight and obesity:

Socio-demographic characteristics:

Table (2) summarizes the distribution of body mass index among primary schoolchildren according to their socio-demographic characteristics. As previously demonstrated girls had more significant overweight and obesity rates compared to boys. Regarding school grade, the highest rate of overweight was reported among children of grade 6 (27.4%) whereas the highest rate of obesity was reported among those of grade three (36.4%). On the other hand, the lowest rate of overweight was observed among children of grade 5 (4.3%) while the lowest rate of obesity was reported among children of grade one (6.3%). This association between school grade and overweight and obesity was statistically significant, $p < 0.001$.

Obesity was more significantly reported among students who had more than two sisters compared to those who had two sisters or less (19.8% versus 6.9%). This difference is statistically significant, $p=0.015$.

There were no statistically significant association between BMI from one side and other studied socio-demographic characteristics such as age, birth weight, living status, family size, number of brothers, birth order and parental education from the other side.

Table (3): Body mass index of primary schoolchildren according to their socio-demographic characteristics

| Socio-demographic characteristics | Body Mass Index | | | | χ^2 -value (p-value) |
|-----------------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Age in years | | | | | |
| 6-9 (n=140) | 14 (10.0) | 89 (63.6) | 17 (12.1) | 20 (14.3) | 4.03 (0.259) |
| >9-12 (n=135) | 6 (4.4) | 92 (68.1) | 21 (15.6) | 16 (11.9) | |
| Gender | | | | | |
| Males (n=132) | 4 (3.0) | 104 (78.8) | 14 (10.6) | 10 (7.6) | 20.56 (<0.001) |
| Females (n=143) | 16 (11.2) | 77 (53.8) | 24 (16.8) | 26 (18.2) | |
| School grade | | | | | |
| 1 st (n=32) | 4 (12.5) | 23 (71.9) | 3 (9.4) | 2 (6.3) | 43.22 (<0.001) |
| 2 nd (n=48) | 6 (12.5) | 29 (60.4) | 8 (16.7) | 5 (10.4) | |
| 3 rd (n=33) | 4 (12.1) | 14 (42.4) | 3 (9.1) | 12 (36.4) | |
| 4 th (n=54) | 2 (3.7) | 43 (79.6) | 5 (9.3) | 4 (7.4) | |
| 5 th (n=46) | 2 (4.3) | 36 (78.3) | 2 (4.3) | 6 (13.0) | |
| 6 th (n=62) | 2 (3.2) | 36 (58.1) | 17 (27.4) | 7 (11.3) | |
| Birth weight | | | | | |
| < normal (n=25) | 2 (8.0) | 16 (64.0) | 5 (20.0) | 2 (8.0) | 8.11 (0.230) |
| Normal (n=240) | 18 (7.5) | 160 (66.7) | 32 (13.3) | 30 (12.5) | |
| > normal (n=10) | 0 (0.0) | 5 (50.0) | 1 (10.0) | 4 (40.0) | |
| Living status | | | | | |
| With both parents (265) | 20 (7.5) | 173 (65.3) | 37 (14.0) | 35 (13.2) | 1.26 (0.739) |
| Either one of them (10) | 0 (0.0) | 8 (80.0) | 1 (10.0) | 1 (10.0) | |
| Family size | | | | | |
| ≤8 (n=223) | 13 (5.8) | 152 (68.2) | 27 (12.1) | 31 (13.9) | 7.45 (0.059) |
| >8 (n=52) | 7 (13.5) | 29 (55.8) | 11 (21.2) | 5 (9.6) | |
| Number of brothers | | | | | |

| | | | | | |
|---------------------------|----------|------------|-----------|-----------|------------------|
| ≤2 (n=190) | 17 (8.9) | 123 (64.7) | 25 (13.2) | 25 (13.2) | 2.68 (0.444) |
| >2 (n=85) | 3 (3.5) | 58 (68.2) | 13 (15.3) | 11 (12.9) | |
| Number of sisters | | | | | |
| ≤2 (n=144) | 10 (6.9) | 103 (71.5) | 21 (14.6) | 10 (6.9) | 10.39 (0.015) |
| >2 (n=131) | 10 (7.6) | 78 (59.5) | 17 (13.0) | 26 (19.8) | |
| Birth order | | | | | |
| First (n=60) | 3 (5.0) | 44 (73.3) | 7 (11.7) | 6 (10.0) | 4.57 (0.600) |
| 2-5 (n=167) | 13 (7.8) | 108 (64.7) | 21 (12.6) | 25 (15.0) | |
| >5 (n=48) | 4 (8.3) | 29 (60.4) | 10 (20.8) | 5 (10.4) | |
| Father`s education | | | | | |
| Illiterate (n=9) | 2 (22.2) | 5 (55.6) | 0 (0.0) | 2 (22.2) | 20.08 (0.169) |
| Primary (n=18) | 0 (0.0) | 15 (83.3) | 2 (11.1) | 1 (5.6) | |
| Intermediate (n=31) | 4 (12.9) | 16 (51.6) | 7 (22.6) | 4 (12.9) | |
| Secondary (n=116) | 11 (9.5) | 74 (63.8) | 18 (15.5) | 13 (11.2) | |
| University (n=84) | 1 (1.2) | 59 (70.2) | 10 (11.9) | 14 (16.7) | |
| Postgraduate (n=17) | 2 (11.8) | 12 (70.6) | 1 (5.9) | 2 (11.8) | |

Table (4): Body mass index of primary schoolchildren according to their socio-demographic characteristics (Cont.)

| Socio-demographic characteristics | Body Mass Index | | | | χ^2 -value (p-value) |
|-----------------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Mother`s education | | | | | |
| Illiterate (n=17) | 2 (11.8) | 10 (58.8) | 2 (11.8) | 3 (17.6) | 14.65 (0.477) |
| Primary (n=28) | 1 (3.6) | 19 (67.9) | 7 (25.0) | 1 (3.6) | |
| Intermediate (n=29) | 4 (13.8) | 15 (51.7) | 6 (20.7) | 4 (13.8) | |
| Secondary (n=98) | 5 (5.1) | 66 (67.3) | 14 (14.3) | 13 (13.3) | |
| University (n=96) | 8 (8.3) | 66 (68.8) | 9 (9.4) | 13 (13.5) | |
| Postgraduate (n=7) | 0 (0.0) | 5 (71.4) | 0 (0.0) | 2 (28.6) | |

Dietary factors:

-History of having breakfast: As seen in table (3), there was no statistically significant association between history of having breakfast and obesity ($p>0.05$).

Table (5): Association between body mass index of primary school children and history of taking breakfast

| Taking breakfast | Body Mass Index | | | | χ^2 -value (p-value) |
|-------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Always (n=117) | 9 (7.7) | 77 (65.8) | 15 (12.8) | 16 (13.7) | 12.55 (0.184) |
| Often (n=51) | 4 (7.8) | 29 (56.9) | 8 (15.7) | 10 (19.6) | |
| Sometimes (n=61) | 1 (1.6) | 49 (80.3) | 7 (11.5) | 4 (6.6) | |
| No (n=46) | 6 (13.0) | 26 (56.5) | 8 (17.4) | 6 (13.0) | |

History of having food at schools:

Table (4) demonstrates that obesity was reported among 13.7% and 15.8% of students who take food always or sometimes, respectively at schools compared to zero among those who had no food at schools. The association was statistically significant, $p < 0.05$.

Among studied food stuffs consumed by school children at schools, intake of milk, cheeps potatoes and cake were significantly associated with higher prevalence of obesity and/or overweight. Having abnormal large amounts of milk was associated with higher significant prevalence of overweight (19.8% versus 10.9%), $p < 0.05$. Also, having abnormal large amounts of cake at schools was associated with higher significant prevalence of obesity (20.5% versus 10.2%), $p = 0.05$. The intake of above normal amount of cheeps potatoes was associated with higher significant prevalence of obesity (32.6% versus 9.5%), $p < 0.001$. The association between intake of other food stuffs (strawberry juice,

beverages, fried potatoes, chocolate, biscuits and fruits) and obesity was not statistically significant.

Table (6): Association between body mass index of primary schoolchildren and their history of having food at schools

| | Body Mass Index | | | | χ^2 -value (p-value) |
|------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|-----------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Food at schools | | | | | |
| Always (n=182) | 10 (5.5) | 128 (70.3) | 19 (10.4) | 25 (13.7) | 17.96 (0.036) |
| Often (n=32) | 4 (12.5) | 22 (68.8) | 4 (12.5) | 2 (6.3) | |
| Sometimes (n=57) | 6 (10.5) | 27 (47.4) | 15 (26.3) | 9 (15.8) | |
| No (n=4) | 0 (0.0) | 4 (100) | 0 (0.0) | 0 (0.0) | |
| Milk | | | | | |
| Within normal (184) | 18 (9.8) | 122 (66.3) | 20 (10.9) | 24 (13.0) | 8.34 (0.040) |
| Abnormal (91) | 2 (2.2) | 59 (64.8) | 18 (19.8) | 12 (13.2) | |
| Sweaty juice | | | | | |
| Within normal (150) | 10 (6.6) | 108 (72.0) | 16 (10.7) | 16 (10.7) | 5.94 (0.115) |
| Abnormal (125) | 10 (8.0) | 73 (58.4) | 22 (17.6) | 20 (16.0) | |
| Beverages | | | | | |
| Within normal (109) | 10 (9.2) | 66 (60.6) | 17 (15.6) | 16 (14.7) | 2.42 (0.490) |
| Abnormal (166) | 10 (6.0) | 115 (69.3) | 21 (12.7) | 20 (12.0) | |
| Cheeps potatoes | | | | | |
| Within normal (232) | 20 (8.6) | 155 (66.8) | 35 (15.1) | 22 (9.5) | 20.4 (<0.001) |
| Abnormal (43) | 0 (0.0) | 26 (60.5) | 3 (7.0) | 14 (32.6) | |
| Chocolate | | | | | |
| Within normal (228) | 17 (7.5) | 152 (66.7) | 32 (14.0) | 27 (11.8) | 1.84 (0.606) |
| Abnormal (47) | 3 (6.4) | 29 (61.7) | 6 (12.8) | 9 (19.1) | |
| Cake | | | | | |
| Within normal (197) | 12 (6.1) | 138 (70.1) | 27 (13.7) | 20 (10.2) | 7.81 (0.050) |
| Abnormal (78) | 8 (10.3) | 43 (55.1) | 11 (14.1) | 16 (20.5) | |
| Biscuits | | | | | |
| Within normal (202) | 13 (6.4) | 140 (69.3) | 24 (11.9) | 25 (12.4) | 4.50 (0.212) |
| Abnormal (73) | 7 (9.6) | 41 (56.2) | 14 (19.2) | 11 (15.1) | |
| Fried Potatoes | | | | | |
| Within normal (214) | 17 (7.9) | 140 (65.4) | 28 (13.1) | 29 (13.6) | 0.74 (0.863) |
| Abnormal (61) | 3 (4.9) | 41 (67.2) | 10 (16.4) | 7 (11.5) | |
| Fruits | | | | | |
| Within normal (200) | 14 (7.0) | 132 (66.0) | 26 (13.0) | 28 (14.0) | 0.90 (0.826) |
| Abnormal (75) | 6 (8.0) | 49 (65.3) | 12 (16.0) | 8 (10.7) | |

-Most frequent meal:

As illustrated in table (5), among primary schoolchildren, the type of frequent taken main meal was not significantly associated with prevalence of obesity, $p>0.05$.

Table (7): Association between body mass index of primary schoolchildren and their most frequent meal

| | Body Mass Index | | | | χ^2 -value (p-value) |
|--|--------------------------------|----------------------------|-------------------------------|--------------------------|------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Breakfast alone or with others (20) | 2 (10) | 9 (45.0) | 5 (25.0) | 4 (20.0) | 7.06 (0.316) |
| Lunch and/or dinner (100) | 7 (7.0) | 65 (65.0) | 17 (17.0) | 11 (11.0) | |
| All meals (153) | 9 (5.9) | 107 (69.9) | 16 (10.5) | 21 (13.7) | |

* Data were missed in two cases

-Habit of eating between meals:

As obvious from table (6), habit of eating between meals was not significantly associated with prevalence of obesity among primary schoolchildren ($p>0.05$).

Table (8): Association between body mass index of primary schoolchildren and habit of eating between meals

| | Body Mass Index | | | | χ^2 -value (p-value) |
|--------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Always (n=44) | 7 (15.9) | 28 (63.6) | 2 (4.5) | 7 (15.9) | |
| Often (n=90) | 4 (4.4) | 64 (71.1) | 11 (12.2) | 11 (12.2) | |
| Sometimes (n=123) | 7 (5.7) | 78 (63.4) | 23 (18.7) | 15 (12.2) | |

| | | | | | |
|-----------------|----------|-----------|----------|----------|-------------------------|
| No (=18) | 2 (11.1) | 11 (61.1) | 2 (11.1) | 3 (16.7) | 12.56 (0.183) |
|-----------------|----------|-----------|----------|----------|-------------------------|

-Frequency of eating Fruits and vegetables:

As displayed from table (7), frequencies of eating vegetables and fruits were not statistically significantly associated with obesity.

Table (9): Association between body mass index of primary school children and frequency of eating fruits and vegetables

| consume of vegetables and fruits (times/day) | Body Mass Index | | | | χ^2 (p-value) |
|---|---|-------------------------------------|--|-----------------------------------|--------------------------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Vegetables | | | | | |
| ≤ 2/ day (236) | 18 (7.6) | 156 (66.2) | 31 (13.1) | 31 (13.1) | 0.87 (0.832) |
| ≥ 3/ day (39) | 2 (5.1) | 25 (64.1) | 7 (18.0) | 5 (12.8) | |
| Fruits | | | | | |
| ≤ 2/ day (240) | 19 (7.9) | 158 (65.8) | 32 (13.3) | 31 (12.9) | 1.44 (0.696) |
| ≥ 3/ day (35) | 1 (2.9) | 23 (65.7) | 6 (17.1) | 5 (14.3) | |

-Habit of eating fast foods:

Table (8) shows that the habit of eating fast foods frequently (more than three times/week) was accompanied with higher prevalence of obesity (26.7%) compared to none among schoolchildren who did not take fast foods. This association between BMI and habit of eating fast foods was statistically significant ($p=0.036$).

Table (10): Association between body mass index of primary schoolchildren and their habit of eating fast foods

| Eating fast food | Body Mass Index | | | | χ^2 (p-value) |
|--------------------------------|--------------------------------|----------------------------|-------------------------------|--------------------------|--------------------|
| | Underweight N=20 No. (%) | Normal N=181 No. (%) | Overweight N=38 No. (%) | Obese N=36 No. (%) | |
| Never (n=7) | 2 (28.6) | 4 (57.1) | 1 (14.3) | 0 (0.0) | 22.14 (0.036) |
| Once/week (n=159) | 15 (9.4) | 108 (67.9) | 17 (10.7) | 19 (11.9) | |
| Twice/week (n=67) | 3 (4.5) | 46 (68.7) | 9 (13.4) | 9 (13.4) | |
| Three times/week (n=27) | 0 (0.0) | 14 (51.9) | 9 (33.3) | 4 (14.8) | |
| >3 times/week (n=15) | 0 (0.0) | 9 (60.0) | 2 (13.3) | 4 (26.7) | |

-Multivariate analysis

Multivariate logistic regression analysis demonstrated that primary school girls were at almost double-folded risk for obesity or overweight compared to primary school boys (aOR=1.93; 95%CI: 1.03-3.60, $p<0.05$). Habit of eating cheeps potatoes at school at abnormal rate was associated with almost three-folded risk for obesity or overweight compared to eating cheeps potatoes within normal (aOR=3.14; 95%CI: 1.17-8.41). School children who had the habit of taking fast foods at least three times/week were more likely to develop obesity or overweight compared to those who did not take fast foods (aOR=1.98; 95%CI:1.08-6.30). Variables of school grade, number of sisters, eating at school, eating cake at school and taking milk at school were not significantly associated with obesity or overweight among schoolchildren. Table 9

Table (11): Adjusted Odds ratio for risk factors of obesity among primary schoolchildren: Multivariate regression analysis

| | Adjusted OR | 95% CI | p-value |
|---|-------------|------------|---------|
| Gender | | | |
| Male ^a | 1.0 | --- | |
| Female | 1.93 | 1.03-3.60* | 0.040 |
| Eating cheeps potatoes at school | | | |
| Within normal ^a | 1.0 | --- | |
| Abnormal | 3.14 | 1.17-8.41* | 0.023 |
| Fast food | | | |
| No ^a | 1.0 | --- | |
| Once/week | 1.16 | 0.69-4.91 | 0.249 |
| Twice/week | 1.19 | 0.71-10.09 | 0.233 |
| At least three/week | 1.98 | 1.08-6.30* | 0.031 |

^a Reference category

*Statistically significant

Terms of school grade, number of sisters, eating at school, eating cake at school and taking milk at school were removed from the final logistic regression model (not significant)

5. DISCUSSION:

The rapid urbanization in gulf countries including Saudi Arabia in the last few decades is associated with unhealthy lifestyles described as the lifestyle syndrome which leads to very high rates of obesity and its consequent morbidity and mortality. In addition, childhood obesity is still considered a sign of healthiness and high social class in such countries. ³⁵

Most of previous local studies about childhood overweight and obesity depended on growth charts which were designed for western nations (usually CDC growth charts were used). These charts might not be appropriate for our children; they could underestimate the problem of obesity and mistakenly assign a child as normal while he is really overweight or

obese. The unique feature of the present study was the use of new growth charts which could be more appropriate for our children i.e. recently published 2007 WHO growth charts which were prepared using data from different countries.³⁴

The prevalence of overweight and obesity of primary school boys in the current study were 10.6% and 7.6%, respectively. This result is lower than that reported in other previous Saudi studies. A study done in 1996 showed that the prevalence of overweight and obesity were 11.7 and 15.8 respectively.³⁶ More recent study concluded that obesity prevalence increased among Saudi school boys to 24.5%.³⁷ In 2008, locally published study done in Al-Hassa showed that the prevalence of overweight of primary school boys was 14.2% while that of obesity was 9.7%, but the age range in that study was 10-12 years.²⁵ In a study conducted among primary school children in Riyadh (2013), a prevalence of 16% and 11% for overweight and obesity were reported among school boys.³²

On the other hand, overweight and obesity were significantly more common in girls in the present survey; prevalence of overweight of primary school girls was 16.8%, while it was 18.2% for obesity. This significant difference could be attributed to the less physical activity practices by girls at schools and at home in comparison to boys. Other local studies done on girls to estimate obesity and overweight locally were few. A study done in Alkhobar in 2003 on 2239 primary school girls concluded that the prevalence of overweight and obesity were 20% and 11%, respectively.²⁶ Recent study done in Riyadh and published at 2008 showed that prevalence of obesity in primary school girls was 14.9%.³⁸ In a study conducted among primary school children in Riyadh (2013), a

prevalence of 20% and 17% for overweight and obesity were reported among school girls.³² Al-Shehri et al (2013)³⁹ carried out a review is to evaluate the prevalence and the trends of obesity among Saudi children (2000-2012). The rates of overweight and obesity among school-age children have reached 23% and 9.3%, respectively.

Overall, a variation in the prevalence of obesity and overweight among children in geographical regions of the Kingdom was noted. This apparent increase in the prevalence of overweight and obesity in recent studies including ours, particularly among girls could reflect more deterioration in the situation in Saudi Arabia regarding childhood obesity which necessitate a need for rapid and effective action to decrease this problem burden.

The prevalence of overweight and obesity among primary schoolchildren in other parts of the world varies considerably. Some studies report higher rates than reported in the present study and some reported comparable results. Krassas et al. reported that the prevalence of overweight in Greek children aged 6–10 years (2001) was 25.3%.⁴⁰ In addition, Núñez-Rivas et al. in Costa Rica reported that the prevalence of overweight and obesity in children aged 7–12 years was 34.5% and 26.2% respectively.⁴¹ In a study done by Manzoli et al in 2005 in Italy, the prevalence of overweight among students aged 6–16 years was 40.6%⁴² while in Spain it was 40.0%.⁴³ In the USA, the prevalence of overweight and obesity among children has been reported to be 22% and 11% respectively.

⁴⁴ However some studies report similar or lower rates than ours. Hajian-Tilaki et al reported prevalence rates of obesity and overweight as 5-8% and 12.3% respectively among

schoolchildren (7-12 years) in Iran.⁴⁵ In India, the prevalence of overweight was reported as 10% among 10–15 year-olds,⁴⁶ and in Turkish students aged 12–17 years this figure was 10.6%.⁴⁷ In a study by Zini et al. in Malaysian primary-school children aged 9–10 years in 2005, the prevalence of overweight and obesity was 16.3% and 6.3% respectively.⁴⁸ The differences seen in the results of these studies could be attributed partially to the effect of genetic, lifestyle and environmental factors, and variations in the age groups of the samples and because of different study methods and definitions of obesity and underweight across the various studies.

This study could not find a significant association between childhood obesity and parental occupational or educational level. These factors were studied internationally and the results were conflicting. Studies from France⁴⁹ and Germany⁵⁰ showed that the risk of childhood obesity was significantly lower in those children with higher parental occupational and educational level. On the other hand; in agreement with our findings, studies from Italy⁵¹ and Turkey⁵² found no association between childhood obesity and parental occupational and educational level.

Taking a healthful breakfast can help prevent childhood obesity. It can also help teens that need to lose a few pounds. Younger children are more likely to eat breakfast than teens. Teens are often in a hurry to leave for school and are often drowsier in the morning and simply do not feel like eating. Providing an appetizing breakfast may encourage them to eat. Something with protein is a good choice but oatmeal is also filling. Some schools now serve

breakfast in the mornings but school breakfasts are often high in sugar and not particularly healthful.⁵³ In the current study, no significant association between taking breakfast and prevalence of obesity among studied school children was reported. However, the high prevalence of overweight and obesity among those who did not take breakfast supports the fact that healthful breakfast can help prevent childhood obesity. From the other side, the relatively high prevalence among those who always take breakfast reflects the unhealthy breakfast full of sugar and fats. So, the issue is not just to have a breakfast, but it should be healthy.

Teenage and childhood fast food obesity epidemic is growing around the world. Part of the reason is advertising, which has positioned fast food as an 'in' thing and this has appeal to the teenage and childhood group.⁵⁴ Fast food obesity is a problem because fast food meal contains low quality carbohydrates; high levels of saturated fat, white bread, and a sugary soda and also has relatively low fiber content. This kind of eating pattern is entirely gives negative effect, which resulted cardiovascular disease and diabetes. This alarming teenage and childhood fast food obesity epidemic is rapidly rising during the past few years. Fast food and obesity is interrelated with each other. For a person who takes fast food meal twice a week, obesity ratio will be raised about 50%.⁵⁴ In accordance with that, in the present study, schoolchildren who consume fast food meals more than three times per week were at almost double risk to develop obesity compared to those not consume fast foods.

As expected, eating abnormal amounts of high-caloric food stuffs as cake and cheeps potatoes was associated with significant increased risk for obesity in the present study.

This study has some limitations that should be mentioned. First of all, it ignores some important risk factors for overweight and obesity such as physical activity and family history of obesity and concentrating only on dietary factors alone. However, the main objective of the present study is estimation of the prevalence of obesity among primary school children in Taif and comparing boys with girls. Second, its cross-sectional design makes it difficult to sort out the causal relationships among variables studied. Despite that, it has an ultimate public health importance.

6. CONCLUSION:

Overweight and obesity among primary school children in Taif city, Saudi Arabia is a public health problem affecting more than one-third of girls and almost one fifth of boys. It confirms the significant association between some unhealthy dietary habits and increase prevalence of childhood obesity such as excessive eating of high caloric food stuffs such as cheeps potatoes as well as frequent intake of fast meals.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the

advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

7. RECOMMENDATIONS:

- 1- Health professionals and policy-makers should focus on primary prevention of childhood obesity, especially in areas with high prevalence aiming at controlling this growing epidemic.
- 2- Primary prevention of childhood obesity could be achieved through implementing strategies at primary schools such as encouraging healthy dietary habits, such as increasing fiber intake, reducing the consumption of junk foods and saturated fat, and increasing levels of physical activity at primary schools will result in important effects on public health.
- 3- Establishment of a uniform and universally accepted set of criteria for defining overweight and obesity in children and adolescents should be the foundation for addressing this emerging public health concern.
- 4- Further study on a larger scale with focusing on all risk factors for childhood overweight and obesity (dietary, physical activity, family history, personal habits, etc...).

5- Supervision of food quality at both at home and school should be done and healthy diet should be advocated.

8. REFERENCES:

1. Gidding SS, Leibel RL, Daniels S, Rosenbaum M, Horn LV, Rmarx GR. Understanding obesity in youth. *Circulation* 1996; 94: 3383-3387.
2. Stunkard A, Mendelson M. Obesity and the body image. I: characteristics of disturbances in the body image of some obese persons. *Am J Psychiatry* 1967; 123:1296-1300.
3. Stunkard A, Burt V. Obesity and the body image. II: age at onset of disturbances in the body image. *Am J Psychiatry* 1967; 123:1443-1447.
4. World health Organization (WHO). Obesity: preventing and managing the global epidemic. Report of a WHO consultation, Geneva, 3-5 June 1997, WHO. 1998.
5. Wilding J. Science, medicine, and the future: obesity treatment. *BMJ* 1997; 315: 997-1000.
6. Willett WC, Manson JE, Stampfer MJ, Colditz GA, Rosner B, Speizer FE. Weight change and coronary heart disease in women: risk within the 'normal' weight range. *JAMA* 1995; 273: 461-465.
7. Al-Nuaim AR. Effect of overweight and obesity on glucose intolerance and dyslipidemia in Saudi Arabia, epidemiological study. *Diabetes Res ClinPract* 1997; 36: 181-191.
8. Ashton W, Nanchahal K, Wood D. Body mass index and metabolic risk factors for coronary heart disease in women. *Eur Heart J* 2001; 22: 46-55.

9. Kordy MN, El-Gamal FM. A study of pattern of body mass index (BMI) and prevalence of obesity in a Saudi population. *Asia Pac J Public Health* 1995; 8: 59-65.
10. Al-Isa AN. Changes in Body Mass Index and Prevalence of Obesity Among Adult Kuwaiti Women Attending Health Clinics. *Annals of Saudi Medicine* 1997; 17: 307-311.
11. Troiano RP, Flegal KM, Kuczmarski RJ, Campbell SM, Johnson CL. Overweight prevalence and trends for children and adolescents: the National Health Examination Surveys, 1963-1991. *Arch Pediatr Adolesc Med* 1995; 149: 1085-1091.
12. Guo SS, Roche AF, Chumlea WC, Gradner JC, Siervogel RM. The predictive value of childhood body mass index values for overweight at age 35 years. *Am J Clin Nutr* 1994; 59: 810-819.
13. Guo SS, Chumlea WC. Tracking of body mass index in children in relation to overweight adulthood. *Am J Clin Nutr* 1999; 70: 145-148.
14. Al-Shammari SA, Khoja TA, Al-Maatouq MA. The prevalence of obesity among Saudi males in the Riyadh region. *Annals of Saudi Medicine* 1996; 16: 269-273.
15. Al-Shammari SA, Khoja TA, Al-Maatouq MA, Al-Nuaim LA. High prevalence of clinical obesity among Saudi females: a prospective, cross-sectional study in the Riyadh region. *J Trop Med Hyg* 1994; 97: 183-188.
16. Al-Nuaim AR, Bamgboy EA, Al-Rubeaan KA, Al-Mazrou Y. Overweight and obesity in Saudi Arabian adult population; role of socio-demographic variables. *J Community Health* 1997; 22: 211-223.

- 17.El-Hazmi MA, Warsy AS. Prevalence of obesity in the Saudi population. *Annals of Saudi Medicine* 1997; 17: 302-306.
- 18.Al-Nuaim AR, Al-Rubeaan KA, Al-Mazrou Y, Al-Attas O, Al-Daghari N, Khoja TA. High prevalence of overweight and obesity in Saudi Arabia. *Int J Obes Relat Metab Disord* 1996;20: 547-552.
- 19.Al-Nuaim AR. Population-based epidemiological study of the prevalence of overweight and obesity in Saudi Arabia, regional variation. *Ann Saudi Med.* 1997; 17(2):195-9.
- 20.Abalkhail B. Overweight and obesity among Saudi Arabian children and adolescents between 1994 and 2000. *East Mediterr Health J.* 2002; 8(4-5):470-9.
- 21.Al Herbish AS, Al Jurayyan N, Olasope A, Abdullah AM, Al Nuaim AA. Childhood obesity: referred cases to a tertiary health center in Riyadh, Saudi Arabia. *Saudi J Gastroenterol* 1999 May;5(2):85-8
- 22.El-Hazmi MA, Warsy AS. The prevalence of obesity and overweight in 1-18 year-old Saudi children. *Ann Saudi Med* 2002; 22(5-6):303-307.
- 23.Al-Rukban MO. Obesity among Saudi male adolescents in Riyadh, Saudi Arabia. *Saudi Med J* 2003; Vol. 24 (1): 27-33.
- 24.Khalid M. Is high-altitude environment a risk factor for childhood overweight and obesity in Saudi Arabia? *Wilderness Environ Med* 2008;19(3):157-63.
- 25.Al-Shehri JA. Childhood obesity prevalence among primary schoolboys at Al-Iskan sector, Holy Makkah, Saudi Arabia. *Int J Med Sci Public Health.* 2014; 3(2): 150-155.
- 26.Lafta RK, Kadhim MJ. Childhood obesity in Iraq: prevalence and possible risk factors. *Ann Saudi Med* 2005 Sep;25(5):389-93.

27. Al-Haddad FH, Little BB, Abdul Ghafoor AG. Childhood obesity in United Arab Emirates school children: a national study. *Ann Hum Biol* 2005;32(1):72-9.
28. Sorkhou I, Al-Qallaf K, Al-Shamali N, Hajia A, Al-Qallaf B. Childhood obesity in Kuwait--prevalence and trends. *Fam Med* 2003 Jul;35(7):463-4.
29. Badawi NE, Barakat AA, El Sherbini SA. Prevalence of overweight and obesity in primary school children in Port Said city. *Egyptian Pediatric Association Gazette* 2013; 61(1):31–36
30. Singh GK, Kogan MD, van Dyck PC. Changes in state-specific childhood obesity and overweight prevalence in the United States from 2003 to 2007. *Arch Pediatr Adolesc Med* 2010;164(7):598-607.
31. Yoshinaga M, Ichiki T, Tanaka Y, Hazeki D, Horigome H, Takahashi H, et al. Prevalence of childhood obesity from 1978 to 2007 in Japan. *Pediatr Int* 2010;52(2):213-7.
32. Muhihi AJ, Mpembeni RN, Njelekela MA, Anaeli A, Chillo O, Kubhoja S, et al. Prevalence and determinants of obesity among primary school children in Dar es Salaam, Tanzania. *Arch Public Health*. 2013 Oct 7;71(1):26.
33. Online Roasoft sample size calculator. Available at: <http://www.raosoft.com/samplesize.html>.
34. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM. CDC growth charts: United States. *Adv Data* 2000; 314:1–27.
35. Al-Ghamdi SH. The association between watching television and obesity in children of school-age in Saudi Arabia. *J Family Community Med* 2013 May-Aug;20(2)-83-89

- 36.al-Nuaim AR, Bamgboye EA, al-Herbish A. The pattern of growth and obesity in Saudi Arabian male school children. *Int J Obes Relat Metab Disord*. 1996 Nov;20(11):1000-5.
- 37.AL-Hazaa HM. Prevalence and trends in obesity among school boys in central Saudi Arabia between 1988 and 2005. *Saudi Medical Journal* 2007; 28 (10):1569-1574.
- 38.Alam AA. Obesity among female school children in North West Riyadh in relation to affluent life style. *Saudi Med J*. 2008; 29(8): 1139-44.
- 39.Al Shehri A, Al Fattan A, Al Alwan I. Obesity among Saudi children *Saudi J Obesity* 2013;1:3-9
- 40.Krassas GE, Tzotzas T, Tsametis C, Konstantinidis T. Prevalence and trends in overweight and obesity among children and adolescents in Thessaloniki, Greece. *Journal of Pediatric Endocrinology & Metabolism*, 2001; 14(Suppl. 5):1319–1326, discussion 1365.
- 41.Núñez-Rivas HP, Monge-Rojas R, León H, Roselló M. Prevalence of overweight and obesity among Costa Rican elementary school children. *Revista Panamericana de Salud Publica*, 2003; 13:24–32.
- 42.Manzoli L, Ripari P, Rotolo S, Di Giacinto G, Bellomo RG, Sorgentone S, et al. Prevalenza di obesità, sovrappeso e ipertensione nei bambini e adolescenti della provincia di Pescara [Prevalence of obesity, overweight and hypertension in children and adolescents from Abruzzo, Italy]. *Annali di igiene: medicina preventiva e di comunità*, 2005; 17:419–431.
- 43.Martínez Vizcaíno V, Salcedo Aguilar F, Franquelo Gutiérrez R, Torrijos Regidor R, Morant Sánchez A, Solera Martínez M, et al. Prevalencia de obesidad y tendencia de

- los factores de riesgo cardiovascular en escolares de 1992 a 2004 : estudio de Cuenca [Prevalence of obesity and trends in cardiovascular risk factors among Spanish school children, 1992–2004: the Cuenca (Spain) study]. *Medicina Clínica*, 2006; 126:681–685.
44. Terrell DF. Overweight and obesity prevalence rates among youth in the Carolinas. *North Carolina Medical Journal*, 2002; 63:281–286
45. Hajian-Tilaki KO, Sajjadi P, Razavi A. Prevalence of overweight and obesity and associated risk factors in urban primary-school children in Babol, Islamic Republic of Iran. *East Mediterr Health J*. 2011 Feb; 17(2):109-14.
46. Sidhu S, Marwah G, Prabhjot. Prevalence of overweight and obesity among the affluent adolescent school children of Amritsar, Punjab. *Collegium Antropologicum*, 2005, 29:53–55.
47. Oner N, Vatansever U, Sari A, Ekuklu E, Güzel A, Karasalihoğlu S, Boris NW.. Prevalence of underweight, overweight and obesity in Turkish adolescents. *Swiss Medical Weekly*, 2004, 134(35–36):529-533.
48. Zaini MZ, Lim CT, Low WY, Harun F. Factors affecting nutritional status of Malaysian primary school children. *Asia-Pacific Journal of Public Health*, 2005; 17:71-80.
49. Salane B and Karl S. Stabilisation of overweight prevalence in French children between 2000 and 2007. *Int J pediatr obes* 2009; 4(2): 66-72.
50. Lamerz A. Social class, parental education and obesity prevalence in a study of six year children in Germany. *Int J obes* 2005 Apr; 29(4): 377-80.
51. Fuiano. Prevalence and risk factor of overweight and obesity in a population of Italian school children: a longitudinal study. *J Endocrinal Invest* 2008 Nov; 31(11): 979-84.

52. Disciqlil G, Telcon N and Soylemz A. Obesity in Turkish children and adolescents: prevalence and non-nutritional correlates in an urban sample. Child care health dev 2009 May; 35(2): 153-8.
53. Martínez Vizcaíno V, Salcedo Aguilar F, Franquelo Gutiérrez R, Torrijos Regidor R, Morant Sánchez A, Solera Martínez M, et al. Prevalencia de obesidad y tendencia de los factores de riesgo cardiovascular en escolares de 1992 a 2004 : estudio de Cuenca [Prevalence of obesity and trends in cardiovascular risk factors among Spanish school children, 1992–2004: the Cuenca (Spain) study]. Medicina Clínica, 2006; 126:681–685.
54. Hajian-Tilaki KO, Sajjadi P, Razavi A. Prevalence of overweight and obesity and associated risk factors in urban primary-school children in Babol, Islamic Republic of Iran. East Mediterr Health J. 2011 Feb; 17(2):109-14.

استمارة بيانات الطالب

Appendix 1

الرقم الكودي: ()

الاسم:

العمر:

المرحلة الدراسية:

المقاييس الجسمية: الطول: م
الوزن: كجم

بسم الله الرحمن الرحيم

المحترم

إلى ولي أمر الطالب /

السلام عليكم ورحمة الله وبركاته

نظراً لما لوحظ عن ظاهرة السمنة لدى أبنائنا الأطفال وما يترتب عليها من أضرار صحية واجتماعية تؤثر على نفسيات هؤلاء الأطفال، فقد حرصت على عمل بحث ودراسة هدفها تحديد نسبة انتشار السمنة بين الأطفال الذين تتراوح أعمارهم بين 6- 12 سنة بالإضافة إلى علاقتها ببعض العادات الغذائية والنواحي الاجتماعية، وهذا لا يتم إلا بتعاونكم وتشجيعكم .

لذا أرجو من سعادتكم تعبئة استمارة الاستبيان المرسلة مع الطالب بكل دقة ووضوح حتى تكون الدراسة واقعية ويستفاد منها الاستفادة المرجوة، مع إعادتها مع الطالب في أسرع وقت مع اطمئنانكم بسرية المعلومات المدونة من قبلكم .

وفي حالة الاستفسار يمكن الاتصال بي على الرقم التالي:

جوال: 0500817684

شاكرين لكم تعاونكم المخلص لما فيه المصلحة العامة.

ولكم جزيل الشكر والامتنان.

أخوكم الباحث

د/ رشيد الغامدي

الإستبانة

1 - الرقم الكودي: ()

أرجو التكرم بالإجابة عن الأسئلة بوضوح بوضع علامة (√) في الخانة المناسبة وملء الفراغات بين

القوسين:

2 - تاريخ ميلاد الطالب: شهر () سنة ()

3 - المرحلة الدراسية : ()

4 - وزن الطالب عند الولادة: () صغير () طبيعي () كبير

5 - مع من يعيش الطالب: 1 ()

مع الوالدين 2 () مع الوالد فقط

3 () مع الوالدة فقط 4 () شخص آخر

حدد ()

6 - عدد أفراد الأسرة: ()

7 - عدد الإخوة الذكور () الإناث ()

8 - ترتيب الطالب بين إخوته: ()

9 - المستوى التعليمي للأم: 1 () لا تقرأ ولا تكتب 2 () ابتدائي

3 () متوسط 4 () ثانوي

5 () جامعي 6 () أعلى من الجامعة

10 - المستوى التعليمي للأب 1 () لا يقرأ ولا يكتب 2 () ابتدائي

3 () متوسط 4 () ثانوي

5 () جامعي 6 () أعلى من الجامعة

11 - هل يتناول الطالب وجبة الإفطار قبل الذهاب إلى المدرسة:

1 () دائما 2 () معظم الوقت

3 () أحيانا 4 () لا

12 - هل يتناول الطالب طعاما في المدرسة:

1 () دائما 2 () معظم الوقت

3 () أحيانا 4 () لا

إذا كان الطالب يتناول طعاما في المدرسة: ففي المعدل الطبيعي للطالب

13 - هل يفضل الطالب تناول الحليب: 1 () نعم 2 () لا

14 - هل يفضل تناول عصير الفراولة: 1 () نعم 2 () لا

15 - هل يفضل تناول المشروبات الغازية: 1 () نعم 2 () لا

16 - هل يفضل تناول بطاطس شيبس: 1 () نعم 2 () لا

17 - هل يفضل تناول الشوكولاتة / الحلوى: 1 () نعم 2 () لا


18 - هل يفضل تناول الكيك: 1 () نعم 2 () لا

19 - هل يفضل تناول البسكوت: 1 () نعم 2 () لا

- 20 - هل يفضل تناول البطاطس المقلية: 1 () نعم 2 () لا
- 21 - هل يفضل تناول الفواكه: 1 () نعم 2 () لا
- 22 - ماهي الوجبات التي يتناولها الطالب بانتظام (يمكن اختيار أكثر من إجابة):
- 1 () الإفطار 2 () الغداء
- 3 () العشاء 4 () الوجبات الثلاثة
- 23 - ما معدل تناول الطالب للخضروات في اليوم (مجموع الخضروات في اليوم مثل طفل يأكل 1 خيار و 1 طماطم و 1 جزر يعتبر 3 مرات):
- 1 () مرتين فأقل 2 () 3-5 مرات
- 3 () أكثر من 5 مرات
- 24 - ما معدل تناول الطالب للفواكه في اليوم:
- 1 () مرتين فأقل 2 () 3-5 مرات
- 3 () أكثر من 5 مرات
- 25 - هل يتناول الطالب طعاماً بين الوجبات الرئيسية :
- 1 () دائماً 2 () معظم الوقت
- 3 () أحياناً 4 () لا
- إذا كان الطالب يتناول طعاماً بين الوجبات, ففي المعدل الطبيعي للطالب:
- 26 - هل يفضل الطالب تناول العصيرات: 1 () نعم 2 () لا
- 27 - هل يفضل الطالب تناول المشروبات الغازية : 1 () نعم 2 () لا
- 28 - هل يفضل الطالب الشوكولاته/ الحلوى : 1 () نعم 2 () لا
- 29 - هل يفضل الطالب تناول الفواكه : 1 () نعم 2 () لا
- 30 - هل يفضل الطالب تناول البطاطس المقلية: 1 () نعم 2 () لا
- 31 - هل يفضل الطالب البسكويت: 1 () نعم 2 () لا
- 32 - هل يفضل الطالب تناول بطاطس الشيبس : 1 () نعم 2 () لا
- 33 - هل يفضل الطالب تناول الكيك : 1 () نعم 2 () لا
- 34 - ما معدل تناول الطالب للمأكولات السريعة في الأسبوع:
- 1 () مرة 2 () مرتين
- 3 () ثلاث مرات 4 () أكثر من ثلاث مرات

Appendix 2

Appendix 3



UNDER PEER REVIEW