

Original Research Article

Effect of *Cardisoma guanhumi* (land crab) extract on hematology and lungs histology in swiss mice infected with *Bordetella pertussis*

Abstract

Pertussis is an acute human respiratory tract disease caused by *Bordetella pertussis*, a known international pathogen that affects infants, children and adults.. This study is aimed at investigating the changes in hematological parameters and histopathological changes of the lungs connected with *Bordetella pertussis* infection in swiss mice and to evaluate the potential of *Cardisoma guanhumi* extract to reverse these adverse changes in relation to erythromycin treatment. The animals were divided into five groups: group 1 was normal control, group 2 was infected with *Bordetella pertussis* without treatment (negative control), group 3 and 4 were *Bordetella pertussis* infected and treated with 300mg/kg and 600mg/kg of the extract respectively and group 5 was infected and treated with 4000mg/70kg of erythromycin in divided doses. The animals were inoculated with a single infectious dose of *Bordetella pertussis* bacteria and were consequently treated with the graded doses of the extract and erythromycin for a period of eighteen days, after the animals were confirmed infected. The mice were humanely sacrificed using diethyl ether anesthesia and blood samples taken for liver function investigation and liver tissue harvested and processed for histological examination. The result showed that *Cardisoma guanhumi* extract reversed the changes in the hematological parameters and pathological changes in the lungs of mice infected with *Bordetella pertussis* in a dose and time dependent manner which suggests prophylactic and curative potentials of *Cardisoma guanhumi* extract against *B. pertussis*.

Introduction

Since the identification of pertussis as a clinical disease more than 1,600 years ago, *Bordetella pertussis* remains a major global pathogen that affects infants, children, and adults (Dorji *et al.*, 2017). *B. pertussis* is a complicated bacterium that expresses numerous bacterial factors with immune modulating functions and produces different bacterial factors responsible for the symptoms seen during the disease (Mattoo and Cherry, 2005). Pertussis is mostly a toxin mediated disease. The bacteria close to the cilia of the respiratory epithelial cells, produce toxins that paralyze the cilia, and cause inflammation of the respiratory tract, which interferes with the clearing of pulmonary secretions (Sheridan *et al.*, 2012). Pertussis antigens appear to allow the organism invade host defenses by enhancing lymphocytosis but impairing chemotaxis (NCDC,

2011). The understanding of this leaves us with the question of why has pertussis remained a major challenge to conquer internationally and what are the imminent dangers if nothing is done to curtail the prevalence of this disease. Numerous studies have progressively proven that pertussis toxin (PT) is the leukocytosis enhancing factor that is produced by *Bordetella pertussis*. Furthermore, other experimental animals, including rats (Samore and Siber 1992), pigs (Elahi *et al.*, 2005) and macaques (Hinds *et al.*, 1996; Pauza *et al.*, 1997), develop leukocytosis (an increase in the number of white blood cell especially during infection) when administered purified pertussis toxin (PT). Mice and baboons inoculated with *B. pertussis* have lowered levels of leukocytosis when induced with PT-specific monoclonal antibodies (Nguyen *et al.*, 2015). In a study by Temple *et al.*, (2009) to determine the influence of living *Bordetella pertussis* on the induction and duration of patho-physiological reactions in mice infected intranasally with graded doses of culture, reported that fatally infected mice showed loss of body weight, spleen atrophy, obvious hypothermia and hypoglycemia (low blood sugar), and highly increased levels of leukocytes and serum immune-reactive insulin while non fatal infected mice showed normal weight gain, almost normal temperature, spleen enlargement, not too pronounced hypoglycemia, lower but obvious increased levels of leukocytes and serum immune-reactive insulin, and histamine sensitization. Leukocytosis (an increase in the number of white blood cell especially during infection) and lymphocytosis (an increase in the number of lymphocytes in the blood) are clear indications of pertussis infection (Beck *et al.*, 2014). Similarly, Momoh *et al.*, (2013) noted that infection with *S. typhi*, similar bacteria to the one under study produced a reduction in pack cell volume (PCV), Red blood cell (RBC) and Hemoglobin (Hb) while there was an increase in monocyte, neutrophil and WBC but there was no significant difference. Crabs are decapod crustaceans which belong to the infra order Brachyura. They are mainly covered with thick exoskeleton. Their lower region is completely hidden under the thoracic cavity. They can be found in most tropical and subtropical regions of the world as reported by Sammy *et al.*, (2009). *Cardisoma guanhumi* is a species of land crab that is found in tropical and subtropical estuaries and other maritime areas of land along the Atlantic coast of the Americas (Renata *et al.*, 2012). They are known to be good sources of essential macro and micro minerals such as potassium, phosphorus, calcium, magnesium, copper, iron, manganese, and zinc. Bae (2010) and Sujeetha *et al.*, (2015) reported the biomedical and nutritional properties of crabs to include Omega3 (a poly unsaturated acid) contained in crab meat which helps in providing protection against heart diseases and inhibit the aggressive behavior. Mahae *et al.*, (2011) noted that the selenium contained in crab meat plays an important role in human's antioxidant defense system by preventing cells and tissues from damage and helps in proper functioning of the immune system and metabolism of thyroid hormone while riboflavin present in them helps in the production of steroids and red blood cells, maintenance of the skin, promote normal growth and iron absorption from the digestive tract and support antioxidant activity. Garry (2015) explained that copper and phosphate content in crab helps in the absorption, storage and metabolism of iron and is involved in the formation of red blood cells. Ming *et al.*, (2010) and Suneeta (2014) in their separate studies reported that crabs lower blood pressure, protect against heart diseases and possess anti

inflammatory properties. . Chitosan derived from crab shell have several properties including anti-microbial and antibacterial properties due to its peculiar characteristics (Mahae *et al.*, 2011). Chitosan fights against numerous pathogenic organisms like fungi, spoilage microorganisms, gram positive and gram negative bacteria (Mahae *et al.*, 2011). However, this is the first attempt at establishing the antibacterial effects of crab extract which was prejudiced by unreliable evidence of the healing potential of crabs in whooping cough among the bonny people of Rivers State, Nigeria This study is aimed at determining the effect of *Cardisoma guanhumi* (land crab) extract on hematology and lungs histology in swiss mice infected with *Bordetella pertussis*

Material and methods

Sample Collection and Identification

Cardisoma guanhumi was caught using a trap in the buguma creek, Rivers State, Nigeria. The samples collected were transferred into perforated plastic containers to allow for air during transportation and was transported to the Pharmacognosy research laboratory, Department of Pharmacognosy, University of Port Harcourt, Nigeria. The samples were identified Mr. Otufu Paciya using Food and Agriculture Organization species identification sheets for fresh water and marine crab species.

Method of Extraction

Using the Shahidi and Synowiecki (1991) extraction method, 60 of the freshly collected crabs were sacrificed and the shell separated from the meat and washed with tap water to remove all impurities. The crab shells and meat were then transferred to the oven and dried at 70°C until they were completely dry. Using a laboratory mortar and pistle, the dried crab shells and meat were ground and sieved into the size of 500µm. 40g of the sieved crab was measured using WANT precision electric weighing balance made by WANT balance instrument company limited, China. into a beaker and 200ml of cord liver oil was added and stirred with magnetic stirrer until it was completely mixed for 20minutes. The beaker was then transferred into a water bath at a temperature of 60°C and allowed for 30mins. The mixture was then filtered with a white handkerchief to drain off the oil and the residue transferred into a beaker. The residue was treated with 2% potassium hydroxide (KOH) at a ratio of 1:20 w/v and was stirred continuously for 2hours at a temperature of 90°C to remove protein from the crab. The sample was filtered under vacumm and the residues were continuously washed until the pH became neutral i.e pH=7. This was done to ensure that all the salt had been removed after removing the protein. The deprotienized crab was transferred into an oven and dried at 60°C until it was completely dry (Shahidi and Synowiecki, 1991). Two point five percent w/v of hydrochloric acid (Hcl) was used at room temperature (20°C) for 6 hours to remove the mineral content of the deproteinized crab at a ratio of 1:20 w/v. The samples were filtered under vacuum and washed with tap water until the pH was neutral. The demineralized crab were then transferred to the oven and dried at a temperature 60°C until completely dried. (Shahidi and Synowiecki, 1991).The demineralized crab was treated with 300ml acetone for 10mins and dried for 2hrs at an ambient temperature and the residues were removed to achieve decolourization. The decolourized sample was washed in running water, filtered and dried at 60°C until it was completely dried to obtain crab chitin (Shahidi and Synowiecki, 1991). Deacetylation of chitin was carried out using the method of Yen *et al.*, (2009). The obtained chitin was treated with 40% w/v aqueous sodium hydroxide (NaOH) in the ratio of chitin to the solution 1:15 w/v at 105°C in a water bath for 2hrs. Thereafter, the chitin was filtered with filter pump and washed with deionized water until pH was neutral to

obtain the extract. The obtained extract was then dried at 60°C for 2hrs in the oven. The dried extract was preserved in a well labeled bottle and kept for the experiment.

Isolation of Test Organism

The test organism *Bordetella pertussis* (ATCC®9340™) was gotten from the American Type Culture Collection (ATCC), USA. The culture media used for isolation according to ATCC is medium 35: Bordet Gengou/Broth medium from a human clinical specimen at a growth temperature of 37°C in aerobic atmosphere. The product was received freeze-dried at 2°C-8°C and stored at -80°C. The bacterium was reconstituted using Regan-Lowe agar (Charcoal blood Agar) in the Department of Microbiology laboratory, University of Port Harcourt.

Experimental Design

A total of one hundred and twenty-two (122) animals (swiss mice) were divided into five groups for the curative treatment study. Group 1 (normal) had 10 animals, group 2 (negative control) had 28 animals; group 3 and 4 consisted of 28 swiss mouse each. Group 1 served as the normal control without treatment but was fed with the normal animal feed and water. Group 2 (negative control group) consisted of *B. pertussis* inoculated mice without treatment. Group 3 consisted of *B. pertussis* infected mice exposed to low dose (300mg/kg) of *Cardisoma guanhumi* extract while group 4 consisted of *B. pertussis* infected mice exposed to high dose (600mg/kg) of *Cardisoma guanhumi* extract and group 5 consisted of *B. pertussis* infected mice exposed to 4000mg/70kg of erythromycin. On day 0, at day 6days interval and on day 18, seven animals were sacrificed using diethyl ether anesthesia; samples of blood were collected and the liver removed for assessment of liver function status and histopathological examination respectively.

Challenging apparently healthy animals with *Bordetella pertussis* infective dose

One hundred and twenty-two animals were intraperitoneally challenged with the infective dose of *Bordetella pertussis* which was calculated to be 5×10^5 cfu/ml. After infection had set in (through physical observation of signs like weakness, non-productive cough, anorexia and the isolation of the organism from the blood of the infected animals on day 0) seven animals were sacrificed and blood samples and liver tissue was collected for preliminary investigation and the rest animals from the other treatment groups were given two times daily of the various doses of the extract and the standard antibiotics (erythromycin) for 18days.

Antibiotic and Extract concentration Preparation

The extract solution for the study was prepared by dissolving 0.5g of the extract in 1ml of dimethyl-sulfoxide (DMSO) solvent to have a stock concentration of 500mg/ml. Since 70kg (70000g) takes 4000mg of erythromycin daily at severe case of whooping cough, then 25g (average weight of test animal) will take $25g \times 4000mg / 70000g = 1.429mg$. This means that 25g will take 1.429mg/ml or 2.858mg/0.5ml (1.429×2) or 5.716mg/0.25ml (1.429×4). 5.716mg/0.25ml was prepared from 500mg tablet of erythromycin tablet thus $500mg/Xml = 5.716mg/ml$. therefore, $X = 500mg \times ml / 5.716mg = 87.47ml$. Hence, 500mg tablet of erythromycin was dissolved in 87.47ml distilled water to prepare the erythromycin solution for the study.

Blood collection

Each animal was anaesthetized with diethyl ether in a desiccator and blood was collected by cardiac puncture method and transferred into a well labeled sample bottles containing anti-coagulant.

Hematological Analysis

The hematological analysis was carried out using the method as described by Randox Laboratories Limited, United Kingdom.

Histopathology studies

The study animals (swiss mice) were subjected to diethyl ether anesthesia in a decicator and dissected aseptically to collect the liver for histopathological studies. The collected tissues were kept in 10% chloroform for preservation and were subsequently trimmed to the size of 3-4mm thickness for fixation. These tissues were fixed, dehydrated, cleared, impregnated, embedded, sectioned and stained with hematoxylin and eosin before mounting according to the method described by Baker (1945).

Statistical Analysis

The results of the measurements are shown as Mean \pm Standard Deviation of Mean. The mean differences were obtained by ANOVA and post hoc with least significant difference (LSD) (Mead & Curnow, 1982).

Results

Effect of *Cardisoma guanhum* extract on Post-Inoculation Treatment on Hematological Indices in *B. pertussis* infected mouse

From the result, inoculation of mice with infective dose of *B. pertussis* reveals a gradual decrease in PCV, Hemoglobin, Platelet, RBC and Eosinophil levels when compared to the normal control but these alterations was reversed when treated with *Cardisoma guanhum* extract. Treatment with *Cardisoma guanhum* extract reveals a steady increase in PCV (table 1) Hemoglobin (table 2) platelet (table 3) RBC (table 4) as the day progresses. Contrarily, the negative control shows a constant decrease in all the parameters. Statistical ANOVA comparison shows a significant difference ($p < 0.05$) between normal control, negative control, standard drug and the treated groups on 6th, 12th and 18th day. Similarly, inoculation with an infective dose of *B. pertussis* shows an increase in WBC, neutrophils, lymphocytes and monocytes when compared with the normal control. However, treatment with *Cardisoma guanhum* extract reversed the variations causing a gradual decrease in the hematological parameters (neutrophils, lymphocytes, WBC and monocytes). ANOVA comparison between the normal control, negative control, standard drug and the treatment group shows a significant difference in WBC on day 6, 12 and 18. Also, neutrophils showed no significant ($p > 0.05$) difference on day 12 and 18 when compared to standard drug. Similarly, there was a significant difference ($P < 0.05$) in lymphocytes on day 6 and 12 but showed no significant ($P > 0.05$) on day 18 when compared to the normal control and standard drug. Finally monocytes count showed no significant difference in relation to standard drug on day 6 and 12 but a significant difference ($P < 0.05$) when compared to negative control on day 18.

Table 1: Effect of Post – IT on *Cardisoma guanhum* extract on PCV (g/dl)] in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	37.00 \pm 0.000	37.00 \pm 0.000	37.00 \pm 0.000	37.00 \pm 0.000

Negative control	24.67±1.155	22.33±.577	18.67±1.155	15.33±1.155
Erythromycin	24.67±1.155	28.33±.577	34.33±.577	36.67±.577
Low Dose	24.67±1.155	26.67±.577 ^{abc}	29.67±.577 ^{abc}	31.33±1.155 ^{abc}
High Dose	24.67±1.155	26.67±.577 ^{abc}	31.67±1.155 ^{abc}	34.67±1.528 ^{abc}

a= Significant (p<0.05) between test groups and control

b= Significant (p<0.05) between test groups and negative control

c= Significant (p<0.05) between test groups and erythromycin

Control = Animal fed with normal feed and water

Negative control = Animal infected with *Bordetella pertussis* without treatment

Low dose = 300mg/kg

High dose =600mg/kg

Erythromycin = standard antibiotics drug

Table 2: Effect of Post - IT on *Cardisoma guanhum* extract on Hgb (g/dl)] in B. pertussis infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	12.30±0.000	12.30±0.000	12.30±0.000	12.30±0.000
Negative control	8.40±.200	7.47±.115	7.00±.173	6.47±.153
Erythromycin	8.40±.200	9.13±.208	11.10±.173	12.20±.100
Low Dose	8.40±.200	8.37±.058abc	9.07±.058abc	9.53±.153abc
High Dose	8.40±.200	8.53±.058abc	10.33±.115abc	11.60±.100abc

Table 3: Effect of Post - IT on *Cardisoma guanhum*i extract on Platelets count ($\times 10^3/\mu\text{l}$) in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	100.00 \pm 0.000	100.00 \pm 0.000	100.00 \pm 0.000	100.00 \pm 0.000
Negative control	56.00 \pm 0.000	53.33 \pm 1.155	50.33 \pm .577	48.33 \pm .577
Erythromycin	56.00 \pm 0.000	73.67 \pm 3.512	85.00 \pm 2.000	99.33 \pm .577
Low Dose	56.00 \pm 0.000	63.67 \pm 1.528 ^{abc}	70.33 \pm 2.517 ^{abc}	76.00 \pm 2.000 ^{abc}
High Dose	56.00 \pm 0.000	66.67 \pm 1.155 ^{abc}	76.33 \pm .577 ^{abc}	91.67 \pm .577 ^{abc}

Table 4: Effect of Post - IT on *Cardisoma guanhum*i extract on RBC ($\times 10^3/\mu\text{l}$) in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	9.29 \pm 0.000	9.29 \pm 0.000	9.29 \pm 0.000	9.29 \pm 0.000
Negative control	4.90 \pm .300	4.50 \pm .200	3.77 \pm .115	3.07 \pm .153
Erythromycin	4.90 \pm .300	6.33 \pm .058	7.90 \pm .200	9.28 \pm .017
Low Dose	4.90 \pm .300	5.50 \pm .173 ^{abc}	6.00 \pm .265 ^{abc}	6.80 \pm .200 ^{abc}
High Dose	4.90 \pm .300	5.83 \pm .058 ^{abc}	6.80 \pm .300 ^{abc}	8.83 \pm .115 ^{abc}

Table 5: Effect of Post - IT on *Cardisoma guanhum*i extract on WBC ($\times 10^3/\mu\text{l}$) in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	5.10 \pm 0.000	5.10 \pm 0.000	5.10 \pm 0.000	5.10 \pm 0.000

Negative control	5.77±.058	6.43±.153	7.70±.173	9.27±.153
Erythromycin	5.77±.058	5.33±.058	5.20±0.000	5.13±.058
Low Dose	5.77±.058	5.73±.058 ^{abc}	5.47±.058 ^{abc}	5.27±.058 ^b
High Dose	5.77±.058	5.63±.058 ^{abc}	5.43±.058 ^{abc}	5.17±.058 ^b

Table 6: Effect of Post - IT on *Cardisoma guanhumi* extract on Neutrophil count ($\times 10^3/\mu\text{l}$) in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	20.00±0.000	20.00±0.000	20.00±0.000	20.00±0.000
Negative control	24.67±1.155	27.33±.577	30.00±1.000	42.00±1.732
Erythromycin	24.67±1.155	21.67±.577	21.00±0.000	20.00±0.000
Low Dose	24.67±1.155	24.00±0.000 ^{abc}	23.00±0.000 ^{abc}	23.00±0.000 ^{abc}
High Dose	24.67±1.155	23.00±0.000 ^{abc}	22.33±.577 ^{ab}	20.67±.577 ^b

Table 7: Effect of Post - IT on *Cardisoma guanhumi* extract on Lymphocyte count ($\times 10^3/\mu\text{l}$) in *B. pertussis* infected mice

	DAY 0	DAY 6	DAY 12	DAY 18
Control	82.00±0.000	82.00±0.000	82.00±0.000	82.00±0.000
Negative	88.00±1.000	91.33±2.082	94.00±1.000	101.67±2.517

control

Erythromycin	88.00±1.000	84.33±.577	83.67±.577	82.00±0.000
Low Dose	88.00±1.000	87.67±.577 ^{abc}	85.67±.577 ^{abc}	84.67±.577 ^b
High Dose	88.00±1.000	85.33±1.155 ^{ab}	84.33±.577 ^{ab}	82.67±.577 ^b

Table 8: Effect of Post - IT on *Cardisoma guanhumi* extract on Monocyte count (x10³/μl) in *B. pertussis* infected mice


	DAY 0	DAY 6	DAY 12	DAY 18
Control	2.00±0.000	2.00±0.000	2.00±0.000	2.00±0.000
Negative control	4.67±.577	6.67±.577	9.00±1.000	11.00±1.000
Erythromycin	4.67±.577	3.33±.577	2.67±.577	2.00±0.000
Low Dose	4.67±.577	4.00±0.000 ^{ab}	3.67±.577 ^{ab}	3.33±.577 ^b
High Dose	4.67±.577	4.00±0.000 ^{ab}	3.33±.577 ^{ab}	2.67±.577 ^b

Effect of *Cardisoma guanhumi* extract on lungs Histo-architecture in *B. pertussis* infected mice

The lungs histo-pathological examination of the control animals reveals a normal structure with clear alveolar spaces, epithelial cells and blood vessels. There was no histologic alteration in the lungs. However, mice lungs tissue infected with *B. pertussis* with no treatment for four days shows various distortions in the lungs tissue such as widened interstitial, interstitial inflammation and bullae formation. The lungs tissues in mice administered with (300mg/kg) of *Cardisoma guanhumi* extract for 12 days shows infected tissues with bullae formation while those treated for 18 days shows no interstitial inflammation. The lungs tissues not treated (negative control) for 18 days shows widened interstitial, interstitial inflammation, hemorrhage and bullae formation. However, Lungs tissues in mice treated with (600mg/kg) of *Cardisoma guanhumi* extract for 6 days, and 12 days shows mild interstitial inflammation, bullae formation and widened interstitial while treatment for 18 days shows no obvious histologic change while for those treated with


erythromycin for 6 days showed interstitial inflammation and bullae formation. However, lungs tissues infected with *B. pertussis* and treated with erythromycin for 18 days shows no obvious histologic change and appeared normal as the control. This is shown in plates 1-14

Histological Plates



alveolar spaces epithelial cells blood vessels

Plate 1: Photomicrograph of lungs tissue of normal mice showing no histologic alteration



widened interstitium inflammatory bullae

Plate 2: Photomicrograph of lungs of mice infected with *B.pertussis* showing interstitial inflammation and bullae formation (Day 0)




Plate 3: Photomicrograph of lungs of mice infected with *B.pertussis* without treatment for 6days showing interstitial inflammation and bullae formation




Plate 4: Photomicrograph of lungs of mice infected with *B.pertussis* treated with 4000mg/70kg of erythromycin for 6days showing interstitial inflammation and bullae formation




Plate 5: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 300mg/kg of *Cardisoma guanhumi* extract for 6days showing interstitial inflammation and bullae formation




Plate 6: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 600mg/kg of *Cardisoma guanhumi* extract for 6days showing interstitial inflammation and bullae formation




Plate 7: Photomicrograph of lungs of mice infected with *B.pertussis* without treatment for 12days showing interstitial inflammation and bullae formation




Plate 8: Photomicrograph of lungs of mice infected with *B.pertussis* treated with 4000mg/70kg of erythromycin for 12days showing no obvious histologic change with clear alveolar spaces and pulmonary vessels




Plate 9: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 300mg/kg of *Cardisoma guanhumi* extract for 12days showing mild interstitial inflammation and bullae formation




Plate 10: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 600mg/kg of *Cardisoma guanhumi* extract for 12days showing interstitial inflammation and bullae formation




Plate 11: Photomicrograph of lungs of mice infected with *B.pertussis* without treatment for 18days showing interstitial inflammation and bullae formation






Plate 12: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 4000mg/70kg of erythromycin for 18days showing no obvious histologic change



alveolar spaces epithelial cells blood vessels

Plate 13: Photomicrograph of lungs of mice infected with *B.pertussis* and treated with 300mg/kg of *Cardisoma guanhumi* extract for 18days showing no histologic alteration



alveolar spaces epithelial cells blood vessels

Plate 14: Photomicrograph of lungs of mice infected with *B. pertussis* and treated with 600mg/kg of *Cardisoma guanhumi* extract for 18days showing normal lungs tissue

Discussion

Several studies have shown that during *B. pertussis* infection, the common hematological changes include leukocytosis (an increase in the number of white blood cell (WBC)) lymphocytosis (an increase in the number of lymphocytes in the blood) and monocytosis (an increase in the number of monocytes in the blood (Back *et al.*, 2014; Temple *et al.*, 2009, Nguyen *et al.*, 2015). Significant decrease in mean levels reduced PCV (Pack Cell Volume), Red Blood Cell (RBC) and Hemoglobin while there is an increased neutrophil (Momoh *et al.*, 2013). The result of this study agrees with the reports from the other researchers. The increase in neutrophil and while blood cell are associated with the invasion of the hemopoietic organ (bone marrow) by *B. pertussis* while monocytosis and lymphocytosis are attributed to the increased release of these cells from the myeloid/ lymphoid tissues in response to the infection (Das and Mukherjee, 2003, Anusuya and Sumathi, 2015). However, the decrease in RBC, PCV and hemoglobin could be as a result of the destruction of Red Blood Cell by the infection (Dangana *et al.*, 2000) hemophagocytosis (a potentially fatal disease of normal but over active histiocytes and lymphocytes that are common in infants) and bone marrow suppression (Khosla *et al.*, 1995). The result further reveals that the post inoculation treatment with *Cardisoma guanhumi* extract reversed the usual trend of significant increase ($P < 0.05$) in WBC, lymphocytes, monocytes, neutrophils and decrease in PCV, RBC, platelets and hemoglobin usually associated with *B. pertussis* infection. Therefore, the post inoculation treatment with *Cardisoma guanhumi* extract reversed *B. pertussis* invasion of the hemopoietic organs, bone marrow suppression, destruction of RBC, hemophagocytosis and the invasion of the macrophages although the level of reversal is time and suggests that prophylactic and curative potentials of *Cardisoma guanhumi* extract against *B. pertussis* when combined with other therapeutic agents. The result of this study agrees with the report of Andreasen and Carbonetti, 2008, Karen *et al.*, 2017 who said infection of *B. pertussis* causes lung tissue inflammation, necrosis and widened interstition. However,

treatment with *Cardisoma guanhumi* slowly reversed the trend with increase in time and dosage. The result confirms the anti-*Bordetella pertussis* property of the extract.

Conclusion

Inoculation of mice with an infective dose of *B. pertussis* produces hematological changes such as leukocytosis (an increase in the number of white blood cell (WBC)) lymphocytosis (an increase in the number of lymphocytes in the blood) and monocytosis (an increase in the number of monocytes in the blood). Significant decrease in mean levels of PCV (Pack Cell Volume), Red Blood Cell (RBC) and Hemoglobin with an increase in neutrophil but treatment with *Cardisoma guanhumi* extract reversed the changes in a dose and time dependent manner. Similar, infection with *B. pertussis* caused distortions in the lungs tissue such as widened interstitial inflammation and bullae formation but treatment with *Cardisoma guanhumi* extract reversed the changes in a dose and time dependent manner.

References

- Andreasen, C. & Carbonetti, N.H. (2008). Pertussis toxin inhibits early chemokine production to delay neutrophil recruitment in response to *Bordetella pertussis* respiratory tract infection in mice. *Infections and Immunology*, 76:5139–5148.
- Bae keun park (2010) “Application of chitin and its derivatives in biological medicine” *international food research journal* **11**(12): 5152-5164
- Beck, T.C., Gomes, A.C. & Cyster, J.G. (2014) CXCR4 and a cell-extrinsic mechanism control immature B lymphocyte egress from bone marrow. *Journal of Experimental Medicine*. **211**:2567–81
- Dangana, A., Ajobiwe, J. and Nuhu, A. (2010). Hematological changes associated with *Salmonella typhi* and *Salmonella paratyphi* in humans. *International journal of Biomedical Health Science*, **6**:219-222
- Das, B.K. and Mukherjee, S.C. (2003). Toxicity of cypermethrin in *Laberohita* fingerlings: Biochemical enzymatic and hematological consequence. *Journal of comparative Biochemistry, Physiology, Toxicology and Pharmacology*, **134**: 109-121
- Dorji, D., Mooi, F., Yantorno, O., Deora, R., Graham, R.M. & Mukkur, T.K. (2017). *Bordetella Pertussis* virulence factors in the continuing evolution of whooping cough vaccines for improved performance. *Medical Microbiology and Immunology*, **10**:17-52
- Elahi, S., Brownlie, R. & Korzeniowski, J. (2005). Infection of newborn piglets with *Bordetella pertussis*: a new model for pertussis. *Infection and Immunology*. **73**:3636–45.
- Garry Kerch, (2015) The potential of chitosan and its derivatives in prevention and treatment of age related disease. *International food research journal*, **13**:2158-2182
- Hinds, P.W., Yin, C. & Salvato, M.S., (1996) Pertussis toxin induces lymphocytosis in rhesus macaques. *Journal of Medical Primatol.* **25**:375–81.

- Karen, M., Scanlon, Y., Snyder G. & Nicolas, H. (2017) Fatal Pertussis in the Neonatal Mouse Model Is Associated with Pertussis Toxin-Mediated Pathology beyond the Airways. *Infectious Immunology*, **85** (11):355-17.
- Khosla, S.N., Singh, R., Singh, G.P. & Trehan, V.K. (1988). The spectrum of hepatic injury in enteric fever. *American Journal of Gastrointestinal*, **83**:413-16.
- Mahae , N., Chalat, C. and Muhamud,P. (2011) Antioxidant and antimicrobial properties of chitosan sugar complex. *International food research journal*, **18** (4) 2011, page no 1543-1551 .
- Mattoo, S. & Cherry, J.D. (2005). Molecular pathogenesis, epidemiology, and clinical manifestations of respiratory infections due to *Bordetella pertussis* and other *Bordetella* subspecies. *Clinical Microbiology Review*, **18**:326–382.
- Mead, R. & Curnow, R. N. (1982). A simple statistical method in Agriculture and Experimental Biology. Chapman Hall, London, UK, 33-46
- Ming Kong, Ke xing & Hyun kin park (2010). “Antimicrobial properties of chitosan and mode of action: A state of art review” *International journal of food microbiology*, **144** (1):50-63.
- Momoh, A. O., Adebolu, T.T. & Ogundare, A.O. (2013). Evaluation of beniseed extract and fermented liquors in treatment of diarrhea in albino rats infected with *Salmonella typhi*. *European Journal of Biology and Medical Science Research*, **1**(2):16-23.
- Nguyen, A.W., Wagner, E.K., Laber, J.R. (2015). A cocktail of humanized anti-pertussis toxin antibodies limits disease in murine and baboon models of whooping cough. *Sci Transl Med*. 7:316-95
- Randox Laboratories Limited (2015). 55 Diamond Road, Crumil. County Antrim, BT294QY, United Kingdom.
- Reitman, S. & Frankel, S. (1957). A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American journal of clinical pathology*, **28**:56-63
- Renata, A. S., Jose, R. F. and Fábio, H. V, (2012). Development of male reproductive system of the blue land crab *Cardisoma guanhumi* Latreille, 1828 (Decapoda: Gecarcinidae). *Acta Zoologica*, Volume 93, ssue 4, pp 390–399
- Sammy, D., Grave, N., Dean, P. and Pentcheff, N. (2009) A classification of living a fossil genera of decapod crustaceans. *Raffles bulletin of zoology*. **21**:1-109
- Schmidt, E. & Schmidt, F. W. (1963) Enzyme. *Biology and Clinical science*, **3**:1-6

- Shahidi, F. & Synowiecki, J. (1991) Isolation and characterization of nutrients and value- added products from snow crab (*Chionoecetes Opilio*) and shrimp (*Pandalus Borealis*) processing discards. *Journal of Agricultural Food Chemistry*, **39** (8):1527–1532.
- Sheridan, S.L., Ware, R.S., Grimwood, K. & Lambert, S.B. (2012) Number and order of whole cell pertussis vaccines in infancy and disease protection. *JAMA* **308**: 454–456
- Sujeetha, M., Sharmila, S., Jayanthi, J. & Ragunathan, G. (2015). Antioxidant property of some extracts derived from the mud crab, scylla serrate. *International Journal of Phytopharmacology*, **6** (2):111-113.
- Suneeta kumara (2014) Extraction and characterization of chitin and chitosan from (Labeo rohit) fish scales. *Raffles bulletin of zoology*, **6**:482-489