

Reference bioelectrical impedance values for children of normal body mass index in southeastern Brazil

Abstract

Objective: The purpose of this study was to determine the reference values of the indexes bioelectrical impedance (BI) for children of normal body mass index in southeastern Brazil of middle-income country.

Methods: Two hundred eighty-one children with normal body mass index were included in the study (135 female and 146 male), aged 4 to 129 months, selected from federal public urban school in São Paulo, São Paulo, Brazil, where bioelectrical impedance values resistance (R) and reactance (Xc) values were measured in order to establish reference values of these parameters.

Results: The anthropometric variables, body mass index, z-scores and bioelectrical impedance parameters were evaluated. For both genders, the mean and standard deviation of anthropometric variables were: age (months): 73.42 ± 34.65 ; weight (kg): 23.5 ± 9.46 ; height (m): 1.16 ± 0.22 ; BMI(kg/m²): 16.65 ± 1.75 ; Xc(ohms): 63.92 ± 9.6 ; R(ohms): 749 ± 75.26 . For analysis, the children were stratified into three groups for each gender, being divided by ages: 4 to 23 months; 24 to 71 months and 72 to 129 months. Linear regression analysis showed R had a significant progressive decrease with age ($p=0.0003$) while Xc had a progressive increase ($p=0.0065$) with age. We analyzed by multiple regression the associations between R and Xc with anthropometric variables by age group in order to establish the reference values, confidence intervals and the tolerance limits for a new individual observation.

Conclusion: The BI reference values were established, in a field where there is a relative lack of publications, and we collected relevant information about resistance and reactance in a population of middle income setting that could be used in epidemiologic studies and could be used reference value in children with altered body composition.

Keywords: bioelectrical resistance, bioelectrical reactance, bioelectrical impedance analysis, children, fluids and electrolyte, dynamic, capacitive properties

INTRODUCTION

Formatted: Heading 1, Justified

Comment [U1]: Few lines about bioelectric impedance can be given as introduction

Comment [U2]: R and Xc measured by ? instrument name can be mentioned here by one word.

Comment [U3]: Key words dynamic and capacitive properties may not be specific to the abstract. You can consider to remove and add BMI

Formatted: Font color: Red

Bioelectrical impedance analysis (BIA) is an fast, and inexpensive method that has been widely applied to evaluate the body composition for over thirty years.^{1,2,3}

Bioelectrical analysis measure two bioelectrical vectors: resistance (R) and reactance (Xc). Resistance the is pure opposition of a biological conductor to the flow of an alternating current through the intra and extra-cellular ionic solution and it is inversely related to the dynamics of body fluids and body composition. For this reason, resistance tends to decrease when there is an increase in free fat mass (hydrophilic) and it tends to decrease when there is an increase in fat mass (hydrophobic).⁴

Reactance is related to the capacitance and it is associated with several types of polarizations and electrochemical gradients produced by cell membrane and tissue interfaces. These vector components originate impedance (Z) and the phase angle (PA), which is the angle formed between Z and R, calculated as the arc tangent of the relation Xc/R . PA is positively associated with Xc and negatively associated with R, and its variations are consequence of alterations of body compositions or in cellular membrane function.^{5,6}

The body bioelectrical impedance technique is useful in the analysis of body composition, as it allows health professionals to manage and prevent nutritional problems. Additionally, the growing interest in the study of body composition and its variations as a method of assessing nutritional status grows over the years as well as recognition of its importance for the assessment of healthy and sick individuals.^{7,8}

BI has a hypothetical inverse relationship to the body's volume and can be used in regression prediction models to estimate total body water (TBW). It is based on a bi-compartmental model, which divides the body into lean mass (LM) - high conductivity, a fact that reduces body resistance (R), and free fat mass (FM) - low conductivity that increases body resistance (R).⁸

Our interest in determining the reference values for a healthy pediatric population with Z-score indexes between ± 2 as well as normal body mass index is justified due to the fact that electrical bioimpedance is easy to perform, allowing non-prolonged training of the technician that will perform the method, and has already demonstrated consistent results, both in adults and children, for body composition estimates, when associated with anthropometry.^{5,7,8,9,10}

Comment [U4]: Explanation for hydrophilic and hydrophobic are same sentences here, just see the meaning and you may modify it.

Comment [U5]: Reference values of normal BMI can be mentioned here

There is a relative lack of publications in the field of bioelectrical parameters reference values on specific population such as low and middle income countries. For this reason, few studies are reported in order to properly assess nutritional individual deviations in relation to these population mean and to analyze the role of bioelectrical parameters on various outcomes in the clinical setting and epidemiological studies.

The purpose of this study was to determine the reference values of the bioelectrical impedance indexes for children of normal body mass index in southeastern Brazil as representative of middle-income country.

METHODS

Data were collected in healthy children aged 4 to 129 months at an federal elementary school in São Paulo city, Brazil. The children belonged to families that have the socioeconomic status of the majority of the Brazilian population, being in the middle class. The protocol was approved by the committee of ethics on research of the Universidade Federal de São Paulo and the school's authorities.

Study Population

Three hundred, twenty seven children of both gender were recruited after interviewing their parents and obtaining a signed written informed consent. The admission criteria for this study were: a) Children with z-score weight-for-height and body mass index between $\pm 2,0$ according to cutoff point as a discriminating nutritional disorders using as reference NCHS curves, b) fasting state major to 3 hours and c) no vigorous physical activity in the 24 hours prior to the tests. The exclusion criteria were: a) undernutrition [z-score ≤ -2], b) obesity [z-score $\geq +2$], c) acutely ill children, and d) children who were under medications.³

Comment [U6]: You may mention the word as after detailed explanation about the procedure instead of interviewing

Anthropometric measurement –

The anthropometric measurements were obtained by the principal investigator who was previously trained to perform the measurements. The anthropometric measurement procedures were undertaken in strict accordance with the methodology described in previously published papers.^{8, 9, 10}

We performed measurements for weight and height in triplicate and the average of these measurements was used. The body weight was measured to a precision of 0.1 Kg with an bean scale in children over 23.9 months of age. In children under 23.9 months, the body weight was measured to a precision of 0.01 kg using an electronic scale. The body-

height was measured by a stadiometer to a precision of 0.1 cm for all age groups. The children were measured without shoes and wearing underwear. The age, body weight and height were used to calculate the Z-score. We used the relationship weight-for-height (W/H index) for the nutritional assessment of the children over 23.9 months of age and for children under 23.9 months, the weight-for-age (W/A index) and weight-for-height (W/H index). The values obtained were compared to standard reference values.

We used version 1.02 of the ANTHRO program from the Nutrition Division of the Disease Control Center (CDC). To calculate the z score, comparisons were made between the z scores obtained with the curves of the National Center for Health Statistics (NCHS), using cutoff values to define the nutritional condition ± 2 z scores.

We determined the body mass indices (BMI) - weight divided by the square of height in meters - for each child, which were also compared with the NCHS values. Thus, only children with a Z score and BMI within the normal values established by the NCHS were included in the study.

Bioelectrical impedance measurements

Whole-body electrical resistance and reactance were measured with a bioelectrical impedance analyzer that measure resistance and reactance independently and separately. (Biodynamics model 310; Biodynamics Corporation, Seattle, WA) of alternate current at 800 μ A and 50 kHz in tetrapolar arrangement.

Oil was removed from the skin by cleaning it with alcohol. No direct contact was made with the child's skin during measurements, and the children were calm and relaxed¹⁴

For children under 18 months of age (where cooperation was more difficult), we made a cylindrical non-conducting cylindric plastic frame with the objective of positioning the children correctly, i.e, in dorsal decubitus with arms and legs separated and in abduction at 30 degrees from the trunk. That frame was not used with older children, and the supine positioning was maintained.

We positioned the electrodes in pairs on the right side of the body in the following anatomical positions: 1- Right hand: The current injector electrode was positioned in the middle of the dorsal surfaces of the hand proximal to the third phalangeal-metacarpal joint.¹¹ The detector electrode was placed 4 cm below the wrist (group 1) or medially between the distal bony prominences of the radius and ulna (group 2 and 3); 2- Right foot: the current injector electrode was positioned in the middle of the dorsal surfaces of the foot to the third metatarsal-phalangeal joint. The detector electrode was placed 4 cm below the

Comment [U7]: Mention the unit as Weight in kg

Comment [U8]: Remove instrument model name from bracket for the continuation of sentence.

Comment [U9]: 2 times cylindrical word is coming in same sentence

Comment [U10]: Why only right side is selected. Is there any supporting article says no difference in right and left measurements

Comment [U11]: Group 1,2,3 are mentioned here, but grouping is explained in results? Grouping can be mentioned here instead of results

ankle (group 1) or medially between the medial and lateral malleoli at the ankle (group 2 and 3). Before each test, the master power switch of the analyzer was turned off and on. After pressing the on key, the analyzer performs a self-test to check the internal calibration in accordance with the recommendation of the manufacturer.

Comment [U12]: Duration of test. How long electrode need to be placed in childrens body surface?

STATISTICAL METHODS

Descriptive analyses were expressed as mean, standard deviation and 95% confidence intervals (CI). The inferential statistical analyses were performed using GCM and REG procedures of the statistical software package SAS (Version 6.0). Bivariate correlations and stepwise maximum R^2 was performed by multiple linear regression analyses in order to evaluate the strength and variability of R and Xc with weight (W), height (H) by age and gender. A p value < 0.05 was considered statistically significant.

RESULTS

The variables were collected in 327 children. 46 children had to be excluded due to the following reasons: 6 were undernourished, fourteen were obese and 26 had other exclusion criteria. After exclusion, the study population consisted of 281 healthy children. Due to low number of children, we previously stratified the children in three age-group: a) 4 to 23 months, b) 24 to 71 months, and c) 72 to 129 months (group 3). The subject characteristics are presented in Table 1.

Linear regression analysis was performed to evaluate if age-group stratification was appropriate to study the variability of the resistance and reactance in relation to anthropometric variables. Figures 1 and 2 shows that the stratification was appropriate. In both graphics there are two inflection points, the first point at 23 months and the second at 71 months. These two points were interpreted as indicative of the resistance and reactance variations imposed by growth and development. The straight lines were significantly different for the resistance ($p=0.0003$) and reactance ($p=0.0065$).

Correlation between bioelectrical impedance components and anthropometric variables

Multivariate regression models were used to analyze the correlations between resistance and reactance with anthropometric variables. The purpose of these models was to establish confidence intervals for R and Xc for normal children and tolerance intervals for a new observation. Tables 2 and 3 lists the multivariable regression equations. Due to

the small number of children for each gender in age group 1, one model was adjusted for both genders.

Pearson's correlation coefficient between anthropometric variables and bioimpedance vector components are described in table 4. Weight and height were negatively correlated with resistance in all age groups. The reactance was positively correlated with weight and height in females.

Boys and girls did not differ in age, body weight and body height but girls had a higher resistance than boys in groups 2 and 3. This difference in body resistance between boys and girls was not found in the infants (group 1). Reactance increases with age, having few variations between genders (table 5).

The regression models were used to estimate R and Xc mean and 90% to 99% confidence intervals (CI) for age group and gender. In addition, we used the regression models to estimate the values expected of the impedance vectors and the tolerance limits 90% to 99% for a new observation.

DISCUSSION

Bioelectrical impedance analysis (BIA) is considered a good method for estimating body composition in the epidemiologic studies and at the bedside. It is safe, non-invasive, reliable, rapid, inexpensive, portable, and it allows to repeated measures could be taken quickly.⁵

We studied separately R and Xc components grouping by age depending on the sample size and gender. The three age groups adopted were based on Waterloo et al.¹² stratification sampling criteria, that clustered the children into relatively homogenous subgroups by age.

In addition, the skin electrodes were placed on anatomical position and those electrodes had their patches width reduced in young children because there is a minimal distance required to avoid interactions between electrodes. These criteria adopted by us were similar to other studies in children where: 1) similar groups of children were considered; 2) skin electrodes were placed in accordance with the child's age; 3) the children were separated in age-groups; 4) Xc vector component was not neglected; and 5) age-related variability was found in these studies.^{1,13,14}

The measures demonstrated that resistance measurements were substantially higher in all age groups than those reported for adults. In healthy American adults, that

Comment [U13]: In which group you gender significance

Comment [U14]: This reference of grouping may be mentioned under methodology.

Comment [U15]: Electrote placement reference could be there in methodology, need not be mentioned 5 points as criteria which all says about age factor

means range from 432 to 485 ohms for men and 551 to 587 ohms for women and in healthy Brazilian adults 552 ± 100 ohms in both genders. Our study demonstrated that resistance values in young children were higher than older children, and these results are similar to those in the previous studies.^{13,5,14}

We observed variability of the resistance and reactance parameters with growth in our study, reinforcing the importance of the reference values of R and Xc by age or age-group and gender in healthy populations of children. The variability of parameters might be reflecting changes during growth as does intra and extra-cellular fluid distribution, cell growth and changes in body mineral and electrolytic content, therefore, reflecting the variability of fluids and body composition in children.

Comment [U16]: Consider the meaning of the sentence

The study showed that resistance decreases with age, which might be because the muscular mass of the limbs increases with growth. These observations reinforce the concept whereby in the infants and toddlers, arms and legs represent a body area with small diameter and length, therefore the resistance is high. With growth, there is an increase of the diameter and length of the limbs, and R decreases due to an increase in the cross-sectional area of the extremities. These observations are according to simple body-composition models where the appendicular skeletal muscles are the primary electrical conductor.^{16,19,20}

We observed differences in the reactance among the three study-groups. This might be due to the differences of capacitance properties of the tissue interfaces and cell membranes. Theoretically, Xc variation among healthy individuals could be due to differences in the capacitive behavior of the tissues associated with variability of the cell size, membrane permeability or intracellular composition during growth.^{21,22} An increase of interstitial fat (anhydrous, meaning that fat is hydrophobic) during maturation reduces both the tissue interface permeability and cell membrane interface permeability, producing an increase in reactance in a critical fixed frequency.²¹

The variability of R and Xc might be explained also by variations that include more and less conductive matter, body temperature, tissue composition, fluid distribution, ionic concentration, nature of fat, as well as anisotropic effects of muscle fibers. These physiological and structural as well as technical factors affect the measurement of both bioelectrical impedance vector components, R and Xc.^{3,4,6}

The limitations of this study is that the sample cannot be considered representative of all millions of Brazilian children because there is difference in the nutritional status

among specific Brazilian regions depending on the socioeconomic levels of population in each region of Brazil. In order to minimizing populational bias, the epidemiologic procedure performed in this study consisted selecting a school with children from families with middle income resources. Our study is the first and only one study already realized in Brazil to establish bioelectrical impedance vectors reference values in children for several age groups and gender.

In summary, we established the normative bivariate 90% to 99% confidence intervals for the mean impedance indexes by group and gender and the bivariate predictive values 90% to 99% tolerance limits for new individual measurements of the resistance and reactance in healthy Brazilian children. Further, changes in resistance and reactance with age are well-established. Our findings add substantial information in a field with relative lack of publications.

APPENDIX

Free software is available to calculate mean estimated value and confidence intervals and tolerance intervals for an additional observation of the bioimpedance vector components from cristina.mangia@unifesp.br

All tables with individual values (weight, height, BMI, z-score, percentiles, resistance, reactance and phase angle) and regression formulas are available for consult at Dr. Cristina Mangia by e-mail.

REFERENCES

1. Davies PSW.1993. Body composition assessment. Arch Dis Child 69: 337- 338.
2. Hammond J, Rona RJ and Chinn S.1994. Estimation in community surveys of total body fat of children using bioelectrical impedance or skinfold thickness measurements. Eur J Clin Nutr 48:164 -171.
3. Lukaski HC, Bolonchuck WW, Hall CB and Siders WA.1986. Validation of tetrapolar bioelectrical impedance method to assess human body composition. J Appl Physiol 60:1327-1332.
4. Lukaski HC.1996. Biological indexes considered in the derivation of the bioelectrical impedance analysis. Am J Clin Nutr 64 (Suppl):397-404.
5. Mattar JA and the Brazilian Group for Bioimpedance Study.1996. Application of total body bioimpedance to the critically ill patients. New Horiz 4: 493-502.

6. Barbosa-Silva MCG, Barros AJD, Wang J, Heymsfield SB and Pierson Jr RN. Bioelectrical impedance analysis: population reference values for phase angle by age and sex. *Am J Clin Nutr* 2005;82:49-52.
7. Walker SP, Grantham-Mc Gregor S, Powell C, Fletcher P, Himes JH. 1990 Bioelectric impedance, anthropometry and body composition in stunted and non-stunted children. *Eur J Clin Nutr* 44:763-768.
8. Fagundes U, Oliva CAG, Fagundes-Neto U. Evaluation of the nutritional status of Indian children from Alto Xingu, Brazil. *J Pediatr (Rio J)* 2002; 78 (5): 383-8
9. Organización Mundial de la Salud.1983. Medición del Cambio del Estado Nutricional. Directrices para Evaluar el Efecto Nutricional de Programas de Alimentación Suplementaria Destinados a Grupos Vulnerables, pp.1-105. OMS, Ginebra.
10. Waterloo JC, Buzina R, Keller W, Lane JM, Nichaman MZ, Tanner JM.1977. The presentation and use of height and weight data for comparing the nutritional status of group of children under age of 10 years. *Bull World Health Organization* 55: 489-498.
11. De Palo T, Messina G, Edefoni A, Perfumo F, Pisanello L, Peruzzi L, Di Iorio B, Mignozzi M, Vienna A, Conti G, Penza R, Piccoli A. 2000. Normal values of the bioelectrical impedance vector in children and puberty. *Nutrition* 16:417- 424.
12. Piccoli A, Fanos V, Peruzzi L, Schena S et al. 2002. Reference values of the bioelectrical impedance vector in neonates in the first week after birth. *Nutrition* 18:383-387.
13. Lukaski HC.1999. Requirements for clinical use of bioelectrical impedance analysis (BIA). *Ann New York Acad Sci* 873:72-76.
14. Baumgartner RN, Chumlea WMC, Roche AF.1987. Associations between bioelectric impedance and anthropometric variables. *Hum Biol* 59: 235- 244.
15. Fuller NJ, Ellia M.1989. Potential use of bioelectrical impedance of the "whole body" and of body segments for the assessment of body composition: comparison with densitometry and anthropometry. *Eur J Clin Nutr* 43:779-791.
16. Organ LW, Bradhan GB, Gore DT and Lozier SL.1994. Segmental bioelectrical impedance analysis: Theory and application of a new technique. *J Appl Physiol* 77:88-112.

17. Heymsfield SB, Wang ZM, Visser M et al. Techniques used in the measurement of body composition: an overview with emphasis on bioelectrical impedance analysis. *Am J Clin Nutr* 1996; 64 (suppl): 478S-84S.
18. De Lorenzo A, Andreoli A, Matthie J and Withers P.1997. Predicting body cell mass with bioimpedance by using theoretical methods: a technological review. *J Appl Physiol* 82:1542-1558.
19. Deurenberg P, Wetstarte JA, Hautvast JGAJ, van der Kooy K.1991. Is the bioelectrical impedance method valid? *Am J Clin Nutr* 53: 179-180
20. Gregory JW, Greene SA, Scrimgeour CM and Rennie MJ.1991. Body water measurement in growth disorders: a comparison of bioelectrical impedance and skinfold thickness techniques with isotope dilution. *Arch Dis Child* 66:220-222.

Table 1. Demographic characteristics of children

		Age (months)		Height (cm)		Weight (Kg)		BMI		R (Ohm)		Xc (Ohm)	
Group		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	Both gender*	10.3	4.6	71.3	6.3	8.7	1.9	8.71	1.85	801	96	51	8
2	M (n=39)	51.7	13.6	106.6	10.9	18.4	4.0	16.02	1.35	750	64	63	9
	F (n=37)	56	13.1	106.5	9.2	18.4	3.1	16.10	1.17	765	64	65	8
3	M (n=94)	97	15.8	130.3	9.2	28.8	6.4	16.79	1.86	720	60	67	8
	F (n=73)	98.8	15.8	131.5	10.6	29.8	7.8	16.92	2.12	750	75	67	8
Total	M (n=146)	76.48	31.96	119	0.20	24.35	8.45	16.67	1.72	729.53	62.98	64.21	9.28
	F (n=135)	70.13	37.18	112	0.28	23.95	14.79	16.63	1.80	770.56	81.78	63.61	9.98
Total	n=281	73.42	34.65	116	0.23	23.5	9.46	16.65	1.75	749	75.26	63.92	9.6

Group 1 = 4 months \leq age \leq 23 months; Group 2 = 24 months \leq age \leq 71 months; Group 3 = 72 months \leq age \leq 123 months.

* Male: n=13 and Female: n=25

Height, cm; Weight, Kg; BMI (Body mass index), kg/m^2 ; R, resistance in ohm (Ω); Xc, reactance in ohm (Ω); SD, standard deviation

Table 2. Prediction of the resistance according to age, body weight, body height for three study-groups by age and genders

Group/sex		N	a_0	a_1	a_2	r^2	SEE	p
G1	Both	38	600.44 ^a	10.86 ^{ns}	-65.89 ^c	0.41	75.39	0.0001
	Male	39	636.82 ^d	3.22 ^{ns}	-12.63 ^c	1.14	60.75	0.07
G 2	Female	37	608.83 ^d	4.00 ^{ns}	-14.73 ^{ns}	0.11	62.71	0.14
	Male	94	467.48 ^d	3.96 ^c	-9.14 ^d	0.29	51.37	0.0001
G 3	Female	73	268.46 ^f	6.50 ^d	-12.54 ^d	0.39	59.43	0.0001

$R = a_0 + a_1 * H + a_2 * W$; R = resistance (ohm); H= Height (cm); W = weight (Kg); Group 1 = 4 months \leq age \leq 23 months; Group 2 = 24 months \leq age \leq 71 months; Group 3 = 72 months \leq age \leq 123 months. ^ap<0.02; ^cp<0.002; ^dp<0.0001; ^ep<0.05; ^fp<0.006. NS = non- significant.

Table 3. Prediction of reactance according to age, body weight, body height for three age study-groups and gender

Group	Gender	N	b_0	b_1	b_2	r^2	SEE	p
1	Both	38	62.92 ^a	-0.24 ^{ns}	0.60 ^{ns}	0.008	7.43	0.87
2	Male	39	27.17 ^{ns}	0.52 ^{ns}	-1.16 ^{ns}	0.085	8.21	0.21
	Female	37	13.15 ^a	0.71 ^a	-1.27 ^{ns}	0.204	6.87	0.02
3	Male	94	50.73 ^a	0.21 ^{ns}	-0.42 ^{ns}	0.031	7.91	0.23
	Female	73	44.12 ^a	0.33 ^{ns}	-0.73 ^a	0.147	7.16	0.0038

$Xc = b_0 + b_1 * H + b_2 * W$; Xc = reactance (ohm); H= Height (cm); W = weight (Kg); Group 1 = 4 months \leq age \leq 23 months; Group 2 = 24 months \leq age \leq 71 months; Group 3 = 72 months \leq age \leq 123 months. ^ap<0.02; NS = non- significant.

Table 4. Correlation of resistance and reactance with body weight, body height for age study-groups and gender

	Group 1	Group 2		Group 3	
	Both genders	Male	Female	Male	Female
Resistance (Ohm)					
Height (cm)	-0.48 ^a	-0.22 ^b	-0.07 ^b	-0.22 ^b	-0.24 ^c
Weight (Kg)	-0.60 ^a	-0.30 ^e	-0.20 ^b	-0.44 ^a	-0.44 ^a
Reactance (Ohm)					
Height (cm)	-0.07 ^b	0.08 ^b	0.46 ^d	-0.04 ^b	-0.04 ^b
Weight (Kg)	-0.05 ^b	-0.00 ^b	0.32 ^c	-0.12 ^b	-0.12 ^b

^ap<0.001; ^b p= NS; ^c p<0.05; ^dp<0.005; ^ep<0.06.); Group 1 = 4 months ≤ age ≤ 23 months; Group 2 = 24 months ≤ age ≤ 71 months; Group 3 = 72 months ≤ age ≤ 123 months.

Table 5. Estimates and 95% tolerance intervals for resistance and reactance for three age-study groups

Group	Gender	Mean	Lower 95% TL	Upper 95% TL
Resistance (Ohm)				
1	Both	880	707	1053
2	Female	765	744	787
	Male	748	728	769
3	Female	749	732	767
	Male	721	708	733
Reactance (Ohm)				
1	Both	51	48	54
2	Female	65	63	68
	Male	63	60	66
3	Female	67	65	69
	Male	67	65	68

Group 1 = 4 months ≤ age ≤ 23 months; Group 2 = 24 months ≤ age ≤ 71 months; Group 3 = 72 months ≤ age ≤ 123 months.

TL= tolerance limits; Lower 95% TL= lower limit; Upper 95% TL= TL upper limit.

Mean estimated value was calculated using the regression models presented in tables 2 and 3. Tolerance limit for the estimated mean were calculated with the expression:

$$\bar{y}_0 \pm zs \sqrt{x_0^T (X^T X)^{-1} x_0}$$

Where $x_0 = (W, H)$ in the estimated regression equation and X the model matrix. Z is the corresponding normal distribution percentile and s the standard error estimate.

Figure 1– Relationship between measured (white) and predicted (black) Resistance values according to age. The regression line predicted for the three age groups studied were significantly different ($p = 0.0003$).




Figure 2 – Relationship between measured (white) and predicted (black) Reactance values according to age. The regression line predicted for the three age groups studied were significantly different ($p = 0.0065$).

