

28 created the vast Niger Delta at its confluence with the Atlantic Ocean's Guinea and Benguela
29 Currents [3].

30 Nigeria is enormously blessed with abundant human, agricultural, petroleum, gas and other
31 unexploited solid mineral resources [4]. However, due to the political instability experienced
32 since her independence in 1960 from British rule, it has experienced decades of political
33 instability; ~~therefore~~ ~~therefore~~, creating social tension and an unpredictable market for
34 businesses [5]. It is worthy to mention that the Nigerian economy is profoundly reliant on the
35 oil and gas sector and classed as the fifth largest oil exporter to the United States [6]. The Niger
36 Delta region of Nigeria literally covers about 36,000 kilometers (14, 000 square miles) of
37 marshland, creeks, tributaries, and lagoons which drain the Niger River into the Atlantic [7].
38 About 12,000 square kilometer of Niger Delta is fragile mangrove forest, and this is arguably the
39 largest mangrove forest in the world. It is also the largest wetland in the world. The biodiversity
40 is very high and the area contains diverse plant and animal species, including many exotic and
41 unique flowers and birds. Implied in this ecology is that the Niger Delta is an easily dis-
42 equilibrated environment. The environment is mostly salt water and associated with shortage
43 of arable land and freshwater. Furthermore, transportation through this ecosystem is very
44 difficult.

45 Today, crude oil is produced in nine States in Nigeria, namely, Rivers, Bayelsa, Delta, Edo, Imo,
46 Abia, Akwa-Ibom, Cross-River, and Ondo. Due to this fact and other political reason, the
47 present-day Niger Delta is technically made up of these nine States as depicted in Figure 2.6,
48 and covers an area of about 41,000 square miles (106,189.50 km²) and harbours Nigeria's
49 proven oil and gas reserves. According to [8], 70% of Nigeria's proven gas reserves are situated
50 on land, while the rest 30% are offshore.

51 Nigeria's Niger Delta is characterized by high biological diversity, abundant natural resources,
52 and extreme poverty. A survey of current knowledge on the biological diversity of the Niger
53 Delta reveals striking global significance across the full range of biological diversity at the
54 genetic, species and ecosystems levels. Biological diversity is the variety of the world's plant
55 and animal life (in this case, the Delta's), including their genetic diversity and the assemblages
56 they form. The Niger Delta region of Nigeria is one of the world's largest wetlands and includes

Comment [KA3]: Where is Figure 2.6, please correct ?

Comment [KA4]: What is this 8

57 by far the largest mangrove forest in Africa. Its biological diversity is of global significance.
58 Within the extremely valuable ecosystem, oil activities are widespread [9]. Particularly, the
59 community which is Eleme is a community in Rivers State and it's one of the oil producing and
60 agro-ecological areas in the Niger-Delta region of Nigeria, a region with abundant natural
61 resources including good weather and fertile land for agriculture.

62 Although the level of agricultural production in that region is very low given the abundant
63 resource endowment, it is the largest oil producing zone in the country. It is the base of
64 Nigerian

65 oil and gas industry, generating over 90% of the nation's economy [10]. Oil exploration and
66 activities have been concentrated in this Niger-Delta region which has over 1000 production oil-
67 wells and over 47,000 km of oil and gas flow lines [11]. These negative impacts of this oil
68 activities includes destruction of wild life, loss of fertile soil, pollution of air and water and
69 damage to the ecosystem of the host communities [12]. The ecological problems observed as a
70 result of oil spill include a brownish vegetation and soil erosion, diminishing resources of the
71 natural ecosystem, fertile land turned barren and adverse effect on the life, health and
72 economy of the people [13].

73 The differences in field management, land use conversion, which involves change in biomass
74 production and nutrient cycling, have influence on soil properties [14]. Change in land use from
75 agriculture to forest brought the development of a large tree biomass and increased the
76 availability of plant nutrient [15]. This type of conversion increased soil organic carbon,
77 microbial biomass and potential nitrogen mineralization rate and reduced the soil bulk density
78 [16]. Land use induced changes in nutrient availability and may influence secondary succession
79 and biomass production, reduce crop production and environmental quality. The changes
80 directly affect some physical chemical and biological properties, such as water retention
81 availability, nutrient cycling, plant root growth and soil conservation [17]. These soils are also
82 known to possess unique morphological characteristics that are strongly influenced by
83 temporary or permanent water saturation and adopted vegetations. This study therefore is to
84 assess the soil heavy metal contents of some land uses in Imo State.

85

Comment [KA5]: Correct the typos

86 2.0 MATERIALS AND METHODS

87 2.1 STUDY AREA

88 The study was carried out in Obinze area of Imo State which lies between latitudes $05^{\circ} 21^1$ and
89 $05^{\circ} 42^1$ N and longitudes $07^{\circ} 48^1$ and $06^{\circ} 53^1$ E. The region consists of tropical rainforest zone
90 with average annual rainfall distribution of 2,250-2800mm. The annual temperature ranges 26-
91 30° C with annual relative humidity range of 85-90% [18].

92 Human activities such as continuous cropping, grazing and bush burning has transformed the
93 natural forest of the area into secondary and grassland soils, but there are some scattered
94 distributions of forest lands in the area. The identified land uses were grass land, continuously
95 cropped land and forest land.

96

97


98

99

100


101

Comment [KA6]: Typos, please write in standard way



103 Figure 1: Map of Imo State Showing the Study Area

104



Comment [KA7]: Image is not very clear.

105

106 Figure 2: Map of the Study Area (-Ohaji/Egbema L.G.A in Imo State).

107

108

109

110

111

112

113

114 2.2 FIELD STUDY

115 Three mapping units were chosen to represent soils that occurred under grass land,
116 continuously cropped land and forest land. In each of the mapping units, representative profile
117 pits were dug up to 90cm delineated and soil samples collected from each of the pedogenetic
118 horizons.

119 2.3 LABORATORY ANALYSIS

120 Routine laboratory analyses were conducted after collection of samples. Soil pH was
121 determined by electrometric method as described by IITA (2010). The walkley and black
122 methods as described by [19] were used in the analysis of organic carbon. Total nitrogen was
123 analyzed using the procedures as described by [20].

124 The Bray 1 method as described by IITA (2010) was used for extractable phosphorus,
125 exchangeable based were determined from the soil samples through normal ammonium
126 acetate solution [21]. The EDTA titration method was used to determine calcium (Ca) and
127 magnesium (mg), while flame photometer was used in the determination of sodium (Na) and
128 potassium (K). The cation exchangeable capacity (CEC) was determined by ammonium acetate
129 saturation method [21]. The electrical conductivity (EC) was determined by measuring the
130 electrical resistance between parallel electrodes immersed in the soil samples using electrical
131 conductivity meter.

132

133

Comment [KA8]: What is IITA?. Describe in full when comes first time in the test body.

Comment [KA9]: Proper noun must be begun with capital letter

Comment [KA10]: Complete the sentence

Comment [KA11]: Complete the sentence

Comment [KA12]: Put full stop

Comment [KA13]: Put full stop

134 The results of some chemical properties of the soils were shown in Table 1 which represented
 135 the grass land, continuously cropped and forest land soils. The soils across the pedons were
 136 generally acidic. The grass land and continuously crop.

137 2.4 HEAVY METAL DETERMINATION

138 Heavy metals like Mn, Fe, Cu, and Zn were determined using double acid method of extraction
 139 and extraction acid read out with AAS.

140 The samples were mixed gently and homogenized and sieved through 2 mm mesh - sieve. The
 141 samples were first dried, and then placed in electric oven at a temperature of 40 °C
 142 approximately for 30 minutes. The resulting fine powder was kept at a room temperature for
 143 digestion.

144 2.4.1 Digestion of Soil Samples

145 lg of the oven dried sample was weighed using a top loading balance and placed in 250ml
 146 beakers separately to which 15ml of aqueous regia (35% HC1, and 70% high purity HNO₃ in
 147 3:1 ratio) was added. The mixture was then digested at 70% till the solution became
 148 transparent The resulting solution was filtered through whatman-Whatman filter paper no
 149 42 and into a 50ml dilute 50ml volumetric flask and diluted to mark volume using deionized
 150 water and the sample solution was analysed for concentration of Cu, Pb, Mn, Zn and Fe
 151 using an atomic absorption spectrophotometer (Perkin-ElmerAAnalyst400).

Comment [KA14]: What is lg, please rectify.

Comment [KA15]: What is this. Perhaps it is aqua regia

Comment [KA16]: HC1 ?? rectify

Comment [KA17]: HNO₃ rectify.

152 2.4.2 Analysis of Soil Samples

153 AAS A Analyst 400 model was used in determining the content of metals in the previously digested
 154 soil samples. The nitrous oxide, acetylene gas and compressor were fixed and compressor
 155 turned on and the liquids trap blown to rid of any liquid trapped. The extractor and AAS control
 156 were turned on. The slender tube and nebulizer piece were cleaned with purifying wire and
 157 opening of the burner was cleaned with an arrangement card. The worksheet of AAS
 158 programming on the joined PC was opened and the empty cathode light embedded in the light

Comment [KA18]: What is AAS, describe in full at first place

159 holder. The light was turned on, beam from cathode adjusted to hit target zone of the
160 arrangement card for ideal light throughput, at that point the machine was touched off. The
161 fine was set in a 10ml graduated chamber containing deionizer water and yearning rate was
162 estimated. The analytical blank was prepared and a series of calibration solutions of known
163 amounts of analyte element (standard) were made. The blank and standards were atomized in
164 turn and their responses were measured. A calibrator graph was plotted for each of the
165 solutions after which the sample solutions were atomized and measured. The various
166 metals concentration from the solution were determined from the calibration based on the
167 absorbance obtained for the unknown samples

168 **2.5 STATISTICAL ANALYSIS**

169 Coefficient of Variation (CV) was used to estimate the degree of variability existing among land
170 uses in the study area as outlined by wilding (1985). The means were separated using least
171 significant difference (LSD) test at 0.05 level of significant incorporated in the statistical analysis
172 system (SAS) package of 9.1 versions (2006).

173 **3.0 RESULTS AND DISCUSSIONS**

174 The results of some chemical properties of the soils were shown in Table 1 which represented
175 the grass land, continuously cropped and forest land soils. The soils across the pedons were
176 generally acidic. The grass land and continuously cropped were slightly acidic. The forest land
177 was medium acidic with a mean pH value of 5.65. Available phosphorus was high in grass land
178 and continuously cropped with mean values of 12.20mgkg^{-1} and $12, 4\text{mgkg}^{-1}$ (Table 1). Organic
179 matter and total nitrogen decreased down the profile in all land uses and were low being less
180 than critical limits of $< 1\%$ for OM and $< 0.9\text{g/kg}$ for N as described by [22-26].

181 In this study, grass land soils organic matter decreased from 0.64 to 0.25, continuous cropped
182 land from 0.55 to 0.21 while in forest land from 0.96 to 0.66%. The organic matter were low in
183 three land studied, but continuously cropped soils were the lowest.

184 Erosion, leaching, and increased intensive agriculture in the area may have depleted the
185 nutrient reserve of the soils. The available phosphorus distributions were very high at the
186 surface horizons and decrease down the profile in all soils studied. The chemical indicator used
187 with the assigned relative weight is shown in this investigation. The soil pH is a critical factor in
188 crop production as it affects the mobility of many pollutants in the soil by way of influencing the
189 rate of their biochemical breakdown, solubility and absorption to soil colloids [27] The results
190 showed moderate acid condition in the continuously cropped soils and medium acid in grass
191 land and forest land respectively. The implications of these are that, the nutrient availability of
192 the soils may be affected. High pH in relation to the forest land soils could cause the released of
193 some toxic amounts of aluminum into the soils [28]. This suggests that these soils were low in
194 nutrient elements composition especially the continuously cropped soils and needs to
195 constantly be supplemented to support productivity.

196 Low values of exchangeable sodium and electrical conductivity (EC) associated with the three
197 land use soil are major soil chemical potentials of the soils as most tolerable tropical plants can
198 be cultivated in them. The organic matter content in the grass land and continuous cropped
199 land were low with mean value of 0.43gkg^{-1} , and 0.41gkg^{-1} . The forest land was higher in
200 organic matter content with mean value of 0.82gkg^{-1} . The low organic matter content
201 associated with the grass land and continuous cropped land were due to the intensive
202 cultivation of the soils, and seasonal bush burning which are more of animal farming ritual in
203 the study areas. The moderate to high organic matter associated with the forest land has been

204 attributed to microbial activities of the soils due to good aeration and decomposition rate of
205 flitters or litters, [24].

206 Heavy metals (Pb, Cu, Mn, Zn and Fe) contents in the three land use soils shows that the
207 concentrations of Cupper (Cu) and Manganese (Mn) were moderate and were below WHO
208 permissible limit. Cu with mean value of 33.4 mg/kg, Mn with mean value of 18.2mgkg, for
209 grass land and continuous cropped land 32.9mg/kg, 17.6mg/kg, while forest land with mean
210 values of 30.7 mg/kg, 17.8mg/kg of Cu and Zn respectively. Cu, Ma and Zn concentrations in
211 grass land and continuous cropped land were significantly higher ($P < 0.05$) than the Fe
212 concentration has a mean value of 1.6mg/kg, while continuous cropped land with mean value
213 of 1.5mg/kg and forest land with a higher mean value of 7.8mg/kg. However the mean value of
214 Mn, Fe in the land use soils was at tolerable limit and Pb was not found in this study while Zn
215 exceeded the maximum tolerable limit in soil for crop production according to [29].

216

217

218

219

220

221

222

223

224

225

226

227

228 **Table 1: SOME CHEMICAL PROPERTIES OF LAND USE SOILS**

Soil depth (CM)	Soil pH	Om	N (mgkg ⁻¹)	P (mgkg ⁻¹)	Ca	Mg	K (mgkg ⁻¹)	Na	CEC	EC dsm ⁻¹
Grass land										
0-15	5.64	0.64	0.84	12.74	2.07	0.84	0.18	0.24	6.57	0.93
15-30	5.52	0.49	0.80	12.64	2.06	0.70	0.18	0.24	6.34	0.62
30-60	5.43	0.35	0.76	12.20	2.05	0.65	0.17	0.13	6.32	0.52
60-90	5.13	0.25	0.40	11.23	2.04	0.55	0.16	0.12	5.64	0.42
Mean	5.43	0.43	0.70	12.20	2.06	0.69	0.17	0.18	6.22	0.50
CV%	12.86	10.18	22.30	23.18	10.84	44.77	14.12	23.45	15.15	3.64
Continuous cropped land										
0-15	5.62	0.55	0.88	12.21	1.67	1.07	0.17	0.24	6.89	0.48
15-30	5.53	0.48	0.84	12.20	1.52	1.04	0.16	0.20	6.75	0.47
30-60	5.22	0.38	0.84	12.08	1.45	0.76	0.14	0.16	6.70	0.44
60-90	5.14	0.21	0.83	12.05	1.43	0.74	0.11	0.15	5.45	0.44
Mean	5.38	0.41	0.85	12.14	1.52	0.90	0.13	0.19	6.45	0.46
CV%	13.08	10.64	20.20	20.57	13.03	16.34	19.86	20.54	7.62	13.78
Forest land										
0-15	5.83	0.96	0.98	12.74	4.73	2.10	0.17	0.86	12.32	5.62
15-30	5.64	0.88	0.98	12.68	4.72	2.07	0.15	0.75	12.09	5.03
30-60	5.61	0.76	0.93	12.66	4.72	2.07	0.14	0.70	11.84	4.60
60-90	5.52	0.66	0.89	12.64	4.65	2.04	0.11	0.65	11.25	4.40

Mean	5.65	0.82	0.96	12.68	4.71	2.07	0.14	0.74	11.88	4.91
CV%	12.12	10.81	11.14	25.28	21.01	10.68	19.0	11.77	3.16	16.50

229

230 **CEC = Cation exchange capacity, EC = Electrical Conductivity, OM = Organic matter**

231 The results on heavy metal contents in different land use soils were shown in Table 2. At the
 232 grass land soils ~~Copper~~ (Cu) decreased from 36.1-31.0, at the continuously cropped it
 233 decreased from 35.6 – 30.1 and forest land, Cu decreased from 30.1 – 29.8mg/kg, means were
 234 2.7, 3.2 and 0.2 mg/kg for GL, CC and FL respectively. Mn had means of 3.5, 1.9 and 0.6 mg/kg
 235 in GL, CC and FL, respectively. Although the Cu and Mn were low in the three land soils
 236 investigated, forest land soils were however very low in Cu and Mn. Zn and Fe decreased from
 237 56.8 – 48.7 and 2.2 – 1.2 mg/kg in GL. In CC land 56.2 – 45.4 and 1.5 – 1.3 mg/kg decreased.
 238 Similar trend also occurred in forest land. There was no Pb found in the three land uses
 239 investigated.

240

241

242

243

244

245

246

247

248

249

250 Table 2: **HEAVY METAL CONTENTS IN DIFFERENT LAND USE SOILS**

251

Soil Depth (cm)	Heavy Metals in mg/kg (WHO/FAO Limit)				
	Pb(85)	Cu (36)	Mn (20)	Zn (50)	Fe(0.5-10)
Grass land					
0 – 15	ND	35.1	20.4	56.8	2.2
15 – 30	ND	33.4	19.4	50.1	1.7
30 – 60	ND	32.9	17.8	50.0	1.3
60 - 90	ND	31.0	15.1	48.7	1.2
Mean	0.	33.4	18.2	51.4	1.6
C.V%	0	2.7	3.5	3.1	0.9
Continuous cropped land					
0-15	ND	35.6	18.3	56.2	1.5
15-30	ND	34.7	18.2	52.1	1.5
30-60	ND	31.2	17.5	48.7	1.5
60-90	ND	30.1	16.2	45.4	1.3
Mean	0	32.9	17.6	50.6	1.5
C V%	0	3.2	1.9	5.6	0.2
Forest land					
0-15	ND	32.5	18.0	56.1	7.8
15-30	ND	30.3	17.8	54.4	7.4
30-60	ND	30.1	17.7	49.8	6.5
60-90	ND	29.8	17.6	47.4	6.3
Mean	0	30.7	17.8	51.9	7.0
C.V%	0	0.8	0.6	4.2	0.1

252

253 Pb = Lead, Cu = Copper, Mn = Manganese, Zn = Zinc, Fe = Iron, ND = Not detected, CV =

254 Coefficient of variation

255

256

257

258

259 4.0 CONCLUSION

260 The investigated soils were sandy with highest occurring at the grass lands. The chemical quality
261 assessment of the soils using the soil management framework (SMAF) showed that the grass
262 land and continuously cropped soils were low to moderate in nutrient reserves and slightly to
263 medium acidic. The forest land soils have good chemical qualities in terms of organic matter;
264 total nitrogen and some high electrical conductivity are among chemical properties. However,
265 the use of both organic and inorganic fertilizers and efficient liming are some of the measures
266 that can improve the efficient use of the soils for crop production. However the mean value of
267 Mn, Fe in the land use soils was at tolerable limit and Pb was not found in this study while Zn
268 exceeded the maximum tolerable limit in soil for crop production according to [30] The main
269 soil factors having impact on the mobility of these elements and the total content in the soil are
270 the cation exchange capacity (CEC,) pH of the soil solution, mechanical composition of the soil
271 humus content and the interaction (Competition) among them [31]. The high concentrations of
272 these heavy metals (Cu, Mn, Zn and Fe) occurring in the land use soils strongly indicate that
273 heavy metals pollution of these soils were due to mineralization of these heavy metals as a
274 result of mining activities in the areas[32-35].

275

276

277

278

279

280

281 **REFERENCES**

- 282 1. Andrews, S. S., Karlen, D. L. and Canbandella, C.A. (2008). The soil management
283 assessment framework: A quantitative soil quality evaluation methods. *Soil Sci. Soc. An.*
284 1.68:1945-1962.
- 285 2. Brady, A. C. and Weil, R. E. (2007). The Nature and properties of soil (13th edition)
286 prentice Hall of India, private limited, New Delhi, India Pp. 976-978.
- 287 3. FAO-Food and Agriculture organization.(1983). Guideline land evaluation for rain forest
288 agriculture soils. Bulletin No. 52. Rome.
- 289 4. Flororunso, O. A., Ohu, I. O., and Adeniji, F. A. (1988).The role of soil spatical variability
290 investigation in the management of the chad Basin Vertisols of NE Nigeria.*Soil Tech.*
291 1:149-156.
- 292 5. IITA-International Institute of Tropical Agriculture.(2010). Automated and semi-
293 automated Analysis.Manual series No. 12 IITA. Ibadan, Nigeria. Pp 72.
- 294 6. LIU, B., Morkved, P.T., Frostegard, A., Bakken, L.R. (2010). Denitrification gene pools,
295 transcription and Kinetics of MO, N₂O and N₂ production as affected by soil pH, FEMS
296 *Microbiology Ecology.* 72:407-417.
- 297 7. Nelson, D.W. and Sommers, L.E. (2008).Total carbon, organic carbon and organic
298 matter. In page; A.L Miller R.H and Keeney, D.R. (ed) Methods of soil Analysis. Part 2
299 Agron.Monograph.NO 9.ASA and SSSA. Madison, W.L. USA.
- 300 8. NMA – Nigerian Metrological Agency., (2011). *Annual Bulletin*, 34:64-74.
- 301 9. Nymore, A.W. (2007). Medel Based System Engineering: and Introduction to the
302 mathematical Theory of Discreet system and to the TricotyledonTheory of system
303 Design. CEC Press Boca Raton. Pp. 212-227.
- 304 10. Obalum, S. E., Opong, E., Igwe, C. A., Watanabell, Y. and Obi, M.E. (2013).Special
305 \variability of uncultivated soils in derived savanna.*Int. Agrophys* 27

- 306 16. Achi, C. (2003). Hydrocarbon exploitation, environmental degradation and poverty: the
307 Niger Delta experience. Diffuse Pollution Conference, Dublin 2003 2B Policy / Socio-
308 Economics.
- 309 17. Adenipekun, C.O. (2008). Bioremediation of engine oil polluted soil by pleurotus tuber-
310 regium Singer, a Nigerian whole rot fungus. *African Journal of Biotechnology*. 7(1): 055-
311 058.
- 312 18. Agboola, O.M., Nwulu, N.I., Egelioglu, F., Agboola, O.P. (2011) Gas Flaring in Nigeria:
313 Opportunity for Household Cooking Utilization. *International Journal of Thermal and*
314 *Environmental Engineering*, 2 (2), 69-74.
- 315 19. Ayoola, T. J., (2011) "Gas flaring and its implication for environmental accounting in
316 Nigeria," *Journal of Sustainable Development*. 4(5).244-250.
- 317 20. Arowolo, A. A. and Adaja, I. J. (2011). "Trends of natural gas exploitation in Nigeria and
318 the implications on the socio-economic stability and governance," in 35th Nigerian
319 Statistical Association annual conference..
- 320 21. Anoliefo, G. O. & Vwioko, D.E (2001). Tolerance of *Chromolaena odorata* L. (K. and R.)
321 grown in soil contaminated with spent lubricating oil. *Journal of Tropical Bioscience* 1:
322 20-24.
- 323 22. Abu, G. O & Dike O. P (2008). A study of natural attenuation processes involved in a
324 microcosm model of crude oil-impacted wetland sediment in Niger Delta. *Bioresource*
325 *Technology*. 99 :4761-4767
- 326 23. Akhionbare, S.M.O(2009) Environment Concept, issues and control of Pollution, M.C
327 Computer press Nnewi Anambra State: pp 82-92, 183-227.
- 328 24. Abdulkareem, A. S. (2005). "Evaluation of Ground Level Concentration of Pollutant Due
329 to Gas Flaring by Computer Simulation: A Case Study of Niger-Delta Area of Nigeria,"
330 *Leonardo Electronic Journal of Practices and Technologies*, Vol. 6, No. 1, pp. 29-42.
- 331 25. AGOC (2003). Gas flaring, oil companies and politics in Nigeria. Alexander's Gas and Oil
332 Connections (AGOC) Publications, 6 (20), 15-16.

- 333 26. Abdulkareem, A. S., Odigure J. O. and Abenege, S. (2009). Predictive model for pollutant
334 dispersion from gas flaring: A case study of oil producing area of Nigeria. *Energy Sources*
335 (Part A), 31(12), 1004-1015.
- 336 27. Akachidike, K. (2008) Remote Stranded Gas-Challenges and Opportunities for
337 Development: Proceedings of the 38th Annual Conference of NSChE, Effurun, Delta
338 State, Nigeria.
- 339 28. Aghalino, S. O. (2009). Gas flaring, environmental pollution and abatement measures in
340 Nigeria, 1969 - 2001. *Journal of sustainable development in Africa*, 11(4), 219-238.
- 341 29. Amadi, A., Abbey, S.D. & Nma, A. (2000) Chronic effects of oil spill on soil properties and
342 microflora of a rain forest ecosystem in Nigeria. *Water, Air and Soil Pollution*, 86(1-4)1-
343 11.
- 344 30. Ajugwo, A. O. (2013). Negative Effects of Gas Flaring: The Nigerian Experience. *Journal of*
345 *Environment Pollution and Human Health*, 1 (1): 6-8.
- 346 31. Anyadiegwu, C.I.C, Muonagor, C.M, Nnakaihe, S.E (2014) Comparative Economic
347 Analysis of Using Natural Gas For Liquefied Natural Gas Production and Converting
348 Natural Gas To Diesel Through Gas-To-Liquid Technology In Nigeria. *International*
349 *Journal of Research in Engineering and Science*, 2(9), 9-26.
- 350 32. Blume, H.-P., Horn, R., Alaily, F., Jayakody, A.N. & Meshref, H. 1980. SAND CAMBISOL
351 FUNCTIONING AS A FILTER THROUGH LONG-TERM IRRIGATION WITH WASTEWATER. *Soil*
352 *Science*, 130(4): 186-192.
- 353 33. Blum, W.E.H. 2005. Functions of Soil for Society and the Environment. *Reviews in*
354 *Environmental Science and Bio/Technology*, 4(3): 75-79.
355 <https://doi.org/10.1007/s11157-005-2236-x>
- 356 34. Blum, W.E.H. (2005). Functions of Soil for Society and the Environment. *Rev Environ*
357 *SciBiotechnol*4, 75-79.
- 358 35. Chaineau, C., Yepremian, C., Vidalie, J., Ducreux, J. & Ballerini, D. (2003) Bioremediation
359 of a crude oil-polluted soil: biodegradation, leaching and toxicity assessments. *Water,*
360 *Air and Soil Pollution*, 144(1)419-440.
- 361