Case study

1 2

3

Percutaneous balloon mitral valvotomy and PCI of Left anterior descending artery done in same sitting

5

4

Abstract: In countries like India where rheumatic heart disease is still a common problem, 6 7 with improvements in diagnosis and treatment, the lifespan of these patients is increased. With increase in the lifespan, these patients may develop coronary artery disease and present 8 9 as acute coronary syndrome. In some cases especially RHD with severe MS, thrombus that 10 develop in left atrium may embolize in one of the coronary arteries, leading to acute coronary syndrome. We report a case of 51 year old female who was a known case of rheumatic heart 11 disease and now presented with acute coronary syndrome. Patient was hemodynamically 12 unstable and underwent Percutaneous Balloon Mitral Valvotomy and Percutaneous coronary 13 14 intervention in the same sitting. This is first of such intervention at our institute and also there are very few such records available online. Patient tolerated the procedure well with 15 significant improvement. 16

Keywords: Percutaneous Balloon Mitral Valvotomy, Percutaneous Coronary Intervention,
 Severe Mitral Stenosis, Left Anterior Descending Artery Stenosis, Rheumatic Heart Disease.

Abbreviations: RHD: Rheumatic heart Disease; MS: Mitral stenosis; PCI; percutaneous
 coronary intervention; BMV: Balloon Mitral Valvotomy; ACS: Acute coronary Syndrome

21 Introduction: Rheumatic Heart disease remains one of the leading cardiac diseases in tropical developing countries like India. Around 25%–30% of all cardiac visits to hospitals 22 are related to RHD¹. Patient with RHD can present with left ventricular (LV) dysfunction due 23 to multiple causes including primary rheumatic myocarditis, secondary to LV remodelling 24 due to altered hemodynamics in valvular pathology, cardio-embolic phenomenon involving 25 coronaries, or co-existing coronary artery disease itself. Mitral Stenosis (MS) is most 26 common valvular pathology in RHD. In countries like India where rheumatic heart disease is 27 28 still a common problem, with improvements in diagnosis and treatment, the lifespan of these patients is increased. With increase in the lifespan, these patients may develop coronary 29 30 artery disease and present as acute coronary syndrome. In some cases especially RHD with severe MS, thrombus that develop in left atrium may embolize in one of the coronary arteries, 31 leading to acute coronary syndrome. Management of RHD patient with significant valve 32 dysfunction and significant coronary artery disease, is primarily revascularisation of affected 33 vessels by Coronary Artery Bypass Graft (CABG) Surgery and Valve replacement. In 34 35 selected patients with pliable mitral valve and single vessel disease, percutaneous 36 interventions are possible, like BMV and PCI. However, there are limited guidelines about 37 management of overtly symptomatic patients with ACS in cases of RHD who may not 38 tolerate major surgical intervention.

40 Case Report:

51 year old female, known case of RHD with MS, admitted in peripheral hospital with 41 multiple episodes of Paroxysmal Nocturnal Dyspnea (PND) in last 1 month. She developed 42 43 acute onset chest pain and NYHA class IV breathlessness on the same day. Patient was transferred to our hospital in the night on oxygen support and ionotropic support. On 44 45 admission- her pulse rate was 110/min, Blood Pressure was 100/80 mmHg on ionotropes and bilateral crepts were present on auscultation. Routine blood investigations were normal, 46 Arterial Blood Gas (ABG) showed hypoxia. Her Electrocardiogram (ECG) showed sinus 47 rhythm with poor r wave progression with ST segment downsloping depression in I avL V4 48 49 V5 V6 and T inversion in I avL (Fig 1). Old ECG obtained from the patient's previous 50 records showed normal progression of r wave with no significant ST T changes (Fig 2). 2D Echocardiography showed Severe Mitral Stenosis (Mitral Valve Area (MVA) by Planimetry -51 0.67 cm², Wilkins score 7/16, MV gradient- 15/10 mmHg) (Fig 7) with Left Ventricular 52 53 Ejection Fraction (LVEF) 35%- basal, mid, distal, anteroseptal, anterior and anterolateral 54 segments hypokinetic (Fig 8). Troponin T was significantly raised (50ng/ml). Patient didn't 55 respond well to the medical management, so she was taken in cath lab for percutaneous 56 balloon mitral valvotomy (PBMV) and urgent coronary angiography (CAG). Right femoral 57 venous and arterial access obtained. Pulmonary artery pressure was 54/20 mmHg, aortic pressure was 138/80 mmHg, PCWP was 23 mmHg (mean) and LV-edp was 12 mmHg (Fig 58 59 6). Gradient across mitral valve was 11 (Fig 6). CAG showed LAD (Left Anterior Descending) artery mid segment thrombotic 90% stenosis (Fig 3). In view of the general 60 condition of the patient, BMV was planned first, because PCI requires injectable heparin as 61 62 anticoagulation and transeptal puncture is usually done without giving any anticoagulation. ACCURA balloon No.26 was used and inflation of 26 mm was given for 1 sec (Fig 5). Post 63 balloon dilatation gradient reduced to 2 mmHg and the MVA improved to 1.54cm² (Fig 9). 64 Patient was taken up for PCI to LAD. Then LAD was stented with Drug Eluting Stent (DES) 65 66 2.75 x 24 mm after predilatation. Stent was post dilated with NC balloon 2.75 x 13 mm. Post stenting check showed TIMI III flow with no residual lesion (Fig 4). Patient tolerated 67 68 procedure well. Patient was shifted to post operative monitoring and care. Post procedure 69 patient improved clinically. Over next few days she was off oxygen support and ionotropes 70 were tapered. Gradually she became ambulatory without symptoms during routine activities. Patient was discharged after 5 days of procedure. Patient is asymptomatic on subsequent 71 follow ups. 72

73 **Discussion:**

In countries like India, patients with RHD are commonly found to have CAD, more often, if 74 presentation is in late middle age or in elderly age group. There is limited data of incidence of 75 primary coronary artery disease in patients of RHD. In study by Jose et al², the overall 76 prevalence of CAD in RHD patients undergoing valve surgery was 12.2%. In another study, 77 done by Dinesh et al³ showed that 9.05% of RHD patients (above 40 years of age) have 78 significant CAD. The Left Anterior Descending (LAD) artery is the most common vessel 79 involved. Acute coronary syndrome in RHD patients can be secondary to cardioembolic 80 phenomenon involving coronaries or due to atherosclerotic disease involving native vessels. 81

82 Acute coronary syndrome secondary to thrombo-embolic phenomenon was reported in studies by Radhakrishnan et al^4 , Niniek Purwaningtyas et al^5 , Cardoz J et al^6 . Ideally 83 transesophagial echocardiography should be done to rule out thrombus in left atrial 84 85 appendage even if the patient is in sinus rhythm. Differentiating these two etiologies, denovo thrombosis or thromboembolic phenomenon, needs intravascular imaging with IVUS 86 87 (Intravascular Ultrasound) or OCT (Optical Coherence Tomography), either of which was not 88 done in our patient due to hemodynamic instability and financial constraints. There are limited available records of patients undergoing PCI and BMV in the same setting. Patients 89 undergoing PCI needs to be heparinised and loaded with dual antiplatelets and in setting of 90 BMV where septal puncture carries inherent risk of pericardial effusion and cardiac 91 92 tamponade this surgery carries high risk to the patient. Most of the patients with coexisting CAD and RHD are referred for valve replacement with CABG. However, sometimes patients 93 may not be stable enough or willing for major operative intervention. In our patients we first 94 started with BMV as mitral valve was pliable and there was no significant Mitral 95 regurgitation. Patient was heparinised after septal puncture and after echo confirmation of no 96 pericardial effusion. Patient was loaded on table with Clopidogrel chewed and kept 97 sublingually, although she was on dual antiplatelets for her ischemic event before procedure 98 as well. The overall procedure was done with due care under fluoroscopic guidance and 99 patient stood procedure well. Similar report of BMV and PCI done in same sitting done by 100 Paul G J et al⁷ was also safe. Paul G J et al⁷ also initiated with BMV and later PCI was done. 101 With our experience and limited available online records it is understood that patients with 102 103 ACS and RHD can undergo both transluminal valvular and coronary intervention in the same 104 sitting with acceptable safety under adequate caution and care.

Figure 1: ECG at the time of presentation. Showing T inversion in I avL, ST sagging in 105

106 lateral leads, Poor R wave progression.



- 108
- 109

110

111 Figure 2: ECG from old records of patient. No significant ST-T changes and Sinus Rhythm.



113 Figure 3: CAG showing LAD mid thrombotic lesion causing significant stenosis.



119 Figure 4: CAG post PCI with Drug eluting stent in Mid AD. Good Result. TIMI III flow



120 121

- 122 Figure 5: Balloon inflation with ACCURA Balloon No. 26 placed across mitral valve and its
- 123 inflation





- 126 Figure 6: Pressure tracing. Blue represent LV pressures and Red represents PCWP (LA
- 127 pressure).





130 Figure 7: ECHO image showing pre BMV MVA of 0.67 cm^2





138 Figure 8: M mode Echo Image showing Anterior wall hypokinesia



139

140 Figure 9: ECHO image showing post BMV MVA of 1.54 cm^2



141

142 Conclusion:

In selected patients with pliable mitral valve and single vessel disease presenting with ACS
who are hemodynamically unstable, percutaneous intervention with BMV and PCI may be

- done to stabilize the patient hemodynamically and symptomatically.
- 146

147 **References:**

148	1.	Padmavati S. Rheumatic fever and rheumatic heart disease in India at the turn of the
149		century. Indian Heart J 2001;53:35-7.
150	2.	Jose VJ, Gupta SN, Joseph G, Chandy ST, George OK, Pati PK, et al. Prevalence of
151		coronary artery disease in patients with rheumatic heart disease in the current era.
152		Indian Heart J 2004;56:129-31

153	3.	Choudhary D. et al (2016) "Prevalence of coronary artery disease in rheumatic heart
154		disease and comparison of demographic and coronary artery disease profile with
155		atherosclerotic coronary artery disease", Advances in human biology 2016; 6:76-83.
156	4.	Radhakrishnan S, Alagesan M, Kaliappan T, Gopalan R. Therapeutic dilemma –
157		Acute coronary syndrome in the presence of severe mitral stenosis. JICC. 2014; 4(2):
158		128-131
159	5.	Niniek Purwaningtyas. Acute Myocardial Infarction in Patient with Mitral Stenosis: A Rare
160		Case J Cardiovasc Dis Diagn 2018, 6:5
161	<mark>6.</mark>	Cardoz J, Jayaprakash K, George R .Mitral stenosis and acute ST elevation myocardial
162		infarction. Proc (Bayl Univ Med Cent).2015;28(2):207–209
163	7.	Paul G. J. et al (2018) "Percutaneous transvenous mitral commissurotomy and
164		coronary intervention in kyphoscoliosis", Indian Heart Journal 2018; 1:151-154.
165		