

## **Original Research Article**

### **COMPARATIVE ANALYSIS OF THE TOP SOIL PROPERTIES UNDER FORESTED AND DEFORESTED ZONES: IMPLICATIONS FOR THE ENVIRONMENT**

#### **Abstract**

An investigation was carried out to examine the properties of top soils between 0 and 30cm under both deforested and forested zones in Bowen University, Iwo, Nigeria. Top soil samples in the deforested zone was taken from the Main Gate area of the institution while that of the forested zone was taken from the forested area opposite staff quarters of the University. The soil samples were subjected to standard laboratory tests in the University central laboratory. The results showed that deforested soil has sandy, clay and silt contents of 72.4%, 9.2% and 18.4% respectively while forested soil has 65.2%, 10.8% and 24% in the same order. Also it was discovered that deforested soil has organic carbon, organic matter, pH, field capacity, moisture and electrical conductivity of 0.32%, 0.55%, 6.8, 0.72g, 126.9g and 230 $\mu$ f/cm respectively while forested soil has 0.45%, 0.77%, 7.1, 0.90g, 0.72g, 129.2g and 275 $\mu$ f/cm in the same order. The implications of this results is that removal of vegetation contributes to the release of carbon into the atmosphere which increases atmospheric heat, alkalinity of soil, loss of soil nutrients and also could pose limits to the survival of plants growth and also susceptibility of soil to surface wash. Thus, it is recommended that effort should be made to checkmate the removal of vegetation and if unavoidable, relevant policies should be put in place for edge development and its maintenance and also, shades growth as remedies to ensure sustainable environment.

**Key words:** Top soil, forested soil, deforested soil, community development, soil properties,

#### **Introduction**

Despite the fact that deforestation has been seen as an act that should be avoided because of its negative impact on human environment, the process is still naturally desirable in certain situations and circumstances both in space and time. For instance, Umana (2018) considered that deforestation is unavoidable in view of the increase in population and also urbanization which also implies demand for forest resources including trees and other biodiversities. Furthermore, Sambe *et al* (2018), also added that in view of the need to embark on both industrial and urban development, deforestation can hardly be avoided. Deforestation is the permanent destruction of forests in order to make the land available for other uses (Inyang and Esohe, 2014). An estimated 18 million acres (7.3 million hectares) of forest, which is roughly the size of the country of Panama, are lost each year, according to the United Nations' Food and Agriculture Organization (FAO,World Wildlife Fund (WWF)). Also, in 2016, global tree cover loss reached a record of 73.4 million acres (29.7 million hectares), according to the University of Maryland, deforestation

40 occurs around the world, though tropical rainforests are particularly targeted. If current  
41 deforestation levels proceed, the world's rainforests may completely vanish in as little as 100  
42 years, according to National Geographic. Deforestation also has impacts on social aspects of the  
43 country, specifically regarding economic issues, agriculture, conflict and most  
44 importantly, quality of life. According to data taken over 2000 to 2005 Nigeria, located in the  
45 western region of Africa, has the largest deforestation rates in the world, having lost 55.7% of  
46 their primary forests (Forest in Nigeria, 2018). Mongabay defines primary forests as forests with  
47 no visible signs of past or present human activities.

48 The annual rate of deforestation in Nigeria is 3.5%, approximately 350,000-400,000  
49 hectares per year (FAO, 2018). The Food and Agriculture Organization of the United Nations  
50 lists the requirements of sustainable forest management as: extent of forest resources, biological  
51 diversity, forest health and vitality, productive functions of forest resources, protective functions  
52 of forest resources, socio-economic functions and a legal, policy and institutional framework  
53 (FAO, 2018). Many aspects of the outline are currently not being met and will continue to have  
54 detrimental effects if not quickly addressed. A lot of damage has been done to Nigeria's land  
55 through the processes of deforestation, notably contributing to the overwhelming trend of  
56 desertification. Desertification is the encroachment of the desert on land what was once fertile  
57 (Ojudgo, 2010). A study conducted from 1901 to 2005 gathered that there was a temperature  
58 increase in Nigeria of 1.1 °C, while the global mean temperature increase was only 0.74 °C. The  
59 same study also found in the same period of time that the amount of rainfall in the country  
60 decreased by 81mm. It was noticed that both of these trends simultaneously had sharp changes in  
61 the 1970s (Omofonmwan and Osa-Edoh, 2008).

62 From 1990 to 2010 Nigeria nearly halved their amount of forest cover, reduced from  
63 17,234 to 9041 hectares. The combination of extremely high deforestation rates, increased  
64 temperatures and decreasing rainfall are all contributing to the desertification of the country.  
65 The carbon emissions from deforestation is also said to account for 87% of the total carbon  
66 emissions of the country (Akinbamiji, 2003).

67 The process of deforestation poses a lot of implications for various natural resources and  
68 most importantly, soil resources (Anyanwu, 2015). Deforestation exposes soils to direct surface  
69 runoff. Tree roots anchor the soil. Without trees, the soil is free to wash or blow away, which can

70 lead to vegetation growth problems. The WWF states that scientists estimate that a third of the  
71 world's arable land has been lost to deforestation since 1960. After a clear cutting, cash crops  
72 like coffee, soy and palm oil are planted. Secondly, the initial increase of soil erosion is almost  
73 certainly due to the removal of the canopy and surface litter that protects the soil surface from  
74 the energy of raindrop impact and surface detachment. Surface cover is probably the greatest  
75 single management effect on soil erosion (Nearing et al., 1994). Soil structure is destroyed by the  
76 plow and the stabilizing effects of root fibers become insignificant as the roots are shredded by  
77 the tillage microbially decomposed following deforestation. As pore space increased due to the  
78 mechanical cultivation, the air exchange increased the available oxygen for microbial decay of  
79 organic matter, particularly the particulate organic matter (POM) that is highly effective at  
80 binding soil particles (Wang, 2002). This factor, coupled with the accelerated erosion rapidly  
81 depleted the SOM in the plow layer and weakened the soil WAS (Zhang and Horn, 2001). Other  
82 effects of deforestation are loss of species and disturbed water cycle. This work aimed to  
83 compare the top soil characteristics under both deforested and forested regions in the  
84 southwestern part of Nigeria. Specific objectives are to: examine the characteristics of top soil  
85 (0-30cm) in the study area and to evaluate the implication of the result on environmental  
86 sustainability.

## 87 **Methodology**

### 88 **Study Area**

89 This study site for this research is Bowen University (7.61464°N, 4.1372°E), Iwo, Osun State,  
90 Nigeria. Bowen University, Iwo, owned by the Nigerian Baptist Convention, became operational  
91 in 2002 at the site of the old Baptist College (a proscribed Teachers' Training College). Since its  
92 inception, the University has embarked on series of projects to enhance its status in the world of  
93 other global institutions. Such projects include, building projects, such as Library, Faculty  
94 buildings like Social and Management, Agriculture, Sciences, College of Health Sciences,  
95 Hostels, landscaping among others. The implications of all these developmental projects are the  
96 clearing of the forest to pave way for the erection of these buildings. Currently, forest trees are  
97 being cleared firstly, to give room for the beautification of the institution with ornamental and  
98 economic trees, and other horticultural plants from the main Gate of the institution to the Chapel  
99 junction and secondly to expand the University Commercial farm of the Faculty of Agriculture.  
100 All these projects have led to the loss of various environmental resources including vegetation,

101 soil resources and exposure of watershed. It is on this premise that, even though some of the  
102 cleared portions are being regrown, the cleared forest could have had effects on soil conditions  
103 including its physico-chemical constituents and on the environment generally. Thus this study  
104 aims at assessing the impacts of deforestation on soil conditions. Specific objectives are to  
105 determine the effects of deforestation on soil moisture and other physical properties and also to  
106 assess the implication of deforestation on the properties of soils in the study area.

#### 107 **Data sources**

108 The data required for this research were soil samples. Soil samples were taken from both  
109 deforested areas and forested areas within the University in April, 2019, after the incidence of  
110 rainfall events. The samples from deforested zone were taken close to the Main University Gate.  
111 This area formerly under forest cover was cleared during the dry season while soil samples under  
112 forest cover were taken under the forest adjacent to the cleared zone. The samples were taken  
113 simultaneously during the daytime with the use of soil auger in triplicate between 0-30cm and  
114 kept in polythene bags. The samples were immediately taken to the laboratory for analysis. Soil  
115 parameters analyzed are soil moisture, bulk density, particle size distribution, organic carbon,  
116 electrical conductivity and soil microorganisms.

#### 117 **Method of analysis:**

118 Six soil parameters analyzed include soil pH, particle size distribution, organic carbon, organic  
119 matter, electrical conductivity, soil moisture contents and the soil bulk density. The analyses of  
120 the parameters were carried out following standard laboratory techniques. The selected  
121 parameters were based on the analysis facilities available and the time frame. The soil pH of the  
122 samples were measured using Testr 2 waterproof digital pH meter with soil as described by  
123 Hendershot *et al.*, (1993) while the soil moisture contents of the two samples were measured  
124 using gravimetric method. Particle size distributions and field capacity were determined with use  
125 of hydrometer method (Adepetu *et al.*, 1984). Organic Carbon was measured using the procedure  
126 according to Golterman *et al.*, (1978); Electrical Conductivity was determined following  
127 Ademoroti (1996) and lastly Bulk Density (BD) was determined using the procedure according  
128 to King (1988).

129

## 130 **Results and Discussion**

### 131 **Bulk Density**

132 Bulk density reflects the soils ability to function for structural support, water and solute  
 133 movement and soil aeration. It is also defined as weight of fiber per unit volume, often expressed  
 134 as  $\text{g/m}^{-1}$ , and is a good index of structural changes (Sreerama *et al.*, 2009). Bulk density increases  
 135 with compaction and tends to increase with depth. From Table 1, the result shows that forested  
 136 soil has the higher bulk density of  $0.9345 \text{ g/m}^{-1}$  compared to  $0.9210 \text{ g/m}^{-1}$  of the deforested soil.  
 137 In other words forested soil is more compact than deforested soil, which is due to vegetation  
 138 found on its surface, because the roots of plants tend to hold soil where it absorbs nutrients from.  
 139 This finding buttresses the observation made by Tefera *et al.* (2007) that soils with higher bulk  
 140 density are more compact.

141 **Table1: Summary of the mean soil physical and organic parameters analysis**

Sample Name	Bulk Density	Soil pH	Organic carbon (%)	Organic Matter (%)	Soil Moisture content ( $\text{g/cm}^3$ )	Field Capacity (g)	Soil Electrical conductivity ( $\mu\text{f/cm}$ )
Deforested	0.9345	6.8	0.32	0.55	126.9	0.72	230
Forested	0.9210	7.1	0.45	0.77	129.2	0.90	275

#### 143 Soil pH

144 Soil pH is the degree of acidity or alkalinity of the soil. Soil pH is very important because it  
 145 directly affects soil nutrients availability. Plant roots can only absorb nutrients after they have  
 146 been transformed into certain ionic form. Soil pH affects the availability of nutrients and how the  
 147 nutrients react with each other. At a low pH, beneficial elements such as molybdenum (Mo),  
 148 phosphorus (P), magnesium (Mg) and calcium (Ca) become less available to plants (Landis *et al*  
 149 2003). From Table 1, the result shows that forested soil has 7.1 and deforested soil has 6.8. The  
 150 result also indicates that deforested soil is slightly acidic and forested soil is slightly alkaline  
 151 which also implies that forested soil have higher nutrient availability compare to deforested soil  
 152 because they tend to transform nutrients into ionic form due to its slight alkalinity. Deforested

153 soil tends to have low population of micro-organism because of its slightly acidic nature as some  
154 **micro-organism** cannot survive acidic soil compared to the forested soil which corroborate the  
155 findings of Landis *et al.* (2003) an acidic soil is dangerous to human and the ecosystem. The  
156 result here indicated that efforts should be made to avoid incessant removal of forest in our  
157 **ecosystem**

### 158 **Organic Carbon**

159 Organic carbon is part of the natural carbon cycle. World's soil holds around twice the amount of  
160 carbon that is found in the atmosphere and in vegetation (Hoyle, 2013). Soil organic carbon is  
161 important for all three aspects of soil fertility, namely chemical, physical and biological fertility  
162 (Viscarra Rossel *et al.*, 2014). From Table 1, Forested soil has 0.45% and Deforested soil has  
163 0.32%, which means that forested soil has good structure, better biological and physical health of  
164 soil, and best buffer against harmful substances compared to deforested soil. This result,  
165 however, further ascertain the findings made by Ingram *et al.* (2001) that soil organic carbon  
166 promotes soil structure by holding the soil particles together as stable aggregates improves soil  
167 and physical properties such as water holding capacity, water infiltration, gaseous exchange, root  
168 growth and ease of cultivation. This findings explains the reason behind the depletion of ozone  
169 layers in the atmosphere. The situation that has generated the current global warming worldwide.

### 170 **Organic Matter**

171 Organic matter binds soil particle into aggregates. From Table 1, forested soil has the highest  
172 organic matter of 0.77% and deforested soil of 0.55%. Forested soil has better supply of nutrient,  
173 better habitat and higher water holding capacity as compared to deforested soil. Deforested is  
174 prone to soil erosion compared to forested soil because the higher the organic matter higher the  
175 soil particles are binded into aggregates thereby buttressing the findings made by FAO and ITPS,

176 (2015) that Organic matter improves soil aggregate and structural stability which, together with  
177 porosity, are important for soil aeration and the infiltration of water into soil. While plant growth  
178 and surface mulches can help protect the soil surface, a stable, well-aggregated soil structure that  
179 resists surface sealing and continues to infiltrate water during intense rainfall events will  
180 decrease the potential for downstream flooding.

### 181 **Soil Moisture Content**

182 Soil moisture content refers to the quantity of water contained in the soil which plays great role  
183 in soil and plant growth relationship. The result from Table 1, shows that forested soil has the  
184 higher soil moisture content of 129.2g compared to 126.9g for deforested soil which may be due  
185 to vegetation cover. It also implies that forested soil has higher regulatory tendency of physical,  
186 chemical and biological activities in the soil as compared with the deforested. Deforested soil is  
187 exposed to direct sunlight thereby losing much of its moisture content. Which confirms Hillel,  
188 (1982) findings that soil moisture content contributes o deeper plant root growth, reduced soil  
189 run-off/leaching and less favourable conditions for insect and fungal diseases.

### 190 **Field Capacity**

191 Field capacity as to do with the amount of soil moisture or water content held in the soil after  
192 excess water has drained away and the rate of downward movement has decreased. From Table  
193 1, forested soil has the highest value for field capacity 0.90g compared to deforested soil of  
194 0.72g which means forested soil is in good condition due to vegetation cover. This implies  
195 reduction in the rate of evaporation and transpiration when compared with deforested soil which  
196 is exposed to direct sunlight with no vegetation cover. This supported Kramer's, (1983, p.71)  
197 observation that the amount of water retained at field capacity decreases as the soil temperature



198 increases. It also means that forested soil has higher soil organic matter content compared to  
199 forested soil because it facilitates soil water holding capacity (Hillel, 1971, p. 165).

200

### 201 **Electrical Conductivity**

202 Soil electrical conductivity is a measure of the amount of salts in the soil. It also means the  
203 salinity of soils. It is an economically friendly method of calculating available nitrogen for plant  
204 growth. Table 1, shows that forested soil has the highest conductivity with value of  $275\mu\text{f}/\text{cm}$   
205 compared to  $230\mu\text{f}/\text{cm}$  for deforested soil. This implies that forested soil has higher nutrient  
206 composition than deforested soil because the lower the value of conductivity the higher the  
207 nutrients availability in the soil. It also implies that forested soil has higher percentage of clay  
208 content and higher Cation Exchange Capacity because the higher the Cation Exchange Capacity  
209 the higher the soil electrical conductivity (Wiatrak et al., 2009). This study reveals that soil with  
210 lower value of sand, higher value of clay and silt have higher electrical conductivity. The farther  
211 the soil pH move away from the neutral point the more electrical conductive they become. The  
212 amount of nutrient in the soil tells the soil electrical conductivity. The higher the soil nutrient the  
213 higher the electrical conductivity. The higher the soil moisture content the higher the soil  
214 electrical conductivity.

### 215 **Particle Size Distribution**

216 Particle size distribution is the proportions by dry mass of a soil distributed over specified  
217 particle-size ranges. The physical and chemical behaviors of soils are significantly influenced by  
218 particle-size distribution which is important for soil interpretations, determination of soil  
219 hydrologic qualities, plant nutrient requirements and classifications (Eshel et al., 2004). From the  
220 result showed shown in Table 2, forested soil has 65.2% of sand, 10.8% of clay, and 24% of silt,  
221 while deforested soil has 72.4% sand, 9.2% clay and 18.4% silt which implies that deforested



222 soils has a better particle size distribution in term of plant growth advantage as compared to  
 223 forested soil which also means that deforested soil has faster water and nutrient movement in the  
 224 soil because particle size. This is used to classify soils for agricultural purposes and also  
 225 influences how fast or slow water or other fluid moves through soil which ascertain the  
 226 observation made by Sandhage-Hofmann *et al.* (2015) that soils with high sand content do not  
 227 get compacted which aid the free flow of water or liquid.

228 **Table 2: Summary of soil sample particle size distribution analysis**

Sample Name	Soil Depth (cm)	Sand (%)	Clay (%)	Silt (%)	Total (%)
Sample A (Under forested zone)	0-30	65.2	10.8	24	100
Sample B (Under deforested zone)	0-30	72.4	9.2	18.4	100

229

### 230 **Conclusion**

231 The properties of top soils under both forested and deforested zones within the Bowen  
 232 University, Iwo, Nigeria were analyzed using standard laboratory methods. The results revealed  
 233 that top soils in the forested area has higher percentage of bulk density, soil moisture, field  
 234 capacity but lower soil pH, soil carbon and electrical conductivity when compared with same  
 235 properties of top soils in the deforested areas. However, the results implied that effort should be  
 236 made to avoid forest removal in order to protect the top soil and also to checkmate the release of  
 237 carbon into the atmosphere which has consequential effect on the depletion of the ozone layer,  
 238 which leads to atmospheric heat. Also there is need to protect the top soil for agricultural  
 239 purposes. Planting of trees as shades and edges should be encouraged in a situation where the  
 240 removal of forest is unavoidable especially as a result of urban expansion, agricultural and  
 241 industrial projects.

242

### 243 **REFERENCES**

- 244 Adepetu, J.A., Obi, A.O. and Amusan, A.A. (1984) *Soil Science laboratory manual*,  
 245 Department of Soil Science, University of Ife, 2<sup>nd</sup> Edition, 33-88pp.  
 246  
 247 Ademoroti, C.M.A. (1996) *Environmental Chemistry and Tixicology*. Foludex Press Ltd.,  
 248 Ibadan, 176-177pp.

- 249 Anyanwu, J. C.1, Egbuche, C. T., Amaku. G. E., Duruora, J.O. and Onwuagba, S.M. (2015) The  
 250 Impact of Deforestation on Soil Conditions in Anambra State of Nigeria. *Agriculture, Fosrestry*  
 251 *and Fisheries*, 4(3-1): 64-69.
- 252 Akinbami, J. (2003) An Integrated Strategy for Sustainable Forest–energy–environment  
 253 Interactions in Nigeria. *Journal of Environmental Management* 69.2: 115-28. Science  
 254 Direct.
- 255 Eshel G, Levy J, Mingelgrin U & Singer M J (2004). Critical evaluation of the use of laser  
 256 diffraction for particle-size distribution analysis. *Soil Science Society of American*  
 257 *Journal* 68:736-743
- 258 FAO and ITPS (2015) Status of the World’s Soil Resources (SWSR)—Main Report. Food and  
 259 Agriculture Organization of the United Nations and Intergovernmental Technical Panel  
 260 on Soils, Rome, Italy. <http://www.fao.org/3/a-i5199e.pdf>.
- 261 Golterman, R.T., Clymo, R.S. and Ohnstad, M.A.M. (1978) *Methods of physical and chemical*  
 262 *analyses of soil and freshwater. IBP Handbook, No (8)*. Blackwell Scientific Publication,  
 263 Oxford Press, 214pp.
- 264 Hendershot, W.H., Lalonde, H. and Duguet, M. (1993) Soil reacytion and exchangeable acidity  
 265 in Carter, M.R. (Ed.) *Soil Sampling and methods of Analysis fo Canadian Society of Soil*  
 266 *Science*, Lewis Publisher, BOCA Raton, Florida, P141-145.
- 267 Hillel D., (1982) Introduction to soil physics. *Applied Ecology and Environmental Sciences*.  
 268 **2014**, Vol. 2 No. 2, 43-47
- 269 Hillel, D. (1971) Soil and Water, Physical Principles and Processes. *Journal of Water Resource*  
 270 *and Protection*, [Vol.6 No.19](#), December 30, 2014
- 271 Hoyle, FC 2013, *Managing soil organic matter: A practical guide*, Grains Research and  
 272 Development Corporation, Kingston, viewed 15 October 2018,  
 273 [https://grdc.com.au/resources-and-publications/all-ublications/publications/2013/07/grdc-](https://grdc.com.au/resources-and-publications/all-ublications/publications/2013/07/grdc-guide-managingsoilorganicmatter)  
 274 [guide-managingsoilorganicmatter](#).
- 275 Ingram, JSI, Fernandes, ECM 2001, 'Managing carbon sequestration in soils: concepts and  
 276 terminology', *Journal of Agriculture, Ecosystems and Environment*, vol. 87, pp. 111–117.
- 277 Inyang, M.P. and Esohe, K.P. (2014) Deforestation, Environmetal Sustainability and Health  
 278 implications in Nigeria- A Review. *Int. J. Sci. Env. Tech.*, 3(2): 502-517.
- 279 King, L.D. (1988) Retention of water by several soils of the southeastern United States. *Journal*  
 280 *of Environmental Quality*, 17(2):246.
- 281 Landis, J.K., Wayne, G. and Ming-Ho Yu. (2003). Introduction to environmental toxicology:  
 282 Impacts of chemicals upon ecological systems. CRC Press, Lewis Publishers, Boca  
 283 Raton, FL.
- 284 Mba, E.H. (2018) Assessment of Environmental impacts of deforestation in Enugu, Nigeria.  
 285 *Resources and Environment*, 8(4): 207-215.
- 286 Nearing, M.A., L.J. Lane, and V.L. Lopes. 1994. Modeling soil erosion. In: R. Lal (Ed.) *Soil*  
 287 *Erosion Research Methods* (2nd Ed.). Soil and Water Conservation Society, Ankeny, IA  
 288 pp. 127-156.
- 289 Odjugo, P. A. (2010) General Overview of Climate Change Impacts in Nigeria. *Journal of*  
 290 *Human Ecology* 29(1): 47-55.
- 291 Omofonmwan, S. I. and Osa-Edoh, G.I. (2008) The Challenges of Environmental Problems in  
 292 Nigeria. *Journal of Human Ecology* 23.1: 53-57.
- 293 Sambe, L.N., Adeofun, C.O. and Dachung, C. (2018) the Economic and Ecological effects of  
 294 deforestation on the Nigerian Environment. *Asian J. Adv. Res. Rep.*, 1(2): 1-25.

- 295 Sandhage-Hofmann, A., Kotze, E., van Delden, L., Dominiak, M., Fouche, H., van der  
296 Westhuizen, H., Oomen, R., du Preez, C., & Amelung, W. (2015) Rangeland  
297 management effects on soil properties in the savanna biome, South Africa: A case study  
298 along grazing gradients in communal and commercial farms. *Journal of Arid*  
299 *Environments*, 120, pp. 14-25.
- 300 Tefera A, Seyoum T, Woldetsadlik (2007). Effect of Disinfection, Packaging, and Storage  
301 Environment on the Shelf Life of Mango. *Biosys. Eng.* 96(2):201–212.
- 302 Umana, K. (2018) Environmental Impacts of Deforestation in Nigeria. *Research cyber, Private*  
303 *Policy*
- 304 United Nations, Food and Agriculture Organization of the United. "[Natural Forest](http://www.fao.org)  
305 [Management](http://www.fao.org)". www.fao.org. Retrieved 15 April 2018.
- 306 Viscarra Rossel, RA, Webster, R, Bui, EN & Baldock, JA 2014, 'Baseline map of organic carbon  
307 in Australian soil to support national carbon accounting and monitoring under climate  
308 change', *Global Change Biology*, vol. 20, pp. 2953–2970, doi:10.1111/gcb.12569.
- 309 Wang, G. H. (2002) Plant traits and soil chemical variables during a secondary succession on the  
310 Loess Plateau. *Acta botanica sinica*. 44(8): 990-998.
- 311 Wiatrak, P., Khalilian, A., Mueller, J., Henderson, W. 2009. Applications of Soil Electrical  
312 Conductivity in Production Agriculture. *Better Crops Magazine*. Vol. 93 (2) 16-17.
- 313 Zhang, B. and R. Horn. (2001) Mechanisms of aggregate stabilization in Ultisols from  
314 subtropical China. *Geoderma*, 99: 123–145.
- 315
- 316
- 317
- 318