

## **Original Research Article**

### **Priming with Methylene Blue Enhances the Antioxidant Properties and Germination Power of Cowpea, Millet and Sorghum Seeds**

Please, authors should check the rules of this journal and adapt to the rules.

#### **Abstract**

The effect of priming cowpea, millet and sorghum seeds with methylene blue (MB) on germination power (GP) was investigated in this work. The seeds were primed by steeping for 6 h in different concentrations of MB solution, germinated over 48 h on wet cotton and the GP determined. Further, soluble protein and total phenolic contents, and the antioxidant properties (scavenging ability for hydrogen peroxide, nitric oxide, hydroxyl radical, and DPPH and iron chelation ability) of the seeds steeped in MB that gave the highest GP were determined. The highest GP was observed for cowpea and sorghum seeds steeped in 80  $\mu$ M MB solution, and 60  $\mu$ M MB solution for millet seeds. At these MB concentrations, a faster rate of germination coupled with an increase in soluble protein and total phenolic content was observed. The increase in germination rate was also characterized by enhanced antioxidant properties of the seeds.

**Comment [User1]:** I suggest inserting a sentence or two on the subject, before describing the objective of the work

**Comment [User2]:** Enter a phrase if search completed

**Keywords:** Antioxidant capacity, Germination power, Methylene blue, Scavenging ability

**Comment [User3]:** Insert the scientific names of the studied species as keywords

## 1. Introduction

Seed priming is the preparation of plants to overcome both biotic and abiotic stress during germination by increasing or boosting the plants defense system. Priming strategies such as hydropriming and osmo-priming have been employed for long periods to improve the germination of seeds (Lutts et al. 2016). In mungbeans (*Vignaradiata*), a faster and more efficient germination was observed upon hydro- and osmo-priming (Umair et al. 2012). Similar increases in germination rates have also been observed in canola (Zheng et al. 1994) and rice (Basra et al. 2005; SukSoon and Jae-Hyeun 2000; Maqsood et al. 2003) seeds upon priming.

In addition to hydro- and osmo-priming, other priming strategies such solid matrix priming, hormoprining and bioprining have also been employed to improve seed germination (Lutts et al. 2016). Chemical priming which involves the treatment of seeds with natural or synthetic compounds is also lately gaining much attention. The use of  $\beta$ -amino butyric acid on *V. radiata* with the aim of increasing tolerance to drought and salt stress has been reported (Jisha and Puthur 2016). Growth regulators and salt have also been used to enhance the germination of wheat seed (Nayyer et al. 1995). Other chemical seed priming agents such as ascorbate to improve the drought and salt resistance of wheat have been reported (Farooq et al. 2013; Fercha et al. 2014). Chemicals such as methylene blue, owing to the presence of the free radical scavenging serial conjugate bonds in its structure, may have the ability to act as a favourable seed priming agent.

Methylene blue (MB), a dye which dissolves in water to produce a deep blue colour, has been shown to have antioxidant properties. In mammalian cells, MB was shown to delay senescence by increasing mitochondrial respiration (Atamna et al. 2008). This effect was attributed to the ability of MB to scavenge for reactive oxygen species (ROS) (Aloni et al. 2010), thereby

reducing oxidative damage to cells (Atamna et al. 2008). The ability of MB to scavenge for ROS might have a beneficial effect on plants.

Seed germination is accompanied by increased production of ROS as a result of increased respiration during germination (Aloni et al. 2010). Thus, in this work, it was postulated that through its role in reducing oxidative damage, MB might have a beneficial effect on seed germination. Accordingly, the efficacy of MB in promoting seed germination was investigated by steeping cowpea, millet and sorghum seeds in MB solution prior to germination. The germination power of the seeds was determined and the effect of antioxidant attributes of MB investigated.

**Comment [User4]:** It is necessary to insert the objective of the study, in a clear and direct way, as the last sentence of the introduction

## **2. Materials and method**

### **2.1 Sample collection and preparation**

Certified cowpea (*Vigna unguiculata*), millet (*Eleusine coracana*, Finger millet) and sorghum (*Sorghum bicolor*) seeds were obtained from a registered Agro-Chemical shop in Cape Coast, Ghana. The seeds were briefly disinfected in 1 % hypochlorite solution and washed with distilled water.

Priming was carried out by steeping 500 seeds in different concentrations of methylene blue solution (0, 10, 20, 40, 60, 80, 100 and 150  $\mu$ M) for 6 h. The methylene blue solution was then discarded, the seeds germinated on wet cotton and the germination power of the seeds determined after 48 h. Three replicate experiments per condition were carried out and the seeds which were not steeped in MB (steeped only in water) used as the control.

In another experiment, seeds were either steeped in MB solution or water (control) for 6 h and the

GP determined every 12 h for a duration of 48 h. This was only carried out for the MB solution which concentration gave the highest GP for the different seeds. After 48 h, the germinated seeds

were homogenized in phosphate buffer, pH 7.0 (1:2,  $\text{g mL}^{-1}$ ). The homogenized samples were centrifuged at 10,000  $\times g$  for 20 min and the supernatant recovered was aliquoted and stored at 20°C for the determination of soluble protein content, total phenolic content, scavenging ability for hydrogen peroxide, nitric oxide, hydroxyl radical, and 2,2-diphenyl-1-picrylhydrazyl (DPPH) as well as iron chelating ability.

Comment [User5]: ??

Formatted: Superscript

Comment [User6]: ??

## 2.2 Determination of soluble protein and total phenolic content

Protein content of the phosphate extract of the seeds was determined based on the Biuret assay using bovine serum albumin as a standard. Biuret reagent was added to the seed extract in a 3:2 ratio, incubated at 37°C for 10 min and the absorbance measured at 562 nm.

The total phenolic content of the phosphate extract of the seeds was determined based on the Folin-Ciocalteu assay using gallic acid as a standard (Ampofo-Asiama and Quaye 2018). Briefly, 0.5 mL of the seed extract was mixed with 2.5 mL of 10% Folin-Ciocalteu's reagent and 2.5 mL of 7.5%  $\text{NaHCO}_3$  solution. The mixture was vortexed and incubated at 45°C for 45 min and the absorbance measured at 765 nm.

## 2.3 Determination of hydrogen peroxide, nitric oxide and hydroxyl radical scavenging ability

Phosphate buffered hydrogen peroxide solution (40 mM) was prepared by diluting 30% aqueous hydrogen peroxide ( $\text{w/w}^{-1}$ ) in 100 mM phosphate buffer, pH 7.4. The hydrogen peroxide solution

Formatted: Superscript

(0.6 mL) was added to the seed extract to a final volume of 3.0 mL. The mixture was incubated in the dark for 10 min and the absorbance measured at 230 nm. The absorbance of the buffered hydrogen peroxide solution was also measured and the percent hydrogen peroxide scavenging ability estimated (El and Karakaya 2004).

To determine the nitric oxide radical scavenging ability, 2 mL of 10 mM freshly prepared sodium nitroprusside in a phosphate buffer was mixed with 0.5 mL of the seed extract. The mixture was incubated for 150 min at 25°C and 0.5 mL aliquot added to 0.5 mL of Griess reagent. This mixture was mixed and incubated at room temperature for 30 min after which the absorbance was measured at 540 nm. Gallic acid was used as the positive control (Boora et al. 2014).

Hydroxyl radical scavenging ability was determined based on the assay of Pavithra and Vadivukkarasi (2015). To varying concentrations of the seed extract, 1.0 mL of iron-EDTA solution was added, followed by 1.0 mL of DMSO (in phosphate buffer) solution and 0.5 mL of ascorbic acid solution. After incubation for 15 min at 90 C, 1.0 mL TCA solution was added. Nash reagent (3.0 mL) was added, the mixture was incubated for 15 min at room temperature and the absorbance measured at 412 nm.

#### **2.4 Determination of iron chelation ability and**

The iron chelating ability was determined based on the assay of (Chew et al. 2009) as modified by (Chai et al. 2014). To 2.0 mL of the seed extract, 0.10 mM FeSO<sub>4</sub> and 0.25 mMferrozine was added and vortexed, allowed to react at room temperature for 10 min and the absorbance measured at 562 nm.

## 2.5 DPPH scavenging ability

The DPPH scavenging activity was determined by adding 0.5 mL of methanolic solution of DPPH to varying volumes of the seed extract. The mixture was vortexed and allowed to stand at room temperature for 30 min and the discoloration of the purple color was measured at 517 nm. Methanol was used as blank and the methanolic DPPH solution as a positive control (Pavithra and Vadivukkarasi 2015).

## 2.6 Statistical analysis

Statistical analysis was carried out using SPSS (IBM, SPSS Statistics 20). The Student's t-test and analysis of variance (ANOVA) were carried out to determine the differences among means at significance level of 0.05%. The reported data are the means of three independent determinations along with the standard deviations.

## 3. Results and Discussion

### 3.1 Steeping in MB enhances germination of seeds

The effect of steeping for 6 h in different concentrations of MB on germination power is shown in Figure 1. After 48 h, the germination power of the control seeds was 73.75, 69.75 and 71.25% for cowpea, millet and sorghum seeds, respectively. Steeping in varying concentrations of MB had differential effects on the germination power of the seeds and as well on seed type. For cowpea (Figure 1A), the lowest (62.50%) and highest (89.75%) GP were observed for seeds steeped in 10 and 80  $\mu$ M MB solutions, respectively. These GP were significantly different compared to the control seeds, i.e. when steeping was done in water. Similarly, the lowest (67.25%) and highest

(82.75%) GP for millet seeds steeped in MB were observed at concentrations of 10 and 60  $\mu\text{M}$  (Figure 1B). Similarly, steeping sorghum seeds in 20 and 80  $\mu\text{M}$  MB resulted in the lowest (67.25%) and highest (82.75%) GP (Figure 1C).

Figure 2 shows the GP as a function of time determined for the different seeds based on the concentration of MB that resulted in the highest GP. It can be observed that the MB primed seeds always had a higher GP than the control seeds at the different time points, although significant differences were observed only beyond 24 h. In cowpea seeds, the control seeds had GP of 39.50 and 73.25% compared to GP of 51.25 and 91.25% for the MB primed seeds after 24 and 48 h of germination, respectively. In both millet and sorghum seeds, higher GP was observed for seeds steeped in MB compared to the control at the different time points. This demonstrates that priming with appropriate concentrations of MB can be useful in enhancing the GP of seeds.

Seed priming has been used to enhance the germination of several seeds (Lutts et al. 2016), including rice (Farooq et al. 2013; Suk-Soon and Jae-Hyeun 2000) and wheat (Basra et al. 2005; Maqsood et al. 2003). Among other functions, it is possible that MB priming better enhances the ability of seed embryos to absorb water to help improve germination. MB priming can also affect germination by enhancing the release of metabolites such as sugars, amino acids, organic acids and ions, as well as repair of genetic materials, resulting in earlier and faster synthesis of DNA, RNA and proteins (Bray et al. 1989). According to Farooq et al. (2013), priming enhances the breakdown of dormancy leading to improved germination of seed.

**Formatted:** Space After: 0 pt, Line spacing: Multiple 1.99 li

**Comment [User7]:** I think that the results obtained in the present work are not well discussed. I think it is necessary to make a stronger connection between your results and the references cited in the text.

### 3.2 Effect on steeping in MB on soluble protein and total phenolic content

The effect of MB priming on soluble protein and total phenolic content is shown in Figure 3. The protein content of the control seeds was 18.03, 6.23 and 8.76 g/100 g DW for cowpea, millet and sorghum seeds, respectively (Figure 3A). MB priming resulted in an increase in the protein content for all seeds compared to the controls (Figure 3A). The increased soluble protein content of MB primed seeds can be attributed to increase in enzyme levels as a result of protein synthesis during germination. According to Hameed et al (2013), this increase in soluble protein shows an improved nitrogen metabolism involving the activity of proteolytic and recycling enzymes (Hameed 2013). MB priming can also ensure proper hydration, which may have resulted in enhanced activity of enzymes that play essential role in germination such as amylases (Umair et al. 2012).

The total phenolic content of the control seeds was 293.09, 205.06 and 607.59 mg GAE /100 g DW for cowpea, millet and sorghum seeds, respectively (Figure 3B). Compared to the controls, MB priming resulted in a significant increase in the total phenolic content of all the seeds (Figure 3B). Germination is an oxidative stress process that is generally associated ROS production (Lutts et al. 2016). Thus, the increased phenolic content can be due to the fact that MB priming causes an increase in the production of compounds, such as phenols, with antioxidant properties to help mitigate the oxidative effects of ROS produced during germination. According to Pal et al. (2017), the increase and possible accumulation of phenolic compounds could be due to both their ROS scavenging ability and the potential to inhibit lipid peroxidation. An increase in phenolic and flavonoids content as well as several photosynthetic pigments was observed following priming of rice seeds (Singh et al. 2016). An increased protein and phenolic content



was also observed in rice seeds primed with selenite (Du et al. 2019), and primed *Moringaoleifera* seeds (Nouman et al. 2014).

**Comment [User8]:** In this item, I also think it is necessary to make a relationship between the results and the references cited. Apparently, results of other research are inserted, but there should be a relationship with the work, and there is not!

### 3.3 Steeping in MB enhances the antioxidant properties of seeds

The effect of MB priming on the antioxidant properties of the seeds is shown in Figures 4 and 5. There was a significant increase in the hydrogen peroxide scavenging ability of the germinated seeds following MB priming (Figure 4A). In cowpea seeds, MB priming increased the hydrogen peroxide scavenging ability by about 38%, while an increase of about 47 and 54% was observed in millet and sorghum seeds, respectively (Figure 4A).

Nitric oxide scavenging ability of the seeds was affected by MB priming. A significant increase of about 31% was observed in MB primed cowpea seeds relative to the control. In millet and sorghum seeds, an increase of 13 and 17% was observed, respectively (Figure 4B). Hydroxyl radical scavenging ability of the seeds also increased upon MB priming (Figure 4C). In cowpea seeds, MB priming significantly increased the hydroxyl radical scavenging ability by about 44%, while an increase of 8 and 14% was observed in millet and sorghum seeds, respectively.

The effect of MB priming on iron chelation ability of the seeds is shown in Figure 5A. An iron chelation ability of 24.75% was observed for the control cowpea seeds. However, upon MB priming, there was an increase of about 6.25% to an overall total of 31%. Increases of about 22 and 24% were observed between the control and the MB primed seeds of millet and sorghum, respectively (Figure 5A). The DPPH radical scavenging activity of all the seeds was affected by

MB priming (Figure 5B). In cowpea seeds, a DPPH radical scavenging activity of 52.75 and 61.25% was observed for the control and MB primed seeds, respectively. Comparing millet and sorghum seeds, an increase of about 11 and 16%, respectively, was observed upon MB priming.

An essential aspect of seed priming is the management of oxidative stress during germination (Lutts et al. 2016). Germination triggers a host of metabolic processes such as mitochondrial respiration and  $\beta$ -oxidation, which lead to increased production of reactive oxygen species (ROS). The accumulation of ROS may affect the germination process and, therefore, increasing the action of antioxidant systems that control ROS levels may help promote germination (Lutts et al. 2016).

Methylene blue has serial conjugate bonds in its structure and so can have the ability to scavenge free radicals thereby possessing antioxidant properties. This attribute of MB has been reported (Atamna et al. 2008), and may help increase the rate at which ROS are scavenged in seeds, particularly during germination. In tomato, a partial protection against abiotic stress root damage was observed following treatment with MB. This protection was attributed to reductions in hydrogen peroxide and malonyldialdehyde levels in the roots (Aloni et al. 2010). MB has also been shown to negate the effect of nitric oxide in maize roots (Gouvêa et al. 1997), and can scavenge nitric oxide radicals. Stohr and Ullrich (2002) demonstrated that MB may have an apoplastic location, where it scavenges NO radicals. In sweet potato tissues, growth stimulation was observed following treatment with MB (Lage and Esquibel, 1997).

**Comment [User9]:** In this item, the comment is the same as the previous one.

#### 4. Conclusions

Selected pulse and cereal (cowpea, millet and sorghum) seeds have been used in this study to demonstrate that priming with the appropriate concentration of MB can enhance the germination

power of seeds. This increase in germination power is accompanied by an increase in soluble protein content that apparently reflect enhanced synthesis of relevant proteins for the needed metabolic efficiency of the emerging plant. ROS are generated as a result of the enhanced metabolic and respiratory activity of the growing plant, but the oxidizing effects are mitigated through accompanying increase in total phenolic content as well as the general antioxidant properties of the germinating seeds.

## References

Aloni B, Karni L, Aktas H. (2010). Methylene blue increases the tolerance of tomato plants to abiotic stresses. *The Journal of Horticultural Science and Biotechnology*, 85:387-393.

Ampofo-Asiama J, Quaye, B. (2018). The effect of pasteurisation on the microbiological and nutritional quality of soursop (*Annona Muricata* L.) juice. *Asian Food Science Journal* 6:1-8.

Atamna H, Nguyen A, Schultz C, Boyle K, Newberry J, Kato H, Ames BN. (2008). Methylene blue delays cellular senescence and enhances key mitochondrial biochemical pathways. *The FASEB Journal*, 22:703-712.

Basra SMA, Farooq M, Tabassam R, Ahmad N. (2005). Physiological and biochemical aspects of pre-sowing seed treatments in fine rice (*Oryza Sativa* L.). *Seed Science and Technology*, 33:623-628.

Boora F, Chirisa E, Mukanganyama S. (2014). Evaluation of nitrite radical scavenging properties of selected Zimbabwean plant extracts and their phytoconstituents. *Journal of Food Processing*, 1-7: ????.

Bray CM, Davison PA, Asharf M, Taylor RM. (1989). Biochemical changes during osmopriming of leek seeds. *Annals of Botany*, 63: 185-193.

Chai T, Mohan M, Ong H, Wong F. (2014). Antioxidant, iron-chelating and anti-glucosidase activities of *Typhadomingensis*Pers (Typhaceae). *Tropical Journal of Pharmaceutical Research*, 13:67-72.

**Comment [User10]:** In conclusion, the authors must correct. In general, it is reported in short sentences, if the objective has been achieved.

**Comment [User11]:** In the references, the authors must follow the rules of this journal, making the necessary corrections.

**Comment [User12]:** Please check that all references cited here are in the text

**Formatted:** Highlight

Chew YL, Goh JL, Lim YY. (2009). Assessment of in vitro Antioxidant capacity and polyphenolic composition of selected medicinal herbs from Leguminosae family in Peninsular Malaysia. *Food Chemistry*, 116:13-18.

Du B, Luo H, He L, Zhang L, Liu Y, Mo Z, Pan S, Tian H, Duan M, Tang X. (2019). Rice seed priming with sodium selenate: Effects on germination, seedling growth, and biochemical attributes. *Scientific Reports*, 9:4311.

Comment [User13]: ????

Formatted: Highlight

El SN, Karakaya S. (2004). Radical scavenging and iron-chelating activities of some greens used as traditional dishes in Mediterranean diet. *International Journal of Food Sciences and Nutrition* 55:67-74.

Farooq M, Irfan M, Aziz T, Ahmad I, Cheema SA. (2013). Seed priming with ascorbic acid improves drought resistance of wheat. *Journal of Agronomy and Crop Science*, 199:12-22.

Fercha A, Capriotti AL, Caruso G, Cavaliere C, Samperi R, Stampachiacchiere S, Laganà A. (2014). Comparative analysis of metabolic proteome variation in ascorbate-primed and unprimed wheat seeds during germination under salt stress.<sup>22</sup> *Journal of Proteomics* 108: 238257.

Comment [User14]: ????

Gouvêa CMCP, Souza JF, Magalhães ACN, Martins IS. (1997). NO<sub>2</sub>-releasing substances that induce growth elongation in maize root segments. *Plant Growth Regulation*, 21:183-187.

Jisha KC, Puthur JT. (2016). Seed priming with BABA (β-amino butyric acid): a cost-effective method of abiotic stress tolerance in *Vignaradiata* (L.) Wilczek. *Protoplasma*, 253:277-289.

Lage CLS, Esquibel MA. (1997). Growth stimulation produced by methylene blue treatment in sweet potato. *Plant Cell, Tissue and Organ Culture*, 48:77-81.

Lutts S, Benincasa P, Wojtyla L, Kubala S, Pace R, Lechowska K, Quinet M, Garnczarska M. (2016). Seed priming: new comprehensive approaches for an old empirical technique, in *New Challenges in Seed Biology-Basic and Translational Research Driving Seed Technology*, eds S. Araujo and A. Balestrazzi (IntechOpen), 1-40.


Comment [User15]: This is a book?? The authors must follow the rules of this journal

Maqsood S, Basra A, Farooq M, Khaliq A. (2003). Comparative study of pre-sowing seed enhancement treatments in fine rice (*Oryza sativa* L.). *Pakistan Journal of Life and Social Sciences*, 1:21-25.

- Nayyer H, Walia DP, Kaistha BL, (1995). Performance of bread wheat (*Triticumaestivum*) seed primed with growth-regulators and inorganic salts. Indian Journal of Agricultural Sciences 65:112-116.
- Nouman W, Basra SMA, Yasmeen A, Gull T, Hussain SB, Zubair M. (2014). Seed priming improves the emergence potential, growth and antioxidant system of *Moringaoleifera* under saline conditions. Plant Growth Regulation, 73:267-278.
- Pal A, Ali MA, Pal AK. (2017). Effect of seed priming on germination behavior, oxidative stress and antioxidant enzyme activities in groundnut (*Arachishypogaea* L) under salinity stress. Bulletin of Environment, Pharmacology and Life Sciences, 6:479-485.
- Pavithra K, Vadivukkarasi S. (2015). Evaluation of free radical scavenging activity of various extracts of leaves from *Kedrostisfoetidissima* (Jacq.) Cogn. Food Science and Human Wellness, 4:42-46.
- Singh A, Gupta R, Pandey R. (2016). Rice seed priming with picomolarrutin enhances rhizospheric *Bacillus Subtilis* CIM colonization and plant growth. PLoS One 11:1.
- Stohr C, Ullrich WR. (2002). Generation and possible roles of NO in plant roots and their apoplastic space. Journal of Experimental Botany, 53:2293-2303.
- Suk-Soon L, Jae-Hyeun K. (2000). Total sugars,  $\alpha$ -amylase activity, and germination after priming of normal and aged rice seeds. Korean Journal of Crop Science 45:108-111.
- Umair A, Ali S, Tareen MJ, Ali I, Tareen MN. (2012). Effects of seed priming on the antioxidant enzymes activity of mungbean (*VignaRadiata*) seedlings. Pakistan Journal of Nutrition, 11:140-144.
- Zheng GH, Wilen RW, Slinkard AE, Gusta LV. (1994). Enhancement of canola seed germination and seedling emergence at low temperature by priming. Crop Science, 34:1589-1593.


Comment [User16]: ???

**Figures**




**Figure 1:** Germination power of cowpea (A), millet (B) and sorghum (C) seeds after 48 h, having previously primed for 6 h in different concentrations of MB.

**Comment [User17]:** Insert the averages in the graph



**Figure 2:** Germination rates of cowpea (A), millet (B) and sorghum (C) seeds primed in MB. The clear bars represent the control seed and the striped bars represent seeds primed in MB solutions that gave the highest germination power.


**Comment [User18]:** Insert the averages in the graph



**Figure 3:** Protein (A) and total phenolic content (B) of germinated cowpea, millet and sorghum seeds. The clear bars represent the control seeds and the striped bars represent seeds primed in MB solutions that gave the highest germination power.


**Comment [User19]:** Insert the averages in the graph





**Figure 4:** Hydrogen peroxide (A), nitric oxide (B) and hydroxyl radical (C) scavenging ability of germinated cowpea, millet and sorghum seeds primed in MB. The clear bars represent the control seeds and the striped bars represent seeds primed in MB solutions with optimum concentration for the specific seeds.

**Comment [User20]:** Insert the averages in the graph



**Figure 5:** Iron chelation ability (A) and DPPH scavenging ability (B) of germinated cowpea, millet and sorghum seeds primed in MB. The clear bars represent the control seeds and the striped bars represent seeds primed in MB solutions with optimum concentrations for the specific seeds.

**Comment [User21]:** Insert the averages in the graph