

Original Research Article

KNOWLEDGE, PRACTICES AND PERCEPTION OF MALARIA AND ITS HOME MANAGEMENT USING ARTEMICININ-BASED COMBINED THERAPY (ACT) AMONG MOTHERS OF UNDER-FIVE

ABSTRACT

Malaria is a serious public health problem, yet preventable and treatable. The disease is one of the world's highest rates of all cause of mortality for children under five, and about one in six children die before their fifth birthday. Hence, mothers of under-five and caregiver have a pivotal role to play in tackling this issue by improving their knowledge and skills concerning the treatment, prevention, and control using the appropriate approach. This study was carried out to assess knowledge, practices, and perception of malaria and its home management using Artemisinin-based Combined Therapy (ACT) in Yemetu community of Ibadan North Local Government. The study was a descriptive cross-sectional survey involving the use of Expanded Programme on Immunization (EPI) to facilitate the sampling and interview of respondents. This included recruiting all the mothers of under-five in Yemetu community who gave consent for the study. Four hundred (400) mothers of under-five in Yemetu community consented to participate in the study and were selected. A validated semi-structured questionnaire interviewed and self-administered questionnaire was used for data collection and respondents were assessed on a 62-points knowledge scale, 5-points practice scale, and 17-points perception scales. Knowledge score ≤ 21 were rated poor, scores $\geq 22 \leq 42$ fair and scores ≥ 43 were considered good. Practice score ≤ 3 was recorded as poor practice while scores ≥ 3 good practice. Perception scores ≤ 9 were considered unfavorable perception and scores ≥ 9 were considered favorable. Descriptive statistics and ~~chi~~ Chi-square tests were used to ~~analyse~~ analyze the data at 95% level of significance.

Respondents' mean age was 29.9 ± 7.0 years and the majority of them (91.1%) of them were Yoruba. ~~Majority~~ The majority (91.0%) of them were married and (91.1%) were Yorubas. Only (23.0%) correctly identified plasmodium as a cause of malaria. The correctly mentioned signs and symptoms of simple malaria were; cold (89.3%), body ache (91.3%) and fever (88.5%). The fairly corrected home management practice steps include; Exposure of baby to fresh air, administration of paracetamol, and then provision of coartem (2.6%) and bathing the baby, use of paracetamol and administration of coartem (1.3%). Negative perception shown by the

34 | respondents include: Only (15.0%) believed that malaria is a disease of the poor and preference
35 | of herbal medicine to medical medicine for treating children at home when they have malaria
36 | episode because it is cheaper (19.8%). Overall, (2.8%) had poor knowledge, the majority
37 | (87.3%) had a fair knowledge, and 9.3% had good knowledge. -There are several gaps in the
38 | respondents' knowledge relating to malaria and its management in under-five. Therefore, there is
39 | a need for peer education/training approach in this regard to upgrade mothers' knowledge and
40 | skills concerning the treatment, prevention, and control of malaria.

41

42 | **Keywords:** Under-five, Home management of malaria, Artemisinin Combination Therapy
43 | (ACT)

44 | 1.0 INTRODUCTION

45 | Malaria remains one of the world's greatest childhood killers and is a substantial obstacle to
46 | social and economic development in the tropics It is a major cause of morbidity and mortality
47 | especially among the vulnerable groups to which children, especially aged less than 5 years
48 | belong. (Idro, Otieno, White, Kahindi, Fegan, Ogutu, Mithwani, Maitland, Neville, and Newton
49 | C.R., 2007). It was observed that malaria accounts for 25 per-cent of infant mortality and 30
50 | percent of childhood mortality in Nigeria thereby imposing great burden on the country in terms
51 | of pains and trauma suffered by its victims as well as loss in outputs and cost of treatments
52 | (WHO, 2000).

53 | The parasite responsible for these deaths—*Plasmodium falciparum*—is transmitted to people
54 | when they are bitten (usually at night) by an infected mosquito. In the human body, the parasites
55 | reproduce in the liver before invading red blood cells. Here, they multiply again before bursting
56 | out and infecting more red blood cells as well as causing a high fever and sometimes damaging
57 | vital organs. The transmission cycle is completed when a mosquito bites an infected person and
58 | ingests parasites with its blood meal.

59 | To reduce the global burden of malaria, this cycle needs to be broken. This can be done in
60 | several ways. First, mosquitoes can be controlled with insecticides. Second, individuals can
61 | avoid mosquito bites by sleeping under insecticide-treated nets. Finally, antimalarial drugs can
62 | be used to reduce the illness and death caused by the malaria parasite and can lessen the
63 | likelihood that a mosquito will pick up the parasite when it bites a person- (WHO, 2005).
64 | Regardless of the fact that it is one of the oldest recorded diseases, malaria remains one of the
65 | world's most deadly infectious diseases. It is arguably, the greatest menace to modern society in

Formatted: Font: Italic

66 | terms of morbidity and mortality. Though preventable, treatable, and curable, there is no known
67 | immunity. Several centuries after its discovery, malaria still remains a devastating human
68 | infection, resulting in 300-500 million clinical cases and three million deaths every year (WHO,
69 | 2005).

70 | It is also believed to contribute up to 11 percent maternal mortality, 25 percent infant mortality,
71 | and 30 percent under-five mortality. It is estimated that about 132 billion Naira lost to malaria
72 | annually in the form of treatment costs, prevention and loss of work time in Nigeria (FMOH and
73 | NMCP, 2009)

74 | Nigeria is known for the high prevalence of malaria and it is a leading cause of morbidity and
75 | mortality in the country. Available records show that at least 50 per-cent of the population of
76 | Nigeria suffers from at least one episode of malaria each year and this accounts for over 45 per
77 | cent of all outpatient visits. (Ojurongbe, Ogungbamigbe, Fagbenro- Beyioku, Fendel, Kremsner,
78 | and Kun -2007).

79 | Malaria is known to have a negative impact on performance and learning in children according
80 | to Holding and Snow (2001). It also aggravates anaemia and malnutrition in children and
81 | pregnant women. (Murphy -and Breman 2001).

82 | There are strategies being promoted for the management of malaria as a result of the emergence
83 | of chloroquine resistance aimed at preventing the occurrence of malaria. The World Health
84 | Organization (WHO) currently recommends Artemisinin-based Combination Therapies (ACTs)
85 | for malaria control. The use of insecticide-treated nets is also now being strongly promoted.
86 | (WHO, 2006). Arigbabuwo, (2010) in his study also opined that prevention is better than cure,
87 | advising that people should learn to maintain personal and environmental hygiene.

88 | Mothers have a crucial role to play in recognition, treatment, and prevention of malaria in under-
89 | five. The knowledge, perception and skills and practices relating to malaria among mothers of
90 | under-five in urban settings are yet to be well investigated. -This study, therefore, focuses on
91 | knowledge, practices, and perception of malaria and its home management using Artemisinin
92 | Combination Therapy (ACT) among mothers of under-five in Yemetu one of the communities in
93 | Ibadan metropolis.

94 | 2.0 Materials and Method

95 The study was a descriptive cross-sectional survey involving the use of Expanded Programme on
96 Immunization (EPI) to facilitate the sampling and interview of respondents. This included
97 recruitment of all the mothers of under-five in Yemetu community who gave consent for the
98 study. Yemetu is a community located in Ibadan North Local Government Area (LGA) in ward
99 3- of Oyo State, Ibadan. South Western Nigeria. With ward number (3). It has a total population
100 of 65,949. Four hundred (400) mothers of under-five in Yemetu community however, consented
101 to participate in the study and were selected. The community had three (3) health facilities: A
102 ~~government~~ government-owned secondary health facility called Adeoyo Maternity Teaching
103 Hospital, and two private health facilities namely Kola Daisi Foundation Center (for primary and
104 community health) and Vine Branch Medical Clinic which are accessible to the people in the
105 community. The total number of under-five children is 13,190- (National Population
106 Commission, 2006-). The community is heterogeneous consisting of people from different part
107 of the country. The Yorubas however constituted the majority and their major occupation is
108 trading.

109

110 **2.1 -Target Population**

111 The populations for this study were mothers of under-five children residing in Yemetu
112 Community of Ibadan North LGA in Oyo State, South West of Nigeria who merited the
113 inclusion criteria for the study.

114

115 **2.2 -Sampling Procedure**

116 The Expanded Programme on Immunization (EPI) sampling technique was used to facilitate the
117 sampling and interview of the respondents. The investigator started data collection by moving to
118 the center of Yemetu community and spinned a bottle. The spinned bottled was allowed to turn
119 round and round unhindered and allowed to come to rest. ~~Interview~~ The interview started from
120 the part of the community to which the mouth of the bottle was pointing. Every third house in the
121 direction was selected and visited and one eligible respondent was selected by balloting for
122 interview if more than one eligible respondent was met in a house. In a house where there was
123 one mother, such a mother was purposively selected for interview if she consented to participate
124 in the study.

125 After reaching the end of the community, the investigator and the research assistants moved back
126 to the center of the community and started recruitment and interview in another direction. This

127 way a total of 422 eligible mothers of under-five who consented to be involved in the study were
128 interviewed.

129 **2.3 Ethical Consideration**

130 All interviews ~~was-were~~ conducted in compliance with the ethics of the health promotion and
131 education profession. Copies of the research proposal were submitted to Oyo state Ethical
132 Review Committee for approval before the study commenced. This was done in order to ensure
133 that the study was conducted in accordance with ethical principle covering studies involving
134 human objects. The research assistants were well trained to obtain informed consent for
135 respondents before ~~the~~ interview. Respondent ~~were-was~~ informed on the purpose of the study
136 and ~~were-was~~ given ~~the~~ option to participate through written or verbal consent or withdraw from
137 participating. Informations provided by the respondent ~~were-was~~ treated with confidentiality.
138 ~~Registration-The registration~~ number was assigned to each questionnaire, no identifiers such as
139 names, address or phone numbers ~~was-were~~ required on the questionnaire.

140 **2.4 Data collection procedure:**

141 A semi-structured questionnaire was used for data collection. The semi-structured interviewer
142 and self-administered questionnaire ~~was-were~~ divided into six sections labeled sections A, B, C,
143 D, and E consisting of ~~open-open~~-ended and ~~close-close~~-ended questions. It was developed from
144 ~~the~~ literature review and adapted questions from related past studies in Nigeria. The structured
145 questionnaire comprised of open-~~ended~~ and ~~close-close~~-ended questions which were used to
146 elicit information on menace of malaria disease, home management and practices, and
147 perception. The Yoruba version of the questionnaire was produced after necessary modification
148 to the English version had been done. It consisted of 28 questions divided into five sections. The
149 questionnaire was validated by the researchers and experts in Public Health they included,
150 pediatricians, statisticians working on malaria control in university of Ibadan. -It was pretested in
151 Ekotodo community in Ibadan North LGA a similar community in Ibadan North LGA. The data
152 were then subjected to descriptive statistics which was basically frequencies and charts.
153 ~~Reliability-The reliability~~ coefficient obtained was determined using the Cronbach's Alpha
154 technique. Any ~~coefficient~~~~coefficient~~ >0.5 is said to be reliable. In the study, reliability
155 coefficient score which is also called chronbach Alpha was calculated to be of 0.733.

156 **3.0 RESULTS**

Comment [DM1]: What type of statistical treatment was applied to this study?

Formatted: Font: Bold

157 | **3.1 -Socio-demographic demographic characteristics of respondents**

158 Table 1 presents the socio-demographic characteristics of the respondents. Respondents within
159 the age group 20-29 years constituted the highest (41.8%) followed by those aged 30-39 (40%).
160 Respondents' aged less than 20 years were the least (6.0%). The mean age of the respondents
161 was 29.9 ± 7.0 years.

162 Most of the respondents (90.8%) were married. Respondents with secondary school (56.5%)
163 topped the list of [the](#) highest level of education. While those with tertiary accounted for (23.8%)
164 of the respondents.

165 Over half of the respondents were traders (57.8%). While artisans constituted (31.3%).
166 Respondents in monogamous constituted the majority (71.0%). (See table 1 for details)

167
168

169 **Table 1: Socio-demographic characteristics of the respondents**

170

Socio-demographic characteristics	Frequency	Percent (%)
Age in years: (n=397)*		
Less than 20 years	24	6.0
20-29 years	166	41.8
30-39 years	161	40.6
40-49 years	46	11.6
Marital status: (n=393)		
Single	29	7.3
Married	364	91.0
Religion: (n=400)		
Christianity	207	51.8
Islam	190	47.5
Traditional	3	0.75
Ethnic group: (n=395)		
Yoruba	360	91.1
Igbo	23	5.8
Hausa	12	3.0
Highest level of education: (n=400)		
Primary	79	19.8
Secondary	226	56.5
Tertiary	95	23.8
Type of tertiary education: (n=75)		
University	24	32.0
Polytechnic	39	52.0
Diploma/nursing	12	16.0

Occupation : (n=396)

Trading	229	57.8
Civil servant	41	10.4
Artisan	124	31.3
Unemployed	2	0.4

Family type: (n=387)

Polygyny	112	29.0
Monogamous	275	71.0

Children aged less than five: (n=397)

One	262	66.0
Two	129	32.5
Three	6	1.5

171

172 *Mean age: 29.9± 7.0

173

174

175

176

177

178 **Table 2: Pattern of use and storage of antimalarial drugs and related medicine for treating**
 179 **under-fives**

Pattern of use of antimalarial	No	%
Ever used an antimalarial to treat under-five (N=398)		
Yes	357	89.7
No	41	10.3
Types of malarial medicine used(N=399)		
Coartem *	268	67.2
Artesunate*	65	16.3
Chloroquine -	28	7.0
Paracetamol -	18	4.5
Ampiclox -	13	3.3
Alabukun-	7	1.8
Pain relieving medicine normally used for treating underfive at home in case of malaria(N=399)		
Paracetamol*	322	80.7
Novagen-	47	11.8
Ibuprofen -	22	5.5
Alabukun -	8	2.0

Places where antimalarial medicines are kept at home(N=399)

Cool dry place*	287	72.1
Inside nylon +	60	15.1
Inside wardrobe+	32	8.0
In the kitchen -	19	4.8

180 **Malaria treatment seeking pathways for under-five during episodes of malaria preceding**
181 **study.**

Where sought treatment	No	%
Hospital*	156	39.1
Patent Medicine Vendors (PMV) ±	120	30.1
Health centre*	50	12.5
Private clinic*	37	9.3
Primary Health Care (PHC)*	30	7.5
Community Medicine Distributors (CMDs)+	6	1.5

182 **Respondents step by step home management of malaria involving under-five child**

Steps taking at home	No	%
Use paracetamol for the baby+	177	46.7
Bath the baby±	52	13.7
Bath the baby and use PCM for the baby+	55	14.5
Use agbo for the baby±	44	11.6
Mop the body with cloth soaked in cold water±	11	2.9
Bath the baby, use paracetamol for him/her and take him/her to the hospital*	25	6.6
Expose to fresh air, give paracetamol and give coartem*	10	2.6
Bath for the baby, use paracetamol and give the baby coartem*	5	1.3

183

184 **3.2 Respondents' practices related to home management of malaria**

185 Respondents pattern of use and storage of antimalarial drugs and related medicine for treating
186 under-fives is highlighted in table 2. ~~Majority~~ The majority (89.7%) of the respondents had used
187 an antimalarial drug to treat their under-five child (ren) while only 10.3% had never used
188 antimalarial drugs. The antimalarial drug normally used included; coartem (34.2%), artesunate
189 (29.0%), amalar (24.3%), while (10.5%) listed fansidar. Respondents were further asked about
190 the type of malarial medicine used. Respondents that used coartem (67.2%) topped the list.
191 Respondents that used paracetamol for pain relieve in treating under-five at home had the highest
192 proportion of (80.7%) A majority (72.1%) kept their related malaria medicine in a cool dry
193 places. (See table 2 for details).

194 Table 2 highlighted malaria ~~treatment-treatment~~-seeking pathways for under-five during
195 episodes of malaria preceding the study. The highest proportion (39.1%) sought for the treatment

196 in a hospital, followed by Patent Medicine Vendors (PMVs) (30.1%). The other listed places are
 197 contained in the table under reference. Respondents' step by step home management of malaria
 198 involving under-five children is specified in table 2. The fairly correct steps mentioned were as
 199 follows: Exposure of baby to fresh air, administration of paracetamol, and then provision of
 200 coartem (2.6%) and bathing the baby, use of paracetamol and administration of coartem (1.3%)
 201 (See table 2 for detail). Categorization of overall respondents' practice score was assessed using
 202 a 5-point scale. Respondents with good practice (4-5points) constituted 94.0%, while the
 203 proportion of respondents with poor practice (0-3) accounted for 6.0%.

204 **Table 3: Respondents' knowledge of causes of malaria**

Causes of malaria [^]	True (%)	False (%)	Don't know (%)	Total
Mosquito	392(98.0)	8(2.0)	0(0%)	400
Too much sun	221(55.8)	175(44.2)	4(1.0)	396
Change of weather	97(24.3)	281(70.3)	22(5.5)	400
Plasmodium	92(23.0)*	253(63.3)	55(13.8)	400
Taking too much palm oil	141(35.3)	230(57.5)	29(7.3)	400
Overwork/too much work	185(46.3)	199(49.8)	16(4.0)	400
Witchcraft	76(19.0)	292(73.2)	31(7.8)	399

Comment [DM2]: Please add footnotes in every Tables.

205
206
207
208
209
210
211
212

Table 4: Respondents' knowledge of factors or condition that can make mosquito breed or multiply

Factors/conditions	True (%)	False (%)	Don't know (%)
Blocked gutters/drains with water	388(97.0)*	11(2.8)	1(0.3)
Improper refuse disposal	385(96.3)*	15(3.8)	0(0)
Stagnant water	340(85.0)*	58(4.5)	2(0.5)
Empty containers or vessels (e.g. bottles, cans, plastics etc.)	217(54.3)*	162(40.5)	21(5.3)

Engine oil in a container that is not covered+	120(30.0)	250(62.5)	30(7.5)
Stagnant water containing spent engine oil+	122(30.5)	246(61.5)	32(8.0)

213

214

215 **Table 5: Respondents' knowledge of signs and symptoms of simple malaria**

Signs and symptoms of simple malaria	Responses		
	Correct (%)	Wrong (%)	Don't know (%)
Inflammation of the skin	169(42.3)	175(43.8)	56(14.0)
Fever	354(88.5)*	38(9.5)	8(2.0)
Nausea+	347(86.8)	45(11.3)	8(2.0)
Diarrhoea	247(61.8)	136(34.0)	17(4.3)
Vomiting	320(80.0)*	71(17.8)	9(2.3)
Cold	357(89.3)*	36(9.0)	7(1.8)
Tiredness	358(89.5)*	34(8.5)	8(2.0)
Catarrh	365(91.3)*	34(8.5)	1(3)
Body ache	365(91.3)*	32(8.0)	3(8)
Itching +	264(66.0)	125(31.3)	11(2.8)
Fatigue	310(77.5)*	75(18.8)	15(3.8)
Sore throat+	230(57.5)	148(37.0)	22(5.5)

216

217 **Table 6: Respondents' knowledge of signs and symptoms of severe malaria**

Symptoms of severe malaria^	Responses			Total
	Correct (%)	Wrong (%)	Don't know (%)	
Fever	361(90.5)*	25(6.3)	13(3.3)	399(100%)
Chills	352(88.0)*	31(7.8)	17(4.3)	400(100%)
Organs dysfunction	154(38.5)*	126(31.5)	120(30.0)	400(100%)
Abnormal bleeding	104(26.0)	167(41.8)	129(32.3)	400(100%)
Clinical jaundice	140(35.0)*	129(32.3)	131(32.8)	400(100%)

Febrile convulsion	175(43.8)*	107(26.8)	118(29.5)	400(100%)
Respiratory distress	130(32.5)*	119(29.8)	151(37.8)	400(100%)
Impaired consciousness	121(30.3)*	118(29.5)	161(40.3)	400(100%)

218

219 **Table 7: Respondents knowledge of preventive measures against malaria**

Preventive measures	True (%)	False (%)	Don't know (%)	Total
Using insecticide treated net	367(91.8)*	31(7.8)	2(.5)	400(100%)
Eating balanced diet	218(54.5)+	171(42.8)	9(2.3)	389(100%)
Clearing of residential environment of grasses/overgrown weeds	322(80.5)*	77(19.3)	1(.3)	400(100%)
Clearing blocked gutters	311(77.8)*	87(21.8)	2(.5)	400(100%)
Bathing daily	136(34.0)+	248(62.0)	15(3.8)	400(100%)
Use of insecticide	292(73.0)*	97(24.3)	11(2.8)	400(100%)
Use of antimalarial drug(SP) by pregnant women	199(49.8)*	175(43.8)	26(6.5)	400(100%)
Having enough sleep	87(21.8)+	283(70.8)	30(7.5)	400(100%)
Not eating too much palm oil	98(24.5)+	272(68.0)	30(7.5)	400(100%)
Not working in the sun for a long time	90(22.5)+	273(68.3)	37(9.3)	400(100%)

220

221

222

223 **Table 8: Respondents' knowledge of malaria related treatment actions involving under-five children treatment**

224

Treatment steps actions^	True (%)	False (%)	Don't know (%)	Total
Tepid sponging	363(92.2)*	25(6.3)	10(2.5)	398
Use of paracetamol	384(96.2)*	14(3.5)	1(0.3)	399
Use of coartem	363(91.0)*	22(5.5)	14(3.5)	399
Use of agbo	267(66.9)	125(31.3)*	7(1.8)	399
Use of chloroquine	256(64.2)	132(33.1)*	11(2.8)	399

Going to a health care facility for treatment 365(91.5)* 27(6.8) 7(1.8) 399

225


226

227 **Table 9: Respondents' knowledge of coartem and paracetamol dosage regimen for children**
 228 **aged 0-5years**

Ages(years)	Coartem dosage regimen [^]	Right	Wrong	Don't know	Total
6months-3years	1 tablet twice daily(3days)*	248(62.2)	93(23.3)	58(14.5)	399
	2 tablet twice daily(3days)	55(13.9)	280(70.7)	61(15.4)	396
	1 tablet thrice daily(3days)	33(8.3)	300(75.8)	63(15.9)	396
3-5years	1 tablet twice daily(3days)	100(25.3)	246(62.1)	50(12.6)	396
	2 tablet twice daily(3days)*	200(50.1)	150(37.6)	49(12.2)	399
	3 tablet thrice daily(3days)	55(13.9)	290(73.2)	51(12.9)	396
Paracetamol dosage regimen					
6months-3years	1/2 tablet twice daily(3days)*	256(64.0)	93(23.3)	51(12.8)	400
	1 tablet twice daily(3days)	74(18.6)	270(68.0)	53(13.4)	397
	1/2 tablet thrice daily(3days)	38(9.6)	308(77.6)	51(12.8)	397
3-5years	1 tablet twice daily(3days)	190(47.9)	169(42.6)	38(9.6)	397
	1 tablet thrice daily(3days)*	143(35.8)	219(54.8)	38(9.5)	400
	1 tablet once daily(3days)	48(12.1)	306(77.1)	42(10.6)	397

229

230 The knowledge of treatment/dosage regimen for coartem and paracetamol for children aged 0-5
 231 years is summarized in table 9. More than half of the respondents (62.2%) were knowledgeable
 232 about the correct treatment/dosage of coartem for children ages 6 months – 3 years. About half
 233 (50.0%) were conversant with the correct treatment/dosage regimen for coartem for children
 234 | aged 3-5 years. ~~Majority~~ The majority (64.0%) knew the correct treatment/dosage regimen of
 235 paracetamol for children aged 6 months – 3years while only (35.8%) were knowledgeable of the
 236 correct treatment/dosage regimen of paracetamol for children aged 3–5 years. (See table 9 for
 237 details).



238

239 Figure: 1 Categorization of respondents' knowledge scores relating to general knowledge of
 240 malaria and home management of malaria.

241 **Table 10: Respondents perception relating to vulnerability to seriousness and of treatment**
 242 **of malaria.**

Perception	Agree	Undecided	Disagree
Perception relating to vulnerability			
My child is not prone to malarial so no need of taking preventive measures	11(2.8)±	4(1.0)	385(96.3)+
I believe malaria is a disease of the poor, our child cannot get it because we are not poor	60(15.0) ±	3(0.8)	337(84.3)+
I make sure my child stays away from people or other children having malaria to avoid getting it.	55(13.8) ±	9(2.3)	336(84.0)+
I am of the opinion that a child that is well fed cannot have Malaria	50(12.5) ±	12(3.0)	338(84.5)+
Perception relating to seriousness			
I do not believe malaria is a serious disease for children	12(3.0) ±	4(1.0)	384(96.0)+
Malaria cannot lead to death of children aged less than five years	7(1.8) ±	9(2.3)	384(96.0)+
Perception relating to treatment			
Malaria infection is caused by witches and wizards, so telling me about using drugs to treat it is a waste of time	6(1.5) ±	28(7.0)	366(91.5)+
I believed malaria infection will disappear on its own without treatment/medicine	11(2.8) ±	12(3.0)	377(94.3)+

243

244

245 **Table 11: Respondents' perception relating to home management of malaria**

Perception	Agree	Undecided	Disagree
I believe chloroquine alone is enough to treat my child of any kind of malaria at home	20(5.0) ±	52(13.0)	328(82.0)+
I am of the opinion that coartem should be used at home only when the child's malaria is serious	27(6.8) ±	68(17.0)	305(76.3)+
Malaria infection in a child is best treated at home with chloroquine than Arthemisinin-based Combined Therapy (e.g coartem, artesunate etc)	24(6.0) ±	70(17.5)	306(76.5) +

The first dosage of malaria drug is enough to treat children when they have malaria	9(2.3) ±	47(11.8)	344(86.0) +
I prefer herbal medicine to medical medicine for treating my child at home when he/she has malaria because it is cheaper for treating under-five with malaria	79(19.8) ±	68(17.0)	253(63.3) +
Traditional medicine used at home is more effective for treating malaria in children aged less than five years	82(20.5) ±	67(16.8)	251(62.8) +
It is better to wait for a day or two to see whether an under-five has malaria before treating him/her at home with malaria medicine	105(26.3) ±	24(6.0)	271(67.8) +
Every mother should keep medicine at home for the home management of malaria when the need arises	301(75.3) ±	9(2.3)	90(22.5) +
It is wrong for a mother to treat her under-five children at home in case of malaria	133(33.3) ±	8(2.0)	259(64.8) -

246

247 3.3 Perception relating to malaria

248 Respondents in this study had a good perception of home management of malaria.
 249 Majority of the respondent believed that ACT is best used in treating under-five children at home
 250 when they have malaria. A similar study was carried out by Ajayi, and Falade (2006); Salako,
 251 Brieger, Afolabi *et al* (2001) where respondents use chloroquine, and
 252 sulphadoxime/pyrimethamine(SP) at home for the treatment of malaria.

Comment [DM3]: A complete discussion is desirable with adequate representation and comparison of data

Formatted: Font: Italic

253

254 4.0 Conclusion

255 This study revealed that the level of awareness and knowledge of malaria among respondents
 256 was fair. However, there are several gaps in the respondents' knowledge relating to the disease
 257 and its management in under-five. The respondents had a poor knowledge of the cause and fair
 258 knowledge of the factors that could promote the breeding of malaria. Advocacy, training, and
 259 public enlightenment are necessary to address the situation.

Comment [DM4]: The conclusion do not matches with the exact outcome of the study. Please be focused, talk on the outcomes, and share your own views prospectively.

260 5.0 Recommendations

261 The recommendations based on the findings of this study are as follow:

262 1. Sustained public enlightenment interventions relating to malaria targeted at mothers of under-
 263 five are needed. These interventions should be aimed at improving their knowledge and their
 264 malaria prevention and control skills.

265 2. Artemisinin Combination Therapy (ACT) is a new strategy for managing malaria.
 266 Training is needed to improve their knowledge and skill relating to the approach.

267 3. Formal health care facilities are commonly used by the residents for the management of
268 malaria in under-five. The capacity of health workers should be enhanced to help upgrade
269 mothers' knowledge and skills relating to [the](#) correct treatment regimen for managing malaria.

270 4. Training on home management of malaria should be organized for respondent. A peer
271 education approach should be used in this regard to upgrade mothers' knowledge and skills
272 concerning the treatment, prevention, and control of malaria.

273 6.0 REFERENCES

274 Ajayi IO. (2006): Development and Assessment of a treatment guideline to improve home
275 management of malaria in children in Ona-Ara local Government, Oyo state, Nigeria.
276 *[Ph.D. dissertation]. Ibadan, Nigeria: University of Ibadan;*

277 Arigbabuwo Adeleye (2010). Malaria: Killer at large. Vanguard Newspaper special report

278 Eckert E (Ed.); Scaling up home-based management of malaria – from research to
279 implementation. World Health Organisation, Geneva, 2004:2

280 Falade C.O., Ajayi, I.O., Yusuf, O.B., Pagnoni,F.(2014) 'High acceptance of artemisinin based
281 combination therapy for the home management of malaria in rural communities in
282 southwest Nigeria' *Malaria World Journal;Vol. 5: 5*

283 Falade C O, Ogundiran M O, Bolaji M O (2006): The influence of cultural perception of
284 causation, complication, and severity of childhood malaria on determinants and
285 treatment preventive pathways. *International Quarterly of Community Health*
286 *Education, a Journal of Policy and Applied Research 2006. 24.4: 347 – 363.*

287 FMOH (2009). Federal Republic of Nigeria training manual for management of malaria in
288 Nigeria Participants' Manual Federal Ministry of Health National Malaria and Vector
289 Control Division, Abuja- Nigeria

290 Holding P.A., Snow R.W., (2001). "Impact of Plasmodium falciparum malaria on performance
291 and learning: review of the evidence". *Am. J. Trop. Med. Hyg.* 64 (1-2 Suppl): 68–75.

292 Idro, R, Otieno G, White S, Kahindi A, Fegan G, Ogutu B, Mithwani S, Maitland K, Neville
293 BG, Newton CR. (2007). "Decorticate, decerebrate and opisthotonic posturing and
294 seizures in Kenyan children with cerebral malaria". *Malaria Journal 4 (57). Online*
295 *biomed Central. 9*

296 Murphy S.C., and Breman J.G. (2001). Gaps in the childhood malaria burden in Africa: cerebral
297 malaria, neurological sequelae, anemia, respiratory distress, hypoglycemia, and
298 complications of pregnancy. *Am. J. Tropical Med. and Hyg.* 64(1-2 Suppl):57-67

Comment [DM5]: References are not uniform and some references are not complete

299 Ojurongbe O, Ogungbamigbe TO, Fagbenro- Beyioku AF, Fendel R, Kremsner PG, Kun JF
300 2007. Rapid detection of Pfcrt and Pfmdr1 mutations in Plasmodium falciparum isolates
301 by FRET and in vivo response to chloroquine among children from Osogbo, *Nigeria*.
302 *Malar J* 6: 41. 3

303 Salako, L.A., Brieger, W.R., Afolabi B.M. (2001).Treatment of childhood fever and other illness
304 in the three rural Nigerian communities. *JTrop pediatr*.

305 WHO (2000) The Abuja declaration and the plan of action. Geneva: World Health Organization.
306 WHO/CDS/RBM/2000.17. Available: [http://](http://www.rbm.who.int/docs/abuja_declaration.pdf)
307 www.rbm.who.int/docs/abuja_declaration.pdf. Accessed 10 March 2007.

308 WHO 2006 Calls for an immediate Halt to provision of single drug artemisinin Malaria Pills
309 WHO recommendations. 2005 , RBM Department; WHO Geneva

UNDER PEER REVIEW