# <sup>1</sup> Original Research Article Validation Of Efficiency Method For Heavy Metals Determination In Kola Nuts (*Cola nitida* Schott & Endl.) From Côte d'Ivoire

#### 9 ABSTRACT

**Aims:** The current study targets the achievement of a reliable process for the determination of heavy metal contents in kola nuts, namely cadmium, mercury, and lead, for better appreciation of the risks incurred from the consumption of such food products.

**Study design:** kola nuts collected from different stakeholders (planters, collectors, stores and centers) were analyzed after the validation of the proposed analytical method.

**Place and Duration of Study:** Central Laboratory for Food Hygiene and Agro-Industry, LANADA in Abidjan, Côte d'Ivoire, running 2018.

**Methodology:** Two references were used for the validation of the analytical method, namely the French standard NF V 03-110 and the European directive 2001/22/EC. The assays were achieved with an flame Atomic Absorption Spectrophotometer (AAS). The heavy metal contents of some samples collected from different sampling place were then determined

**Results:** From the data, a significant regression chart was recorded for the heavy metals detection graphs, with significant correlation coefficients (R<sup>2</sup>> 0.99). The linearity domain was validated between 0.5  $\mu$ g/L and 1.5  $\mu$ g/L for cadmium, 15  $\mu$ g/L and 45  $\mu$ g/L for lead and from 10  $\mu$ g/L to 100  $\mu$ g/L for mercury. In addition, the LOD were 0.03  $\mu$ g/L, 1.85  $\mu$ g/L and 2.92  $\mu$ g/L, while the LOQ were 0.07  $\mu$ g/L, 6.52  $\mu$ g/L and 3.32  $\mu$ g/L for cadmium, lead and mercury, respectively. The relative standard deviations of the repeatability and reproducibility assays are below 4%, whereas standard additions of heavy metals are fully recovered, with percentages close to 100%. Contents of cadmium, lead and mercury in kola nuts are respectively valued at 22.97±9.01  $\mu$ g/kg, 1065.57±613.76  $\mu$ g/kg and 33.88±31.58  $\mu$ g/kg from the farmers and 24.99±7.79  $\mu$ g/kg, 296.51±98.18  $\mu$ g/kg and 39.74±34.66  $\mu$ g/kg from the storage centres.

**Conclusion:** This analytical method could help in ensuring effective sanitary control at different critical points of kola nut distribution channel for promoting a good management of the toxicity concerns in such products

10

12

<sup>11</sup> Keywords: Cola nitida, heavy metals, Cadmium, Lead, Mercury, validation method.

#### 13 **1. INTRODUCTION**

- Native from the tropical forests of Western Africa, *Cola* genus (*Sterculiaceae*) includes about 40 species and the nuts for the commercial extract are derived, almost exclusively, from two species of Cola, either *Cola nitida* (Vent.) Schott and Endl. or *Cola acuminata* (Beauv.) Schott and Endl [1, 2, 3]. In Côte d'Ivoire, the most common cola crops are from *Cola nitida* species generally grown in the districts of the Mountains, Comoe, Lagoons and Down-Sassandra [4]. The cola plant greatly grows with annual rainfall over 1000 mm and on soft and well-drained soils [5]. The main interest of the cola crop lies in the production of fruits known as kola nuts
- According to Asogwa [6], one of the major constraints of the cola cultivation lies in the infertility of the soils. So, for improving the yield, numerous farmers usually use chemical fertilizers and pesticides [7]
- which can reduce both the plant's strength and the fruits quality in long-range. Of course, this practice really succeeds in improving the growth and yield of the plant [8]. Unfortunately, it also brings toxic elements into the crops [9].
- 26 Otherwise, the post-harvest preservation of the raw crops is another significant constraint for the cola
- stakeholders [1, 10]. Indeed, kola nuts are generally consumed fresh [11]. Yet, the fresh crops state easily allows proliferation of microbes, ants and other parasites. In order to control the crops postharvest enemies and to keep the fruits fresh, the farmers and traders generally soak the raw kola nuts
- in organic pesticides solutions [12, 13].
- 31 The use of chemicals in the cola sector is observed in the planted soils and during the crops carriage
- and processing. Indeed, the kola nuts distribution channel is generally from farmers to the big storage,
   processing and export centres, with temporary stay from rural collectors and small urban stores [4].
- During their processing, carriage and sale, heavy metals could be laid on the kola nuts stock [14]. According to Nordström [15] and Vine [16], the use of pesticides results in negative impact on the
- According to Nordstrom [13] and vine [16], the use of pesticides results in negative impact on the
   environment and human health. Furthermore, the works of Biego et al. [13] and Aikpokpodion et al. [3]
   showed the presence of organochlorine pesticides in kola nuts at concentrations over the maximal
   values admitted by the *Codex Alimentarius*.
- According to Adeosun et al.[8], 90% kola nut production is daily consumed by the populations during
- 40 native ceremonies such as weddings, baptisms, friendly meeting, funeral and the sacrifices rituals [17, 18, 19]. This high consumption is due to the alleged properties of nuts in particular stimulation of the nervous system, energy, and dietary properties [20, 21]. Some Studies highlighted correlations between such properties and the large amount of alkaloids, polyphenolics and carbohydrates compounds in kola nuts [22, 23]. However, accounting the various anti-nutrient substances as pesticides, mycotoxins, and heavy metals also found in kola nut, the consumption of this raw product is a source of public health toxicity concerns [13, 24].
- The heavy metals, namely lead (Pb), cadmium (Cd), and mercury (Hg) are known as strict and toxic contaminants for living beings, even at very lower concentrations [25, 26]. The accumulation in the food chain is one of the harmful properties of these heavy metals. They reach foods led by the air, household and industrial waste, animal dung and fertilizers [27]. The permanent exposure of the human being to lower measures of these heavy metals is reported to be co-factor of some neurological, carcinogenic and digestive diseases [28]. In addition, they represent the third source of food risk for human and animal after mycotoxins and microorganisms [26].
- For the consumer's health, the European Commission worked about a regulation laying down the maximum limits of heavy metals residues in fruits. From the resulted standard, the maximal values agreed are 0.05 mg/kg for cadmium and 0.5 mg/kg for mercury and lead [29]. However, international trades involve the systematic control of foodstuffs to insure their safety. In order to promote the distribution and consumption of foodstuffs, the analysis and control methods regarding toxic substances need to be effective and sound. The current study exhibits and validates a running method for the sure assessment of cadmium, lead, and mercury in kola nuts from Côte d'Ivoire.

#### 62 2. MATERIAL AND METHODS

#### 63 2.1 Sampling and pre-treatment

Sampling was achieved in accordance with the Regulation N° 333/2007of the European Commission [30]. Thus, 2 kg of kola nuts samples were collected in each district: Mountains, Comoe, Lagoons, and Down-Sassandra. The samples precisely derived from farmers, rural collectors, urban stores and big storage centres. Kola nuts were cut into small pieces with clean stainless knife. Then, they were dried at room temperature (30 ± 2 °C) for four weeks away from sun light ground in a hammer mill and kept in polyethylene sealed bags before their achievement at Central Laboratory for Food Hygiene and Agro-Industry (Abidjan, Côte d'Ivoire) to be analyzed.

- 71
- 72 2.2 Reagents

All reagents used in this study were of pure analytical grade, unless otherwise specified, were purchased from Merck, Germany : nitric acid 65%,hydrogen peroxide 35%, tin II chloride, steaming hydrochloric acid 37%, standards of cadmium, lead and mercury and ultrapure water at 18 MW.

76

#### 77 2.3 Apparatus and conditions of quantification of heavy metals

An Atomic Absorption Spectrophotometer (AAS, type VARIAN SPECTRAA 110) with GTA 110 furnace was used for the determination of cadmium and lead. Regarding the determination of mercury, the AAS was equipped with a VGA77 vaporization unit in the presence of a solution of 10% tin-II chloride previously prepared with 37% fuming hydrochloric acid. Nitrogen was used as a vector gas for the analysis. The operating conditions of the AAS device are shown in Table 1.

83

#### **Table 1. Operating Conditions of the Atomic Absorption Spectrophotometer**

	Cadmium	Lead	Mercury
Current intensity (mA)	4	10	4
Width of the slot (mm)	0.5	1	0.5
Wavelength (nm)	228.8	217	253.7
Coefficient of variation (%)	1	1	1
Integration time (seconds)	5	5	5
Number of repetitions	3	3	3

.

85

#### 86 **2.4 Validation of analytical method**

The validation of this analytical method for the determination of heavy metals (cadmium, lead, and mercury) was performed according to the French Standard (AFNOR, NFV03-110-1998) and the European directive 2001/22/EC [31, 32]. The process includes the study of the linearity for the calibration range, the determination of the limits of detection and quantification (LOD and LOQ values), the calculation of the relative standard deviation regarding repeatability and reproducibility assays, and the calculation of the recovery percentage for the analysis accuracy essays.

#### 94 2.4.1 Evaluation of the linearity

The adequacy of the calibration curve to the linear design was examined using 5 replications of a 5 independent points range. The linearity was assayed including the working range. The 5 calibration points were:

- 98 0.5 μg/L, 0.8 μg/L, 1 μg/L, 1.2 μg/L, 1.5 μg/L for cadmium
- 99 15 μg/L, 20 μg/L, 25 μg/L, 30 μg/L, 45 μg/L for lead;
- 100 10 µg/L, 25 µg/L, 30 µg/L, 50 µg/L, 100 µg/L for mercury. 101

#### 102 2.4.2 Limits of detection and quantification

The limits of detection (LOD) and quantification (LOQ) were calculated from the analysis of 10 separate assays of blank matrices. These parameters were measured using the following formulas:

#### LOD=Mx+3S

#### LOQ =Mx+ 10S

109 With: LOD:Limits of detection, LOQ: Limits of quantification, Mx: Average from 10 assays of blank 110 matrices, S: Standard deviation of blank values

## 111 **2.4.3 Assessment of the repeatability and reproducibility**

- 112 The repeatability of the analysis was probed with 10 assays of reference sample. For the 113 reproducibility, 5 separate assays were achieved with the reference sample at several days intervals.
- 114

106

107

108

#### 115 2.4.4 Analysis of the recovery

The extraction rate was determined from addition of various standards concentrations of heavy metals to uncontaminated solutions of kola nut samples. The concentrations of the added standard were 0.018 mg/kg, 0.116 mg/kg and 0.044 mg/kg for cadmium, lead and mercury, respectively. Ten

separate assays were achieved to assess the recovery rate allowed by the method of heavy metal

120 determination.

121

128

# 122 **2.5 Method of heavy metals mineralization**

An Aliquot of 0.5 g homogenate of each sample was heat mineralized with 7 ml of concentrated nitric acid (65%) and 1 ml of hydrogen peroxide (35%) using a microdigest for 20 min [33]. The mineralizate was reduced with a 10% chloride tin II solution previously prepared with 37% steaming hydrochloric acid for the mercury [34]. The resulted mineralised was then diluted in high quality ultrapure water and investigated with Atomic Absorption Spectrophotometry.

#### 129 2.6 Statistical analysis

130 The data were statistically treated using Statistical Program for Social Sciences (SPSS 20.0, SPSS for windows, USA) at 5% significance. Mean concentrations of cadmium, lead and mercury were 131 calculated; then the relative standard deviations were used as values of repeatability and 132 reproducibility. The square of Pearson correlation coefficient ( $R^2$ ) was calculated to appreciate the 133 134 linearity. The recovery rate was estimated to express the extraction yield. The mean concentrations 135 and the concentrations' variation range of the heavy metals allowed the description of contamination 136 range of the cola samples. The comparison of the concentrations recorded with the reference values 137 was performed using other 5% risk of conformity test.

138

# 139 3 RESULTS

#### 140 **3.1 Validation parameters for the quantification of the heavy metals**

- 141 The validation data deal with the values of linearity, repeatability, reproducibility, soundness, and limits 142 of detection and quantification involved from the heavy metals determination.
- 143 The results of the linearity analysis are recorded in Table 2. All the analytes exhibited good linearity 144 over the evaluated range with significant correlation coefficients ( $R^2 > 0.99$ ).
- Table 3 displays the statistical validity of the linearity over the full calibration range according to the statistical Fisher rule. Indeed, the F1 values calculated for regression (3743, 5758 and 6365 for Cd, Pb and Cd, respectively) are higher than the critical Fischer value (8.10). On the other hand, the F2 values are calculated for the error trend (2.97, 4.22, and 4.66 for Cd, Pb, and Hg, respectively) are lower than the critical Fisher value (4.94).
- Values of the limits of detection (LOD) and quantification (LOQ) of the heavy metals assessed are showed in Table 4. Values reported are 0.03 µg/L, 1.85 µg/L, and 2.92 µg/L for cadmium, lead, and mercury, respectively. Whereas the LOQ values are 0.07 µg/L (Cd), 6.52 µg/L (Pb) and 3.32 µg/L (Hg).

#### 155 Table 2. Calibration equationand determination coefficient from the heavy metal assessment

Heavy metal	Calibration equation <sup>a</sup>	Coefficient of determination (R <sup>2</sup> )
Cadmium (Cd)	y = 0.1836x + 0.0274	0.9978
Lead (Pb)	y = 0.0182x + 0.0949	0.9983
Mercury (Hg)	y = 0.0011x - 0.0021	0.9981

156 <sup>a</sup>: y - absorbance; x - concentration ( $\mu$ g/L).

# 157 Table 3: Linearity traits deriving from the experimental domain calibration

Heavy metals	F1	Critical F- value	Rating of the regression trend	F2	Critical F-value	Calibration domain
Cadmium (Cd)	3743		Acceptable	2.97		Acceptable
Lead (Pb)	5758	8.10	Acceptable	4.22	4.94	Acceptable
Mercury (Hg)	6365		Acceptable	4.66		Acceptable

<sup>158</sup> 

F1: F-value for the regression trend; F2: F-value for the statistical error trend

159

#### 160 Table 4: Minimal concentration (µg/L) for the detection and the quantification of the heavy

1	C	1
1	D	т

metals from the determination method used

Heavy metals	Limit of detection	Limit of Quantification
Cadmium (Cd)	0.03	0.07
Lead (Pb)	1.85	6.52
Mercury (Hg)	2.92	3.32

#### 162

163 The relative standard deviation (RSD) calculated from the repeatability assays are 3.32%, 2.11%, and 164 2.74% for cadmium, lead and mercury, respectively. Regarding the reproducibility, the RSD values are 165 3.98%, 3.28%, and 3.33% for respective cadmium, lead and mercury (Table 5).

The mean extraction yields resulting from the recovery of the measures added compared to the standard recovery of the heavy metals studied are 97.72% (Hg), 102.78% (Cd), and 104.31% (Pb) as showed in Table 6.

169

170

173

Table 5: Values measured (µg/L) and relative standard deviation (%) from the investigation of the repeatability and reproducibility during the determination of heavy metals assessed.

	Repeatability			Reproducibility		
Heavy metals	Standard solution (µg/L)	Value measured	RSD <sub>-value</sub>	Standard solution (µg/L)	Value measured	RSD <sub>-value</sub>
Cadmium	0.8	$0.78 \pm 0.2$	3.32	0.8	0.77 ± 0.03	3.98
Lead	30	$30.72 \pm 0.65$	2.11	10	10.5 ± 0.34	3.28
Mercury	15	14.94 ± 0.41	2.74	15	14.19 ± 0.47	3.33

<sup>174</sup> 

RSD-value: value of the relative standard deviation

175

# 176 Table 6. Data of the recovery traits for the heavy metals studied

Heavy metals	Standard recovery (mg/kg)	Recovery value measured (mg/kg)	Recovery rate (%)
Cadmium	0.018 ± 0.001	0.0185 ± 0.002	102.78 ± 3.7
Lead	0.116 ± 0.006	0.121 ± 0.009	104.31 ± 4.3
Mercury	$0.044 \pm 0.002$	$0.043 \pm 0.003$	97.72 ± 5.1

177 178

#### 3.1.2 Trends of heavy metals concentrations in cola samples

Table 7 shows the variation of the heavy metals concentrations of the cola samples from farmers andbig storage centres. Values are reported on the dry matter basis.

181 The mean concentrations of heavy metals from the farmers were 22.97  $\mu$ g/kg, 1065.57  $\mu$ g/kg, and 182 33.88  $\mu$ g/kg for cadmium, lead, and mercury, respectively.

From the big storage centres, the kola nuts record means of 24.99 μg/kg, 296.51 μg/kg and 39.74
 μg/kg of cadmium, lead, and mercury, respectively.

185 186

#### 187 Table 7 Mean kola nuts concentrations (µg/kg dry matter) in heavy metals

Heavy metals	Farmers		Bigstorage centres	
-	Mean±SD	[min-max]	Mean±SD	[min-max]
Cadmium	22.97±9.01	3.35-33.80	24.99±7.79	3.49-36.16
Lead	1065.57±613.76	465.49-2421.93	296.51±98.18	133.37-3270.08
Mercury	33.88±31.58	nd-81.35	39.74±34.66	nd-108.42

189 The cumulative mean concentrations of heavy metals for the overall samples studied are displayed in 190 fig. 1.The highest content was found from the farmers (1122.42 µg/kg); whereas the big storage 191 centers showed the lowest heavy metals cumulative content (660.75 µg/kg).



193



Fig. 1. Cumulative mean concentrations of three heavy metals from the kola nuts studied.

196

194

195

#### 197 3.2 Discussion

198 The linearity of every determination method is the ability, within a range of digits, to provide 199 information values or results which are proportional to the amount of the analyte to be measured from 200 the material studied [35]. The assays performed for the linearity highlighted the normality of the distribution across the calibration range set at 0.5 µg/L to 1.5 µg/L, 15 µg/L to 45 µg/L and 10 µg/L to 201 202 100 µg/L for cadmium, lead and mercury, respectively. According to the current linearity data, the method for determining heavy metals (Cd, Pb and Hg) could be considered as a reliable process for 203 204 kola samples.

From the statistical Fischer test, the results showed that the variance proportion due to the error of the 205 206 experimental design is not higher than the variance of the experimental error. For each heavy metal measured, the calculated F-value is lower than the critical F-value corresponding to a Fisher variable at 207 1% statistical significance. The results obtained show that the linearity domain is valid and the 208 209 regression design is also acceptable. In addition, overall Pearson determination coefficients(R<sup>2</sup>) recorded during the study were closed to 1. The determined second-order mathematical models are 210 211 therefore valuable for forecasting the main responses [36].

212 The limits of detection (LOD) and quantification (LOQ) are similar to the values reported by Labat et 213 al.[37].These authors showed LOD values reaching 0.02 µg/L for the cadmium and 1.3 µg/L for the 214 lead; whereas the LOQ values were respectively recorded at 0.03 µg/L and 1.4 µg/L. However, the 215 slight variations of LOD and LOQ values could be explained by the soundness of the apparatus. Indeed, these authors have used the inductively coupled plasma added with a Mass Spectrometry 216 217 (ICP-MS). According to Alsac [38], some chemical elements result in higher LOQ values in ICP-MS 218 compared to the ICP-AES used in the current study.

Regarding the accuracy of the determination method as estimated by repeatability and reproducibility 219 220 essays, overall relative standard deviations are lower, below 5%. The RSD-values were ranged from 221 2.11% to 3.32% for the repeatability and from 3.28% to 3.98% for the reproducibility. This observation is in the same trend as the work of El Alami [39] stating that the lower experimental error involved by 222

- standard deviation reflects the closeness between the values obtained from various measurements of 223 224 the same object under specified conditions. Both reproducibility and repeatability analyses performed
- 225 using our experimental design are sounded.
- 226 For the recovery assessment, the results showed recovery rates between 97.72% and 102.78% from the reference sample. In addition, there wasn't any significant difference with the evaluation of the 227 228 conformity.
- 229 The data of this study are in accordance with the operating conditions recommended by the FAO [40]
- 230 as acceptable analysis technique for heavy metals determination, since the results highlighted 231 reliability and good precision of the mineralisation and analysis operations.

232 Thereafter, the kola nuts samples recorded significant contents in heavy metals (Cd, Pb, and Hg). The 233 results evidenced various contents according to the origin of the kola sample and the heavy metal 234 assessed. From overall cumulated heavy metals concentrations, the samples originating from the farmers recorded the highest value. The high concentrations of heavy metal in kola nuts can be 235 attributed to the cultivation techniques used by farmers. However, the excessive use of chemical 236 237 fertilizers is a source of contamination of agricultural soils and kola nuts [7]. Also, during bush fires, 238 heavy metals are emitted into the environment as particles during combustion and contaminate kola 239 nuts [41]. According to Dauguet [26], the changes of heavy metals concentrations in food stuffs 240 derives from natural atmospheric conditions (volcanism, dust of erosion), anthropogenic activities 241 (industry, transport), and human contributions (fertilizers, phytosanitary products, animal dung, urban 242 sludge, etc.). 243

## 244 4. CONCLUSION

The study showed that the method suggested for the heavy metals determination from kola nuts is really suitable. The assays resulted in linear calibration curve within the heavy metals concentration range considered. The determination method is reproducible and repeatable, and is therefore trusted. It is easily implementable in every laboratory equipped with Atomic Absorption Spectrophotometer. It's also sensitive and does reveal any matrix effect (good specificity). Thus, this method could help in ensuring effective sanitary control at different critical points of kola nut distribution channel for promoting a good management of the toxicity concerns in such products.

#### 252 253 COMPETING INTERESTS

- 254 Authors have declared that no competing interests exist.
- 255 256

# 257 **REFERENCES**

# 258

- 259 1. Opeke L.Tropical commodity tree crops, Spectrum Books Limited, Ibadan. Nigeria. 2005;86.
- 260 2. Famaye A. Handbook on Kola Production. PAMMA PRESS. Iloro Street, Off Stadium Road, Akure,261 Ondo State, Nigeria. 2012;12.
- 3. Aikpokpodion E, Oduwole O, Adebiyi S. Appraisal of Pesticide Residues in Kola Nuts Obtained from
   Selected Markets in Southwestern, Nigeria. J Sci Res Rep. 2013;2(2):582-597.
- 4. Deigna M, Kouadio K, Konan N, Biego G. Diagnosis in Production and Post-harvest Processing of
   Nuts of *Cola nitida* (Malvaceae) in Côte d'Ivoire. J Agric Ecol Res Int. 2016;9(2):1-11.
- 5. Seri Z. Projet de plan d'actions de l'APPEXCO-CI 2013-2015. Centre d'Etudes et de Recherches
   Appliquées (CERA). 2012;50. French.
- Asogwa E, Oluyole K, Ndubuaku T, Uwagboe E. Kolanut production, processing and marketting in
   the south eastern states of Nigeria. A-E J Env Sci. 2012; 12:463-468.
- 7. Adebiyi S, Uwagboe O, Agbongiarhuoyi E, Ndagi I, Aigbekaen E. Assessment of Agronomic
   Practices among Kola Farmers in Osun State, Nigeria. W J Agric Sci. 2011;7(4):400-403.
- Adeosun S, Adejobi K, Famaye A, Idrisu M, Ugioro O, Nduka B. Combined effect of kola testa
   based organic manure and NPK fertilizer on soil, leaf chemical composition and growth performance
   of kola (Cola nitida). Res J Agr Env Man. 2013;2(7):183-189.
- 9. Trueby P.Impact of Heavy Metals On Forest Trees From Mining Areas. In International Conference
   On Mining And The Environment III, Sudbury, Ontario, Canada; 2003.
- 277 Available: www.x-cd.com/sudbury03/prof156.html.
- 10. Ndubuaku T, Asogwa E, Hassan A. Distribution of kolanut weevil (*Balanogastris kolae*)
  (Coleoptera: Curculionidae) in *Cola nitida* stored in baskets. Afr j plant sci. 2015;9(1):13-16.
- 11. Adebayo S, Oladele O. Medicinal Values of Kolanut in Nigeria: Implication for Extension Service
   D.L Sci J. 2012;9(2):887-891.
- Agbeniyi S. Efficacy of Milton solution and wood ash in the control of storage rot of kolanut (*Cola nitida*) (Vent.) Schott and Endl. Nigerian J Tr Crop Res. 1999;3:67-75.
- 13. Biego G, Yao K, Ezoua P, Chatigre K, Kouadio L. Niveaux de contamination en pesticides organochlorés des noix de *Cola nitida*. Int J Biol and Chem Sci. 2009;3(6):1238-1245. French.
- 14. Al-Jassir M, Shaker A, Khaliq M. Deposition of heavy metals on green leafy vegetables sold on
   roadsides of Riyadh city, Saudi Arabia. Bul Env Cont Toxicol. 2005;75:1020-1027.
- 15. Nordström M. Concentrations of Organochlorines Related to Titers to Epstein-Barr Virus Early Antigen IgG as Risk Factors for Hairy Cell Leukemia. Environnemental Health Perspective.
- 290 2000;108:441-445.

- 16. Vine M. Plasma 1,1 Dichloro-2,2- bis (p-chlorophenyl) ethylene (DDE) in Levels and Immune
  Response. A J Epidemiol. 2001; 221:53-63.
- 17. Tindall R. The culture of Cola: social and economic aspects of a West African domesticate.
  Ethnobotanical Leaflets. 1998;2:1-3.
- 18. Ojo G, Nwoha P, Ofusori D, Ajayi S, Odukoya S, Ukwenya V. Microanatomical effects of ethanolic extract of *Cola nitida* on the stomach mucosa of adult wistar rats. Afr J Trad Compl Alter Med.
- 297 2009;7:47-52.
- 19. Ajai I, Ochigbo S, Jacob O, Ndamitso M, Abubakar U. Proximate and mineral compositions of different species of kola nuts. Euro J Appl Eng Sci Res. 2012;1:44-47.
- 300 20. Odebode A. Phenolic compounds in the kola nut (*Cola nitida* and *Cola aculinata*)
- 301 (Sterculiaceae) in Africa. Revista de Biologia Tropical. 1996;44(2):513-515.
- 302 21. Onomo P, Niemenak N, Ndoumou O., Lieberei R. Change in amino acids content during 303 germination and seedling growth of Cola sp. Afr J Biotechnol. 2010;9(35):5632 5642.
- 22. Niemenak N, Effa O, Lieberei R, Omokolo N. Purine alkaloids and phenolic compounds in three Cola species and *Garcinia kola* grown in Cameroon. S Afr J Bot. 2008;74:629-638.
- 306 23. Nyamien Y, Adje F, Niamke F, Chatigre O, Adima A., Biego G. Caffeine and phenolic compounds
  307 in *Cola nitida* (vent) Schodt and Endl and Garcinia kola Heckel grown in Côte d'Ivoire. Brit J Appl Sci
  308 Technol. 2014;4 (35):4846-4859.
- 24. Dongo N, Manjula K, Orisajo B. Occurrence of ochratoxin A in Nigerian kola nuts. Afr Cr Sci Conf
   Proc. 2007;8:2133-2135.
- 25. Salama A, Radwan M. Heavy Metals (Cd, Pb) and Trace Elements (Cu, Zn) Contents in Some
   Foodstuffs from Egyptian Market. E J Agr Sci. 2005;17(1):34-42.
- 26. Dauguet S, Denaix L, Nguyen C, Royer E, Levasseur P, Potin G. Mesure des flux d'éléments
  traces (Pb, Cd, As, Cu, Zn) dans les sols, végétaux, porcs et lisiers des exploitations porcines du Sud
  -ouest. Innovations Agronomiques. 2011;17:175-190. French.
- 27. Verloo M. Les métaux lourds dans les denrées alimentaires: origine et évolution des teneurs.
  Symposium "Les oligo-éléments dans l'alimentation en Belgique Données récentes" Institut
  Danone.2003;3. French.
- 319 28. Achour S, Iken I, Amarti A. L'intoxication au plomb: caractéristiques toxicologiques et clinico biologiques au Maroc. 2014;21:9-12. French.
- 321 29. Règlement (CE) N°420/2011.de la Commissiondu 29 avril 2011 modifiant le règlement (CE)
- N°1881/2006 portant fixation de teneurs maximales pour certains contaminants dans les denrées
   alimentaires. J O UE. 2011;4. French.
- 324 30. Règlement (CE) N° 333/2007 de la Commission du 28 mars 2007 portant fixation des modes de 325 prélèvement d'échantillons et des méthodes d'analyse pour le contrôle officiel des teneurs en plomb, 326 en cadmium, en mercure, en étain inorganique, en 3-MCPD et en benzo(a)pyrène dans les denrées
- alimentaires. J O UE. 2007;10. French.
- 31. Association Francaise de Normalisation. Analyse des produits agricoles et alimentaires: Procédure
   de validation intra-laboratoire d'une méthode alternative par rapport à une méthode de référence.
   Paris : Edition AFNOR. 1998;40. French.
- 331 32. Directive 2001/22/CE: mode de prélèvement d'échantillons et de méthode d'analyse pour le
  332 contrôle officiel des teneurs en mercure, plomb, cadmium et 3-MCPD dans les denrées alimentaires.
  333 Journal officiel. 2001;77:14-21. French.
- 33. AOAC official method 999.10. Determination of Lead, Cadmium, Zinc, Copper, and Iron in Foods
   by Atomic Absorption Spectrometry after Microwave Digestion. J. AOAC int. 2005, 83:1189-1203.
- 336 34. AOAC official method method 974.14. Mercury in fish by flame atomic absorption spectroscopy. J.337 AOAC int. 1976.
- 338 35. OIV: Organisation Internationale de la Vigne et du Vin: Recueil des méthodes internationales
  339 d'analyse des vins et des mouts. Résolutions adoptées à Paris (France) 3ème A.G. 17 juin 2005.
  340 2006;2:416. French.
- 341 36. Chan S, Lee C, Yap C, Wan A, Ho C. Optimization of extraction conditions for phenolic 342 compounds from limau purut (*Citrus hystrix*) peels. Int Food Res J. 2009;16:203-213.
- 343 37. Labat L, Dehon B, Dhorne C, Lhermitte M. Dosage de métaux par ICP-MS dans différents milieux 344 biologiques. Annales de Toxicologie Analytique. 2003;4 (15):281-286. French.
- 38. Alsac N. Dosage des métaux lourds (As, Cd, Cr, Cu, Ni, Pb, Zn et Hg) dans les sols par ICP-MS.
  Annales de Toxicologie Analytique. 2007; 1 (19):41p. French.
- 347 39. El Alami S. Validation de dosage des ions fluorures (dosage par électrodes spécifiques), 348 Université Sidi Mohamed Ben Abdellah Fès, Maroc. Projet de fin d'études.2012; 34. French.
- 40. FAO: Food and Agriculture Organization Validation of analytical methods for food control. Vienna:
- 350 Report of joint FAO/IAEA Expert Consultation. 1997;11:18.

- 41. Yahaya M., Ezeh C., Musa Y. & Mohammad S. Analysis of heavy metals concentration in road sides soil in Yauri, Nigeria. Afr J P Appl Chem. 2010, 4(3), pp 22-30.