

## Response of BARI Bush Bean-2 at different levels of nitrogen fertilization

### ABSTRACT

A field experiment was carried out at Sher-e-Bangla Agricultural University Farm during the *Rabi* season of 2008 to investigate the response of BARI Bush Bean-2 at different levels of nitrogen fertilization in respect of various yield components. The soil of the experimental area belongs to the Modhupur Tract (AEZ No. 28) which had shallow red brown terrace soil. The selected plot was medium highland and the soil series was Tejgaon. The experiment was laid out in a RCBD with three replications. The experiment comprises 4 levels of nitrogen in the form urea (0, 40, 80 and 120 kg ha<sup>-1</sup> N). The results obtained revealed that different levels of nitrogen showed significant variations on the different parameters studied. Among the different treatments of the study the treatment N<sub>80</sub> (80 kg N) gave the highest pod length (11.15 cm), pod diameter (2.93 cm), average single pod weight (5.42 g), total pod weight plant<sup>-1</sup> (25.63 g), pod yield plot<sup>-1</sup> (1789.58 g) and pod yield hectare<sup>-1</sup> (4.37 ton). Thus the findings of the experiment revealed that the use of 80 kg N maximally influence the yield and yield contributing characters of BARI Bush Bean-2 in the selected red brown terrace soil of the Tejgaon series.

**Keywords:** Yield; Bush Bean; Nitrogen

### 1. Introduction

Being a major grain legume crop of the world Bush bean (*Phaseolus vulgaris* L.) is consumed in all parts of the globe for its edible seeds and pods [1]. It belongs to the family Leguminosae, and sub-family Papilionaceae [2]. In Bangladesh, this crop is known as “Farashi Sheem” [3]. It is also known as french bean, kidney bean, common bean, basic bean, snap bean, raj bean, navy bean, haricot bean, string bean, pole bean, wax bean and bonchi [4]. Bush bean originated in Central and South America [5]. Now it is cultivated in many parts of the tropics, sub-tropics and throughout the temperate region [6]. It is an annual herbaceous vegetable crop-producing straight or slightly curved pod. Bush beans are green, tender, rounded shaped pods, 4 to 6 inches (10 to 15 cm) long. The pods are eaten whole, including the immature seeds, when they are still young, juicy and tender. Its green pods are rich in protein, carbohydrate, fat, thiamin, riboflavin, Ca, Fe and niacin as well as fiber [7,8]. It also contains various immune system-boosting antioxidants e.g. flavonoids and carotenoids [9]. Due to its high levels of flavonoids, this power-packed legume has been shown to help manage and regulate diabetes symptoms in many patients and reduce the risk of heart diseases [10]. Bush bean has a delicate texture and sweet flavor. Fresh green beans are sweeter than canned. This crop is also valuable in Bangladesh mostly for exporting tender beans to the European market [11]. In recent years, the Hortex Foundation, Vegetable Export Center of BRAC and some other organizations facilitated the export of this vegetable of Bangladesh. Although, to play an important role in the economy of Bangladesh and increasing popularity in vegetable consumption, the yield of Bush bean in Bangladesh is lower than the world average. In this context, the yield of Bush bean has become an important component of vegetable production and cash income. Many factors influenced the production of BARI Bush Bean-2. Among them, fertilizer especially nitrogen is the most effective and critical input for increasing crop yield [12]. In agro-ecological zones the requirement of fertilizer for any crop varies with the cultivars and soil types [13]. Nitrogen is one of the most limiting nutrients to plant growth [14]. [15] Found a maximum pod yield of Bush bean from 120 kg N ha<sup>-1</sup>. By using 120 kg N ha<sup>-1</sup>, [16] found the maximum plant height, the number of branches plant<sup>-1</sup> and green pod yield<sup>-1</sup> of Bush bean. [2] Stated that the maximum plant height (39.03 cm), number of branches plant<sup>-1</sup> (18.25), pod length (14.10 cm), pod weight (5.37 g) and pod yield (8.26 t ha<sup>-1</sup>) were recorded in the treatment receiving 120 kg N. The short growth cycle of this crop is one of the reasons that Nitrogen

fixation is not as efficient as other legumes because carbohydrate production by the Bush bean plant occurs at the same time as the plants and the rhizobium needs a maximum carbohydrate supply and there is a heavy competition for carbohydrate between the bacteria and Bush bean [17]. Nitrogen is the most essential elements for crop production. It encourages vegetative growth and increases the leaf area of plants, which helps in photosynthetic activity. It stimulates the root growth and development of the plant. Furthermore, it helps in the uptake of other nutrients from the soil. The vegetative growth and yield of BARI Bush Bean-2 were increased with a successive increase in the dose of nitrogen [18]. BARI Bush Bean-2 can fix atmospheric nitrogen in its root zone. An optimum amount of nitrogen for the whole growth period is necessary to produce the maximum yield of good quality zharshemor BARI Bush Bean-2 [19]. For that reason, nitrogen fertilizer should be applied in an optimum dose to maximize its utilization for crop production. Therefore the present experiment carried out to investigate the response of BARI Bush Bean-2 at different levels of nitrogen fertilization in respect of various yield components.

## 2. Methods and materials

The research work was conducted in *Rabi* season at Sher-e-Bangla Agricultural University Farm, Sher-e-Bangla Nagar, and Dhaka during the *Rabi* season of November 2008 to March 2009. It is located at 90.335°E longitude and 23.774° latitudes. The treatment of the experiment was different doses of Nitrogen viz. 0, 40, 80, 120 kg ha<sup>-1</sup>. The experiment was laid out in Randomized Complete Block Design (RCBD) with 3 replications. The unit plot size was 2.4 m x 1.5 m (3.6 m<sup>2</sup>). The distance between the two plots was 50 cm and between blocks was 50 cm. The plots selected for the experiment was plowed on the 30<sup>th</sup> of November 2008 and cross-plowed several times to obtain a good tilth. The manures and fertilizers like Cow dung, TSP (Triple Super Phosphate), MoP (Muriate of Potash), Gypsum, and Boric acid were applied at the rate of 5000, 2000, 60 and 5 kg ha<sup>-1</sup>, respectively [20]. The entire quantity of Cow dung, Gypsum, and Boric acid were applied during the final land preparation while full doses of TSP and half of MoP were at seed sowing time and rest half MoP 30 days after seed sowing. Seeds of BARI Bush Bean-2 were sown on 03 December 2008. The seeds were sown per hill at a depth of 5 cm and the seeds were covered with pulverized soil just after sowing and gently pressed with hands. Intercultural operations were done whenever required for getting better growth and development of the plants. Data were collected on pod length, pod diameter, average single pod weight, total pod weight plant<sup>-1</sup>, pod yield plot<sup>-1</sup> and pod yield hectare<sup>-1</sup>. The data obtained from the experiment were analyzed statistically using the MSTAT computer package program to find out the significance of the difference among the treatments. The mean values of all the treatment were calculated and analysis of variances for all the characters was performed by the 'F' (variance ratio) test. The significance of the differences among the pairs of treatment means was estimated by the Duncan Multiple Range Test (DMRT) at 1 and 5% level of probability [21] for the interpretation of results.

## 3. Results and Discussion

### 3.1 Pod Length

Nitrogen influenced significantly in respect of pod length. Pod length increased gradually due to increasing dose of nitrogen fertilizer. The length of pod increased significantly with the increasing levels of nitrogen up to 80 kg N ha<sup>-1</sup>. The highest length of the green pod was found (11.15 cm) in treatment N<sub>80</sub> which is statistically identical with N<sub>40</sub>, N<sub>120</sub> treatments and the lowest (10.30 cm) was found in the plot treated with N<sub>0</sub> treatment (Table 1). This result is in full agreement with [22, 23]. They reported that pod length was significantly influenced by optimum dose of N. Nitrogen is a part of the chlorophyll molecule which activates the normal metabolic process of leaves. The leaves become metabolically most active those contain the highest level of N. So the addition of nitrogen enhances the vegetative growth viz. pod length of BARI Bush Bean-2.

### 3.2 Pod diameter

At different levels of nitrogen application there were found significant variations in pod diameter. By the increasing levels of nitrogen the diameter of pod increased significantly up to 80 kg N ha<sup>-1</sup>. The topmost diameter of the pod was found (2.93 cm) in treatment N<sub>80</sub> which is statistically alike with N<sub>120</sub> treatment

and the lower most (2.73 cm) was found in the plot treated with  $N_0$  treatment (Table 1). These results are in alike with the findings of [15, 24]. An adequate supply of nitrogen favors the formation of carbohydrates as it is a part of the chlorophyll molecule. The ideal level of N promotes the metabolic process and produces more food that reserved in fruits. That's why pod diameter increases with the application of an optimum dose of N.

### 3.3 Average single pod weight

A significant influence was observed in respect of average single pod weight at different rates of nitrogen. The highest average single pod weight was found (5.42 g) in treatment  $N_{80}$  which is statistically similar with  $N_{40}$ ,  $N_{120}$  treatments while the lowest (4.78 g) was found in the control plot (Table 1). It was clearly observed that there was a positive effect between optimum level of nitrogen and the pod weight plant. [25] Also reported similar results. The increment in the yield component may be due to the proper use of nitrogenous fertilizer which resulted in boosting in biosynthesis of the photosynthates and finally the yield.

### 3.4 Total pod weight plant<sup>-1</sup>

Statistically, a significant variation was found for different levels of nitrogen with total pod weight plant<sup>-1</sup>. The highest total pod weight per plant was observed (25.63 g) in treatment  $N_{80}$  which is statistically identical with  $N_{40}$ ,  $N_{120}$  treatments and the lowest (22.85 g) was found in the zero treated plot (Table 1). This result is similar to that of [26]. About 50% at higher doses of applied N remains unavailable to a crop due to N loss through leaching. So, optimum dose of N gave the best result.

### 3.5 Pod yield plot<sup>-1</sup>

The effect of nitrogen on pod yield plot<sup>-1</sup> showed a statistically significant difference among the treatments at the harvesting stage of BARI Bush Bean-2. Pod yield plot<sup>-1</sup> increased significantly by the increasing rates of nitrogen up to 80 kg N ha<sup>-1</sup>. The maximum pod yield plot<sup>-1</sup> was found (1789.58 g) in treatment  $N_{80}$  which is statistically alike with  $N_{40}$  treatment followed by  $N_{120}$ . The lowest on pod yield plot<sup>-1</sup> (1454.17 g) was found in control treatment (Table 1). Similar result was found by [22, 27]. On the other hand [21] found significantly influenced by optimum dose of nitrogen on green pod yield. Nitrogen is an integral part of proteins, phytochromes, compounds, coenzymes, chlorophyll and nucleic acids. All the biochemical processes occurring in plants are mainly governed by nitrogen and its associated compounds which make it essential for the growth and development of BARI Bush Bean-2.

### 3.6 Pod yield hectare<sup>-1</sup>

Pod yield hectare<sup>-1</sup> was recorded at the harvest stage of BARI Bush Bean-2 and it was observed that there were significant variations in relation to nitrogen application. Pod yield hectare<sup>-1</sup> increased significantly by the increasing levels of nitrogen up to a certain dose. The highest pod yield hectare<sup>-1</sup> was found (4.37 ton) in treatment  $N_{80}$  which is statistically identical with  $N_{40}$  treatment followed by  $N_{120}$ . The lowest (3.55 tons) was found in the control treated plot (Table 1). [22] Found the similar result and stated that pod yield of bush bean increase with the increment of nitrogenous fertilizer. Among all the essential nutrients applied to the plants nitrogen is the major one which has a key role in the process of photosynthesis. Increased rate of photosynthesis by the best dose of nitrogen gave more yield because large amount of dry matter, more assimilates were produced and transported to fill the seeds as a result of more applied nitrogen.

## Conclusion

This experiment was conducted to investigate the response of BARI Bush Bean-2 at different levels of nitrogen fertilization in respect of various yield components. Though the different treatments of the experiment ( $N_{80}$  and  $N_{120}$  kg ha<sup>-1</sup>) performed similarly but 80 kg ha<sup>-1</sup> N performed best in all the studied characters. As for maximum output, an over-fertilized condition of nitrogen did not bring out any significant advantage so excessive use of N fertilizer should avoid. So this dose of nitrogen fertilizer can be recommended for commercial cultivation of BARI Bush Bean-2.

## REFERENCES

1. Heuzé V, Tran G, Nozière P, Lebas F. Common bean (*Phaseolus vulgaris*). Feedipedia, A programme by INRA, CIRAD, AFZ and FAO. 2015;14:50.
2. Ali QS, Zeb S, Jamil E, Ahmed N, Sajid M, Siddique S, Shahid M. Effect of various levels of nitrogen, phosphorus and potash on the yield of French Bean. Pure and Applied Biology. 2015; 4(3):318.
3. Rashid MM. SabjiBiggan (In Bengli). 1st ed. Bangla Academy, Dhaka, Bangladesh. 1993;387-390.
4. Tindall HD. Vegetables in the Tropics. McMillan Education Ltd., New York. 1988;527.
5. Swiader JM, Ware GM, McCollum JP. Producing Vegetable Crops. 4<sup>th</sup> ed. Interstate Publishers, Inc., Danville, Illions, USA. 1992;233-249.
6. Purselove JW. Tropical Crops: Dicotyledons. Longman. New York. 1992;52.
7. Pierce LC. Legumes In: Vegetable: Characters, Production and Marketing. John Wiley and Sons, New York. 1987;561-567.
8. Rashid MM. SabjiBiggan. In Bangla. 2nd Ed., Rashid Publishing House, Dhaka. 1999;396 -399.
9. Fatema R, Rahman J, Shozib H, Nazrul M, Fatima K. Genetic Diversity and Nutritional Components Evaluation of Bangladeshi Germplasms of Kidney Bean (*Phaseolus vulgaris* L.). Journal of Genetic Resources, 2019;5(2):83-96.
10. Baloch MS, Zubair M. Effect of nipping on growth and yield of chickpea. J Anim Pl Sci., 2010; 20(3):208-210.
11. Kakon SS, Bhuiya M SU, Hossain SMA, Sultana N. Flowering Behaviour and Seed Yield of French Bean as Affected by Variety. International Journal of Applied Sciences and Biotechnology, 2015;3(3):483-489.
12. Arya MPS, Kalara GS. Effect of phosphorus dose on the growth, yield and quality of summer mung (*Vignaradiata* L. Wilezek) and soil nitrogen. India J. Agric. Res., 1988;22(1):23-30.
13. Mitra SK, Sadhu ML, Bose TK. Nutrition of Vegetable Crops. Nayapokash, Calcutta. 1990;157-159.
14. Hirel BJ, Gouis B, Ney A, Gallais F. The challenge of improving nitrogen use efficiency in crop plants: towards a more central role for genetic variability and quantitative genetics within integrated approaches. J. Exp. Bot., 2007;58(9):2369-2387.
15. Prajapoti MP, Patel HA, Prajapati BA, Patel LR. Studies of nutrient uptake and yield of French bean (*Phaseolus vulgaris* L.) as affected by weed control methods and nitrogen levels. Legume Res., 2004;27(2):99-102.
16. Sharma SK. French bean green pod and seed production as influenced by nitrogen and phosphorus application. Annals of Agricultural Research. 2001;22(1):130-132.

17. Van Schoonhooen A, Voysests O. Common Bean: Research for Crop Improvement. CAB International, Oxon. 1991;980.
18. Tewari SP, Singh RP. Effect of rate and time of application of nitrogen on growth and productivity of French bean (*Phaseolus vulgaris* L.). Legume Res., 2000;23(2):110-113.
19. Habbish IG, Ishaq MK. Nodulation of legume on Sudan.III. Response of haricot bean to inoculation. Expl.Agric., 1974;10:45-50.
20. BARI (Bangladesh Agriculture Research Institute)..AdhunikKrishiProjukti Hat Boi. 2019; 146.
21. Gomez AK, Gomez AA. Statistical Procedures for Agricultural Research. Second Edition. John Wiley and Sons, New York, U.S.A.1984.
22. Nasrin S, Al-Amin M, Mabud AG, Sani MNH. Growth and Yield Response of Bush Bean (*Phaseolus vulgaris* L.) as Influenced by Different Levels of Nitrogen and Phosphorus Application. Asian Journal of Research in Crop Science, 2019;1-8.
23. Rahman A. Influence of nitrogen and plant spacing on French bean. A thesis of Horticulture Department. Bangladesh Agricultural University, Mymensingh, 2001.
24. Begum A, Ahad A, Kaiser MO, Islam MM, Anam MK. Effect of sowing dates and fertilizer treatments on the reproductive variability of French bean (*Phaseolus vulgaris*) Pak. J. Biol. Sci., 2003;6(22):1897-1901.
25. Sharma SR, Prakash O, Ahlawat IPS. (Response of French bean (*Phaseolus vulgaris* L.) varieties to plant density and nitrogen application. Indian J Agron., 2008;46(2):277-281.
26. Sharma SK, Sing R, Singh ON. Influence of nitrogen levels and Rhizobium inoculation on yield, quality and nitrogen uptake of French bean (*Phaseolus vulgaris* L.). Res. Crops., 1996;3(3):524-528.
27. Chandel RS, Singh R, Singh RS, Singh ON. Influence of nitrogen levels and Rhizobium inoculation on yield, quality and nitrogen uptake of French bean (*Phaseolus vulgaris* L.). Research on Crops. 2002;3(3):524-528.

UNDER PEER REVIEW

**Table 1. Effect of nitrogen on the growth and yield of BARI Bush Bean-2**

Treatment	Pod length (cm)	Pod diameter (cm)	Average single pod weight (g)	Total pod weight per plant (g)	Pod yield per plot (g)	Pod yield per hectare (ton)
N <sub>0</sub>	10.30 b	2.73 c	4.78 b	22.85 c	1454.17 c	3.55 c
N <sub>40</sub>	10.55 ab	2.74 b	5.17 ab	24.42 ab	1600.00 ab	3.90 ab
N <sub>80</sub>	11.15 a	2.93 a	5.42 a	25.63 a	1789.58 a	4.37 a
N <sub>120</sub>	10.90 ab	2.90 ab	5.00 ab	24.99 ab	1537.50 b	3.750 b
CV (%)	7.28	7.47	10.48	7.01	15.43	15.41
Level of significance	*	*	*	*	*	*