

Original Research Article

Qualitative Risk Assessment of *Campylobacter jejuni* in Street Vended Poultry in Informal Settlements of Nairobi County

ABSTRACT

Aim: To determine the food safety risks of consumption of street-vended poultry products, to evaluate the determinants of microbial safety and the risk rank of these products.

Study design: A cross-sectional survey was done in the Korogocho and Kariobangi North slums among the consumers and vendors to assess their food safety knowledge and practices. Swab samples of the cooking equipment, utensils, and personnel, raw and cooked portions of poultry were collected for microbial quality evaluation. The most prevalent microorganism was assessed for its qualitative risk rank using the Risk Ranger software.

Place and duration of study: The study was carried out in the capital city of Kenya, Nairobi, from June 2018 to July 2018.

Methodology: A total of 15 vendors were exhaustively sampled and included in the study with the food safety and hygiene practices evaluated using a food safety checklist. The snowballing sampling technique was used to locate all the vendors. Samples of raw and cooked street vended poultry products were subjected to microbial analysis. All samples were collected in sterile polythene bags followed by transportation to the laboratory of the Department of Food Science and Technology of the University of Nairobi and microbial analysis.

Results: *Campylobacter jejuni* contamination, in both raw and cooked poultry products, was $8.95 \pm 0.94 \log_{10}$ CFU g⁻¹ and $4.66 \pm 2.67 \log_{10}$ CFU g⁻¹ respectively; the probability of contamination of raw street-vended poultry was found to be 48.96 %. The mean weekly intake of the poultry was reported 140.0 g per person. The probability of campylobacter infection in an individual consumer was found as 7.12×10^{-3} with the predicted illnesses among the population found as 1.11×10^6 cases. The qualitative risk estimate from the study was reported as 67, above the limit of 48 for medium risk.

Conclusion: The study concluded that *Campylobacter jejuni* posed high food safety risks as a resultant from the consumption of street-vended poultry.

Keywords: Campylobacter jejuni; Qualitative; Risk assessment; Street-vended; Poultry; Informal settlements

1. INTRODUCTION

Informal food vending either along the streets or market places contributes to the daily food intake of the urban and peri-urban poor [1]. The utilization of street foods is so high in developing countries, especially in informal settlements [2]. The increasing popularity of street vended poultry is coupled with the increased intake of ready to eat poultry, especially in urban areas [3]. This is because street-vended poultry is known to be affordable, accessible and ready-to-eat, thereby the high utilization in the urban households. Street-vending in sub-Saharan African countries and other developing countries is informal with less regulation of the sector [4], thus the value and quality of food in this sector are usually not easily documented.

Despite the ever-increasing utilization of street-vended poultry products, safety concerns still linger over these foods. These foods are often sold in compromised hygienic conditions and are usually left open for display [5]. Microbial pathogens including but not limited to *Salmonella*, *Escherichia coli*, *Listeria monocytogenes*, *Staphylococcus aureus* and *Campylobacter jejuni* are prevalent in these products [6–9]. A study done on street-vended chicken in Taiwan and the Philippines reported a salmonella contamination level of 7% and 8%, respectively [6]. Considering that poultry is a vehicle for various microorganisms, its contamination with disease-causing microorganisms such as *Salmonella* and *Campylobacter* becomes a public health concern [8]. Factors that predispose these foods to contamination include personal and equipment hygiene, working environment and time and temperature abuse for the cooked food [10].

Campylobacter species including *C. jejuni* and *C. coli* are potent food pathogens found in poultry products and indeed several risk assessment reports have attributed chicken consumption to several campylobacter incidents [3, 11]. The two microorganisms are part of a group of *Campylobacter* species that causes campylobacteriosis, a major foodborne infection from poultry products [12]. Campylobacteriosis is highly infectious and a leading cause of bacterial gastroenteritis [13]. The EU notification rate for campylobacteriosis in 2012 was found to be 55.49 per 100,000 populations [14]. *C. jejuni* and *C. coli* account for almost 95% of the global incidence of campylobacteriosis [15]. Additionally, *C. jejuni* infection can also lead to auto-immune conditions of Guillain-Barré syndrome (GBS) and Miller Fisher syndrome [12].

The risk posed by the contamination of *C. jejuni* in street vended poultry needs to be managed through guided policy action with a scientific justification of its course. However,

quantitative risk assessment of *C. jejuni* in street-vended chicken in Nairobi County was difficult due to limited specific data on the food-borne illnesses associated with this microorganism in the country [16]. In as much as the qualitative risk assessment obtains a categorical or descriptive risk estimate rather than a numerical value as in the case of quantitative, the outcome is still valid for guided action [17]. In Europe, risk assessment has become a norm due to availability of tools, reliable data, and experts; however, the converse is true in Kenya; making the present study to be of such a priority. This study used the qualitative risk assessment technique to rank the risk resulting from the contamination of the street vended poultry with *C. jejuni* in the informal settlements of Nairobi County; an area of study that poses serious public health concerns.

2. MATERIALS AND METHODS

2.1 Study Area

A cross-sectional study was conducted in the informal settlements; Korogocho (1.2504° S, 36.8909° E GPS coordinates) and Kariobangi North (1.2534° S, 36.88815° E GPS coordinates), Nairobi County (1.2921° S and 36.8219° E GPS coordinates), of Kenya. The 2009 national population census estimated the population of Nairobi County to be over three million, with over half of the population living in slums (Kenya National Bureau of Statistics, 2015). Street vended foods were collected in the following regions within Korogocho; Gitathuru, Nyayo, Kisumu Ndogo, Paradise and Kariobangi North.

2.2 Sampling and sample collection

Nairobi County was purposively selected for the study because of its populous nature. Korogocho and Kariobangi North areas were also purposively selected for the study as they are largely informal settlements with the population being those of low-income. A total of fifteen vendors were exhaustively sampled and included in the study with the food safety and hygiene practices evaluated using a food safety checklist. The snowballing sampling technique was used to locate all the vendors. Samples of raw and cooked street vended poultry products were subjected to microbial analysis. The respondents were purposively selected from the villages to include all the vendors selling the ready-to-eat (RTE) chicken products. All samples were collected in sterile polythene bags followed by transportation to the laboratory of the Department of Food Science and Technology of the University of Nairobi and microbial analysis done for *Escherichia coli*, *Salmonella*, *Staphylococcus aureus* and *Campylobacter jejuni*. The samples were stored at a temperature of 4 °C and analyzed within 24 hours of collection.

2.3 Microbial analysis

2.3.1 Determination of *Escherichia coli*

Based on ISO method 9308-1:2000 [18], the *E. coli* was accordingly enumerated. About 10g of the sample was homogenized in 90ml peptone water. Decimal serial dilutions of the homogenized solution in sterile peptone water were prepared and plated in duplicate on the selective agar media. Blue-green colonies for *E. coli* were counted after 48 hours of incubation at 44°C. The number of colony-forming units (CFU) of presumptive *E. coli* per gram of sample was calculated.

2.3.2 Determination of *Salmonella spp.*

The ISO method ISO 6579 [19], was used to enumerate the salmonella species. A sample of 10 g was weighed, homogenized in buffered peptone water and incubated at 37±1°C for 18±2 hours. From pre-enrichment broth, the inoculums were transferred to Rappaport-

Vassiliadis broth and selenite cystine broth and then incubated at $41.5\pm 1^{\circ}\text{C}$ and $37\pm 1^{\circ}\text{C}$ for 24 hours for selective enrichment. A loopful of the selective enrichment was streaked onto solid selective media: xylose lysine desoxycholate agar (XLD). XLD agar was incubated at $37\pm 1^{\circ}\text{C}$ and observed after 24 ± 3 hours for typical salmonella transparent red halo and a black centre.

2.3.3 Determination of *Staphylococcus aureus*

EN ISO method ISO 6888-1:1998 [20], was used for the detection and enumeration of *Staphylococcus aureus*. In a sterile pipette, 0.1ml of the appropriate sample test dilutions were transferred in duplicate onto the Baird Parker agar (BPA). The plates were then incubated at $35\text{-}37^{\circ}\text{C}$ for 24 ± 2 hours, then re-incubated for further 24 ± 2 hours. Observation ensued for typical colonies appearing black or grey, shining and convex, 1-1.5mm in diameter after 24 hours and 1.5-2.5mm after 48 hours of incubation, surrounded by a clear zone but a partially opaque zone. Coagulase positive staphylococci were then expressed as cfu/g of sample.

2.3.4 Determination of *Campylobacter jejuni*

The analysis was conducted according to ISO 10272-1:2017 [21] procedures which specify a horizontal method for the detection and enumeration of *Campylobacter* spp.

2.4 Risk assessment tools

Data generated from secondary sources from published articles in renowned databases including Science Direct, Elsevier, Springer, Hindawi and reports by global bodies like FAO and WHO were used. The data was used to respond to a set of eleven questions posed by the risk ranger and risk rank obtained [22]. The risk estimate was generated in a risk ranger which represented the relative risk of campylobacteriosis due to consumption of street-vended poultry. The spreadsheet uses its in-built functions to convert qualitative responses into numerical values that it uses to generate a risk rank [23]. The risk estimate generated by the Risk Ranger is usually on a scale of 0-100.

2.5 Data analysis

Genstat version 15 was used to analyse the microbial data. ANOVA was used to establish significant differences in the log counts of microbial pathogens. The LSD was used to separate means that were significantly different. Descriptive statistics including mean and SD of the mean microbial contamination levels were also generated. The risk estimate was generated from the risk ranger [23]. The risk estimate generated was on a scale of 0-100.

3. RESULTS AND DISCUSSION

3.1 Hazard Identification

Campylobacter jejuni which was the most prevalent microorganism in both raw and cooked poultry products at 8.95 ± 0.94 and 4.66 ± 2.67 log CFU g⁻¹ respectively; as compared to Salmonella, E. coli and Staphylococcus which had contamination levels of 6.42 ± 1.64 , 6.60 ± 1.25 and 6.92 ± 1.32 log CFU g⁻¹ in raw poultry respectively and 2.22 ± 1.88 , 2.67 ± 1.98 and 2.86 ± 1.61 in cooked poultry respectively $p < 0.05$. These findings are different from those established in another study done on street-vended poultry in Egypt where *Staphylococcus aureus* was found to be the most prevalent food pathogen [3]. Cardinale et al. (2015) in his study on street vended poultry in Madagascar established that there was no contamination of the products with *C. jejuni* as proper heat treatment of the products addressed the problem.

Campylobacter jejuni is a gram-negative, non-spore forming and motile microorganism [12]. The microorganism also has flagella which serve a role in its invasion [25]. The microorganisms account for about 90% of all human infections by *Campylobacter* sp. in human beings [26]. In recent times, the microorganism has been associated with enteritis and gastroenteritis in both the adult and pediatric patients [27]. About 30% of the cases of campylobacteriosis have been attributed to the consumption of poultry [28].

The microorganism induces food poisoning through intake of contaminated water or food [25]. *Campylobacter* isolation in patients suffering from diarrhoea in Kenya was reported as

12%, higher than for both *Salmonella* and *Shigella* [29]. Additionally, the prevalence of *Campylobacter* spp. in chicken in the informal settlements of Kenya was reported as 60-64% in the retail market. Mageto et al. (2018) reported that 32.5-76.5% of the *campylobacter* isolates from chicken in Nairobi County were *C. jejuni*. Another study by Nguyen et al. (2016) reported that 61.3% of the *C. jejuni* isolates from chicken in Kenya showed multi-drug resistance.

3.2 Hazard Characterization

The clinical manifestations of the infection are gastroenteritis, meningitis and acute cholecystitis [12]. The severity of the illness due to the infection by the microorganism was established as low as it was reported by Smith (1985) that the overall fatalities were 0.059 per 100,000 population. [33] reported that the illnesses would at times result in death thus rarely medical attention due to diarrheal episodes. Other diseases that are resultant from campylobacter infection include Guillain-Barré syndrome (GBS) and Miller Fisher syndrome. Gastroenteritis due to campylobacter infection is usually occasioned by diarrhoea, malaise, fever and abdominal pain and sometimes vomiting, inflammation of the intestinal mucosa, presence of blood in faeces and disruption of the epithelial cells [25]. The incidence of foodborne illnesses with diarrheal symptoms, in the study area, was found to be 52.9%. A study done in the informal settlements of Yaoundé, Cameroon, found that 59.5% of the diarrheal cases there were attributed to infectious microorganisms like *C. jejuni*. Deogratias et al. (2014) also reported a prevalence of 9.7% of campylobacter infection among under-five-year-old children in Tanzania with diarrheal infection. The global burden of campylobacteriosis in 2013 was reported as 7.5 million DALY [35].

The global data on GBS and Miller Fisher syndrome are so limited [35]. The GBS is characterized with sensory symptoms including sensation in the legs, rapidly progressive distal weakness, loss of vibration and proprioception and respiratory symptoms. WHO (2014) reported the disability weight of GBS at 0.445, lower than the one for gastroenteritis and enteritis. The illness, however, has life-long disability.

3.3 Exposure Assessment

Contamination of the poultry occurs at any given stage of the process of preparation of the poultry. The process of preparation of the street vended poultry in the informal settlements follows the schematic illustration shown in Figure 1. More than eight in every ten (87.5%) of the households in the informal settlements had the whole family as consumers of the street vended. Similar findings were reported in the informal settlements in India where both the adults and children were found to be consumers of street-vended [37]. The consumption of the street-vended chicken is either with or without further processing. Eight in every ten (82.8%) of these consumers had an intake of at least once a week. Namugumya and Muyanja (2011) also reported that poultry and meat products were one of the most frequent dishes of urban communities in Uganda. All the raw and cooked samples which were sampled were contaminated with *C. jejuni*. The microbial load of *C. jejuni* on the raw and cooked portion of chicken was reported as 8.95 ± 0.94 and $4.66 \pm 2.67 \log_{10}$ CFU g⁻¹ in this study. Another study that evaluated the raw portions of chicken in Burma and Ngara that

were reported by the vendors as the sources of the chicken found that they had a microbial load of $>4 \log \text{CFU g}^{-1}$ [39]. The high level of contamination in raw poultry was attributed to handling whereas undercooking was found to be responsible for the contamination in the cooked portions.

The informal settlements of Nairobi County host majorly the low economic class. The informal settlements in Nairobi that would be of greatest interest including Kibera, Mukuru Kwa Njenga, Mathare and Korogocho slums have an estimated population of 1.7 million people [40]. The low-income status of this area occasions the largest proportion of the residents to opt for the compromised quality of products. The study established that street food consumption in the area stood at 86%, which involved intake by the general household including children under the age of five years.

Through derivation from studies by Carron et al. (2018) and Mageto et al. (2018), the occurrence of *C. jejuni* in street vended raw poultry in the informal settlements was established as 48.96%. The high level of contamination in these raw portions are due to poor food handling practices [41]. The raw portions of poultry are usually roasted or deep-fried before the sale. Proper heat treatment has been proven as an effective strategy in eliminating the microbial counts of *C. jejuni* to undetectable levels [42, 43]. However, roasting as one of the fastest heat techniques has been indicated as one of the improperly practised food preparation techniques that enhance food safety risks and heat resistance in microbes [44]. Karoki et al. (2018) in his study showed that roasting would not reliably reduce the microbial counts in meat. Heating temperatures of $\geq 70^\circ\text{C}$ for about two minutes would reliably eliminate *C. jejuni*, though the cooking at the informal settlements is highly questionable.

The cooked samples of poultry had an average *C. jejuni* count of $4.66 \pm 2.67 \log_{10} \text{CFU g}^{-1}$ which was above the infective dose for *C. jejuni*; the infective dose of *C. jejuni* has been indicated to be low, $2.7\text{-}2.9 \log_{10} \text{CFU g}^{-1}$ [45]. The average weekly consumption sizes of poultry were found to be 140g per person. The intake levels found in this study were lower than those reported for both the children and adults who were reported to have consumption levels of 300g and 450 g respectively [37]. Further contamination of the cooked poultry products has been attributed to the poor post-processing handling that included contamination from the display surfaces and hands. This study established that all the display surfaces of the vendors were contaminated with *C. jejuni* averaging at $6.84 \pm 0.71 \log_{10} \text{CFU g}^{-1}$. Three-quarters of the vendors (76.9%) did not cover their food on display. There were no systems for control against post-process contamination. In as much as 72.1% of the vendors had been oriented on food safety issues, none of them (0%) had any formal training or expertise in food handling. Food safety training improves the food safety of processed food [46].




Fig. 1. : Process of preparing street vended poultry in informal settlements. Sources [47, 48].

3.4 Risk Characterization

No study has established the quantitative estimate of the risk posed by consumption of street vended poultry in the informal settlements. There is also no documented information on the process controls of the preparation of street foods. The information from the three previous steps of qualitative risk assessment was combined in the Risk Ranger software for the generation of a risk estimate. The probability of illness per day in a considered consumer was found as $7.12E-03$. Another study that evaluated the risk of *Campylobacter* infection due to consumption of ham reported that the probability of illness in an individual was $2.20E12$ [49]. Predicted illnesses in the population were found to be $1.11E06$.

The risk estimate generated for the consumption of finding to be 67. The level of risk posed is interpreted as to be a high risk, >48 [22]. The risk estimate is also higher than that posed by chicken consumed either in rural or urban china which was 52 and 49 respectively [50]. Another study in South Korea agreed that the outdoor, eating, of chicken and other poultry feeds, poses additional risks than the indoor [51]. This calls for better controls to be put in place to manage the currency risk. Proper cooking of the food and that which will reliably eliminate all hazards will reduce the risk posed in the consumption of street-vended poultry.

Table 1: Summary of Risk Ranger input data

Risk criteria	General population
Dose and severity	
Hazard severity	Mild hazard
Susceptibility	General, all members of the population
Probability of exposure	
Frequency of consumption	Weekly
Proportion consuming	Most (75%)
Size of population	1.7 million
Probability of consumption	
Probability of raw product contamination	48.96%
Effect of processing	The process usually (99% of cases) eliminates hazards
Possibility of recontamination	Yes- Major (50% frequency)
Post-process control	Not controlled-no systems, untrained staff (10-fold increase)
Increase to infective dose	Slight (10 fold increase)

Further cooking before eating	Meal preparation usually eliminates (99%) hazards
Probability of illness per day in the considered consumer	7.12×10^{-3}
Predicted cases of illnesses in the population	1.11×10^6
Risk ranking (0-100)	67

4. CONCLUSION

In as much as the street vended poultry have a greater preference and affordability, the food safety risks of consuming the products are very high. The risk of campylobacteriosis due to intake of street vended poultry is very high among the dwellers of the informal settlements. Less surveillance and disregard of important food hygiene and preparation practices creates this gap that endangers the lives of the consumers.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Oguttu JW. Participatory risk analysis of street vended chicken meat sold in the informal market of Pretoria , South Africa By. 2015;(April):1-254.
2. Climat R. Microbial safety aspects of street foods in Haiti. 2013.
3. Abd-El-Malek AM. Cooked poultry meat and products as a potential source of some food poisoning bacteria. IOSR J Environ Sci Toxicol Food Technol. 2017;11(06):23-29. doi:10.9790/2402-1106032329
4. Cardinale E, Abat C, Bénédicte C, Vincent P, Michel R, Muriel M. Salmonella and Campylobacter contamination of ready-to-eat street-vended pork meat dishes in Antananarivo, Madagascar: A risk for the consumers? Foodborne Pathog Dis. 2015;12(3):197-202. doi:10.1089/fpd.2014.1864
5. Tesfaye W, Bereda TW, Emerie YM, Reta MA, Asfaw HS. Microbiological Safety of Street Vended Foods in Jigjiga City , Eastern Ethiopia. (2).
6. Manguiat LS, Fang TJ. Microbiological quality of chicken- and pork-based street-vended foods from Taichung, Taiwan, and Laguna, Philippines. Food Microbiol. 2013;36(1):57-62. doi:10.1016/j.fm.2013.04.005

7. Food Standards Australia New Zealand. Scientific Assessment of the Public Health and Safety of Poultry Meat in Australia.; 2005.
8. Ologhobo AD, Omojola AB, Ofongo ST, Moiforay S, Jibir M. Safety of street vended meat products - chicken and beef suya. *African J Biotechnol.* 2010;9(26):4091-4095.
9. El-Shenawy M, El-Shenawy M, Mañes J, Soriano JM. *Listeria* spp. in Street-Vended Ready-to-Eat Foods . *Interdiscip Perspect Infect Dis.* 2011;2011:1-6. doi:10.1155/2011/968031
10. Rane S. Street Vended Food in Developing World: Hazard Analyses. 2011;51(1):100-106. doi:10.1007/s12088-011-0154-x
11. FAO, WHO. Risk Assessment of *Campylobacter* Spp . in Broiler Chickens and *Vibrio* Spp . in Seafood Report of a Joint FAO / WHO Expert.; 2002.
12. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global epidemiology of *campylobacter* infection. *Clin Microbiol Rev.* 2015;28(3):687-720. doi:10.1128/CMR.00006-15
13. Marotta F, Garofolo G, Donato G Di, et al. Population diversity of *campylobacter jejuni* in poultry and its dynamic of contamination in chicken meat. *Biomed Res Int.* 2015;2015:1-10. doi:10.1155/2015/859845
14. EFSA. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2017. *EFSA J.* 2014;16(12). doi:10.2903/j.efsa.2018.5500
15. Mughal MH. *Campylobacteriosis: A Global Threat.* *Biomed J Sci Tech Res.* 2019;11(5):1-5. doi:10.26717/bjstr.2018.11.002165
16. WHO, FAO. *FAO/WHO Guide for Application of Risk Analysis Principles and Procedures during Food Safety Emergencies.* Rome, Italy: FAO and WHO; 2011.
17. FDA/CFSAN. *DRAFT Qualitative Risk Assessment Risk of Activity / Food Combinations for Activities (Outside the Farm Definition) Conducted in a Facility Co-Located on a Farm.* Washington D.C., USA; 2012.
18. ISO. *Water Quality—Detection and Enumeration of Escherichia Coli and Coliform Bacteria.* Vol 2000.; 2000.
19. ISO. *ISO 6579. Microbiology of Food and Animal Feeding Stuffs. Horizontal Method for the Detection of Salmonella Spp.* 4 Ed. 2002. The International Organization for Standardization, Amendment 1: 15 Jul. 2007. Vol 2002. Geneva, Switzerland: ISO; 2002.
20. ISO. *Microbiology of Food and Animal Feeding Stuffs- Horizontal Method for Enumeration of Coagulase Positive Staphylococci (Staphylococcus Aureus and Other Species).* Vol 1999. 1st ed. Geneva, Switzerland: ISO; 1999. doi:10.1088/0957-4484/25/12/125707
21. ISO. *Microbiology of Food Chain: Horizontal Method for Detection and Enumeration of Campylobacter Spp.;* 2017.
22. FAO. *Application of risk assessment in the fish industry.*
23. Ross T, Sumner J. A simple , spreadsheet-based , food safety risk assessment tool. *Int J Food Microbiol* 77. 2002;77:39-53.
24. Cardinale E, Abat C, Benedicte C, Vincent P, Michel R, Muriel M. *Salmonella and Campylobacter Contamination of Ready-to-Eat Street-Vended Pork Meat Dishes in Antananarivo, Madagascar: A Risk for the Consumers? Food borne Pathog Dis.* 2015;12(3):197-202. doi:10.1089/fpd.2014.1864
25. Quetz da JS, Lima IFN, Havt A, et al. *Campylobacter jejuni* infection and virulence-associated genes in children with moderate to severe diarrhoea admitted to emergency rooms in northeastern Brazil. *J Med Microbiol.* 2012;61(4):507-513. doi:10.1099/jmm.0.040600-0

26. James C, Daramola B, Reyers F, Purnell G, Turner R, James SJ. Qualitative Risk Assessment to Support a Policy Decision on Partially - Eviscerated (Effilé) Poultry Production.; 2014.
27. Facciola A, Riso R, Avventuroso E, Visalli G, Delia SA, Laganà P. Campylobacter: from microbiology to prevention. *J Prev Med Hyg.* 2017;58(2):E79-E92.
28. Chlebicz A, Śliżewska K. Campylobacteriosis, Salmonellosis, Yersiniosis, and Listeriosis as zoonotic foodborne diseases: A review. *Int J Environ Res Public Health.* 2018;15(5):863. doi:10.3390/ijerph15050863
29. Carron M, Chang YM, Momanyi K, et al. Campylobacter, a zoonotic pathogen of global importance: Prevalence and risk factors in the fast-evolving chicken meat system of Nairobi, Kenya. *PLoS Negl Trop Dis.* 2018;12(8):1-18. doi:10.1371/journal.pntd.0006658
30. Mageto L, Ombui JN, Mutua FK. Prevalence and risk factors for Campylobacter infection of chicken in peri-urban areas of Nairobi, Kenya. *J Dairy, Vet Anim Res.* 2018;7(1):22-27. doi:10.15406/jdvar.2018.07.00184
31. Nguyen TNM, Hotzel H, Njeru J, et al. Antimicrobial resistance of Campylobacter isolates from small scale and backyard chicken in Kenya. *Gut Pathog.* 2016;8(1):1-9. doi:10.1186/s13099-016-0121-5
32. Smith GS. Fatalities Associated With Campylobacter jejuni Infections. *JAMA J Am Med Assoc.* 1985;253(19):2873. doi:10.1001/jama.1985.03350430085033
33. WHO. Campylobacter.
34. Deogratias AP, Mushi MF, Paterno L, et al. Prevalence and determinants of Campylobacter infection among under five children with acute watery diarrhea in Mwanza, North Tanzania. *Arch Public Heal.* 2014;72(1):1-6. doi:10.1186/2049-3258-72-17
35. WHO, OIE, FAO, Universiteit Utrecht. The Global View of Campylobacteriosis: Report of an Expert Consultation, Utrecht, Netherlands, 9-11 July 2012. Utrecht, Netherlands; 2013. doi:ISBN 978 92 4 156460 1
36. WHO. WHO estimates of the global burden of diseases. *Who.* 2014;46(3):1-15. doi:10.1016/j.fm.2014.07.009
37. Selvan P, Preethi NS. Exposure assessment of Bacillus cereus in street vended chicken biryani – a deli food of South India. *Int J Curr Microbiol Appl Sci.* 2018;7(12):2894-2899.
38. Namugumya BS, Muyanja C. Contribution of street foods to the dietary needs of street food vendors in Kampala , Jinja and Masaka districts , Uganda. *Public Health Nutr.* 2011;15(8):1503-1511. doi:10.1017/S1368980011002710
39. Odhiambo WA, Kebira AN, Nyerere A. Prevalence of Campylobacter jejuni and other bacterial pathogens in selected foods and drinks served in fast food kiosks in Ngara and Burma markets in Nairobi. *Int Acad J Heal Med Nurs.* 2017;1(1):32-57.
40. Mutisya E, Yarime M. Understanding the grassroots dynamics of slums in Nairobi: The dilemma of Kibera informal settlements. *Int Trans J Eng Manag Appl Sci Technol.* 2011;2(2):197-213.
41. Alimi BA. Risk factors in street food practices in developing countries: A review. *Food Sci Hum Wellness.* 2016;5(3):141-148. doi:10.1016/j.fshw.2016.05.001
42. Josefsen MH, Löfström C, Hansen TB, Christensen LS, Olsen JE, Hoorfar J. Rapid quantification of viable Campylobacter bacteria on chicken carcasses, using real-time pcr and propidium monoazide treatment, as a tool for quantitative risk assessment. *Appl Environ Microbiol.* 2010;76(15):5097-5104. doi:10.1128/AEM.00411-10
43. de Jong AEI, van Asselt ED, Zwietering MH, Nauta MJ, de Jonge R. Extreme heat resistance of food borne pathogens Campylobacter jejuni, Escherichia coli , and Salmonella typhimurium on chicken breast fillet during cooking. *Int J Microbiol.* 2012;2012:1-10. doi:10.1155/2012/196841

44. Karoki WH, Karanja DN, Bebora LC, Njagi LW. Isolation, characterization, and quantification of bacteria from African sausages sold in Nairobi County, Kenya. *Int J Food Sci.* 2018;2018:1-9. doi:10.1155/2018/3861265
45. Epps SVR, Harvey RB, Hume ME, Phillips TD, Anderson RC, Nisbet DJ. Foodborne *Campylobacter*: Infections, metabolism, pathogenesis and reservoirs. *Int J Environ Res Public Health.* 2013;10(12):6292-6304. doi:10.3390/ijerph10126292
46. Adesokan HK, Akinseye VO, Adesokan GA. Food safety training is associated with improved knowledge and behaviours among foodservice establishments workers. *Int J Food Sci.* 2015;2015:1-8. doi:10.1155/2015/328761
47. Muth MK, Beach RH, Karns SA, Taylor JL, Viator CL. *Poultry Slaughter and Processing Sector Facility-Level Model.*; 2006.
48. Carron M, Alarcon P, Karani M, et al. The broiler meat system in Nairobi, Kenya: Using a value chain framework to understand animal and product flows, governance and sanitary risks. *Prev Vet Med.* 2017;147:90-99. doi:10.1016/j.prevetmed.2017.08.013
49. Lee J, Ha J, Kim S, Lee H, Lee S, Yoon Y. Quantitative microbial risk assessment for *Staphylococcus aureus* in natural and processed cheese in Korea. *Korean J Food Sci An Vol.* 2015;35(5):674-682. doi:10.3168/jds.2015-9611
50. Wang J, Guo YC, Li N. Prevalence and risk assessment of *Campylobacter jejuni* in chicken in China. *Biomed Environ Sci.* 2013;26(4):243-248. doi:10.3967/0895-3988.2013.04.002
51. Jeong J, Lee J, Lee H, et al. Quantitative microbial risk assessment for *Campylobacter* foodborne illness in raw Beef offal consumption in South Korea. *J Food Prot.* 2017;80(4):609-618. doi:10.4315/0362-028x.jfp-16-159