

## Colorectal adenocarcinoma in East Africa: A narrative literature review

### Author Contributions

The author read, edited and approved the final manuscript.

### ABSTRACT

This paper is a review of work done on colorectal adenocarcinoma in East Africa showing geographic spread, age and sex ratios, clinical presentation, management and predominant histopathology. A steady increased incidence of CRC in East African countries is currently being documented however this is associated with a higher CRC-associated morbidity and mortality. Whilst the male: female ratio varies between 1.2:1 to 1.88:1, up to 38% of CRC diagnosis are in patients younger than 40 years, in contrast to only 1.9% of CRC patients in Western developed countries such as the USA. Generally rectal carcinoma is more common than colon carcinoma and abdomino-perineal resections are commonly performed in up to between 54% - 71% due to the advanced stage of presentation of rectal tumours in East Africa. The late stage presentation and delayed effective treatment in East Africa may result in a higher morbidity in CRC patients. Interestingly there is a significant incidence of mucinous adenocarcinoma sub-groups compared to Western developed countries which carry a poor prognosis. A significant proportion of CRC patients have been found to have histological and demographic features which suggest that MSI-tumours and these tumours are more common in younger patients. However only a few authors have looked at the possibility of mismatch repair mutations in the genetic aetiopathogenesis of colorectal adenocarcinoma in East Africa.

*Keywords: CRC: colorectal carcinoma; HNPCC: hereditary nonpolyposis colorectal carcinoma; MSI: microsatellite instability; MSI-H: high amount of microsatellite instability*


### INTRODUCTION

Colorectal cancer is the fourth leading cause of cancer death in the world (1,2). It is the third most commonly diagnosed malignancy in the world and is responsible for 1.4 million new cases and approximately 700,000 deaths in the year 2012 (2). CRC is the second most common cancer in women worldwide (614,000 cases; 9.2% of the total) and the third most common cancer in men (746,000 cases; 10% of the total) (3). In Africa colorectal carcinoma is a rare disease and it currently represents 2-6% of all malignant tumors (4,5).

The crude incidence of CRC in Sub-Saharan Africa is 3.69/100,000 for women and 4.38/100,000 for men (overall 4.04/100,000) with a wide variation in the geographical distribution of CRC reported on the African continent. A high incidence in South Africa of 11.9/100,000 whilst low incidences reported in East Africa (6.5/100,000) and West Africa (3.8/100,000 in women; 4.5/100,000 in men) has been reported (3, 4, 5).

In between the years 1957 and 1964, 105 cases of colorectal carcinoma were reported in Kenya giving an incidence of 2.5/100,000 (6). Eshelman (1966) and Hiza (1974) reported a low incidence of colorectal carcinoma in Tanzania with only four cases of rectal carcinoma and none of colon carcinoma reported (7,8,9). However, in a recent study by Siegel from Sub-Saharan Africa has shown that in urban areas the incidence of colorectal carcinoma is indeed increasing (6). Generally, in many East African countries there has been an increased incidence in the incidence rate of CRC. In Kenya there has been an increased incidence of colorectal carcinoma (4.8%) and it is currently the fourth commonest carcinoma in the country

(10). The population incidence of CRC in Kenya is currently 9.6 per 100,000 populations (10). In Uganda, the Kampala Cancer Registry has shown that colorectal carcinoma has a low incidence however there are increases occurring especially among women (11). In 1977, Ssali et al, reported a low incidence of colon carcinoma at 2.8% and stated that the incidence may be rising (12). The age standardized incidence rate has since remained oscillating around 7.8 per 100,000 population in males for the years 1991-2010 (11). The age standardized incidence rate has increased from 5.2 per 100,000 population for the year period 1991-1995 to 9.0 per 100,000 population for the year period 2006-2010 in females (11). This gives a 4.1% annual percentage change in the incidence of colorectal carcinoma in females in Uganda which is a greater increase than in males (11). Figure 1 shows a bar chart depicting the steady increase in incidence of CRC in East African countries over the last few decades. The increase in the percentage change in age-standardized incidence rates between the year 1990 and the year 2017 is 22.1% in Uganda, 17.5% in Kenya, 5.7% in South Sudan, 13.6% in Somalia, 14.3% in Eritrea and 7.2% in Djibouti (11).



**Figure 1:** A bar graph showing the age standardized incidence rates of colorectal adenocarcinoma per 100,000 person-years in East African countries depicting a steady increase in the incidence of CRC between the years 1990 to the year 2017

Despite the incidence of CRC being low in East Africa, there is a higher CRC-associated morbidity and mortality (1). The increasing incidence of CRC being reported in East Africa is in turn associated with an increased rise in the burden of non-communicable diseases in East Africa (6). Improved cancer awareness,

an increase in reporting and a shift towards a western diet may be attributing towards the rising incidence of colorectal carcinoma in East Africa (1,6). The increased incidence may also probably be due to an increase in risk related behaviors such as smoking, alcohol, obesity and dietary changes particularly in urban areas of Sub Saharan Africa (4).

The rarity of colorectal carcinoma in the past and the current steady increase in CRC cases in East Africa makes confirmed cases today worthy of attention. This literature review looks at the work done on colorectal carcinoma by researchers from countries in East Africa to illustrate the results of these authors from the last 20 years.

## **DISCUSSION**

Several factors are reported to be linked to the incidence of colorectal adenocarcinoma which include environmental and hereditary factors. Changes in dietary preferences is a result of an important environmental influence which is related to economic development (13). Economically developed countries such as the UK and USA have a high incidence of CRC however in East Africa a steady increase is being reported (1,6). Over the past two decades an increase in the incidence of CRC affecting rural populations in East Africa of low socio-economic status has also been reported (14). This paper looks at as many publications available in East Africa to illustrate the rate and pattern of CRC in this part of the world over the past years.

### **Age and sex distribution of colorectal carcinoma**

In East Africa a significant proportion of patients are young with a median age at diagnosis of 41-59 years (15-18). Between 19% to 38% of patients younger than 40 years of age present with CRC (16,17). This is in contrast to only 1.9% of patients in the USA who are younger than 40 years and present with CRC (19). Among African-American patients, early-onset CRC has also been described (20). In Kyadondo in Uganda, colorectal carcinoma has been found to be more common in women whilst in the Western world men have a higher incidence of CRC (11). In Uganda, colorectal carcinoma is diagnosed at a median age of 55 years (11). 30.6% of CRC patients are under 50 years of age, 6.9% of CRC patients were under 30 years of age and 69.4% of CRC patients were either 50 years or above in a study from Uganda (11). This finding suggests that genetic factors may play a part in the aetiology of this early onset CRC in African patients (20). The male to female ratio in East African studies currently report a ratio of 1.2:1 to 1.88:1 (14,16,17,21).

### **Presentation of patients with colorectal carcinoma**

Patients present in a way that is determined by the anatomical location of the CRC in the large bowel. In East Africa several studies have shown that there is a predominance of left sided colorectal tumors in particular rectosigmoid tumors (17, 22, 23). In Uganda, Dijkhoorn identified only 2 tumors (5.1%) that were in the right colon whilst 37 tumors (94.9%) were located in the left colon (24). 32 of these 37 tumors left colonic tumors were located in the rectum (86.5%) (24). In Kenya, Saidi H, found 50% of the tumors located in the rectum, 17.1% in the sigmoid colon, 12.9% in the ascending colon, 11.4% in caecum and 7.1% in the ascending colon (17). Table I shows studies from East Africa showing the anatomical site location of CRC and percentages together with the most frequent site of CRC from the data presented.

In developed countries a right-sided preponderance of colorectal carcinoma (proximal shift) is generally seen (25,26). In a Louisiana population, Minardi et al found CRC to be evenly distributed throughout the colon and rectum (27). Whilst only 30% of patients had rectal involvement in a study reported by Kam et al in Singapore (28). The reason for this anatomical difference between East African countries and developed countries is not clear however the reason could be that the rectum and proximal sigmoid colon could be more accessible to investigate by rigid proctosigmoidoscopy than the right colon (29).

Authors	Saidi H et al., 2007	Chalya PL et al., 2013	Dijxhoorn DN et al., 2014	Medhin LB et al., 2019	Deressa et al., 2019
Caecum (%)	11.4	12			
Ascending colon (%)	7.1	7.5	5.1	-	7
Transverse colon (%)	-	3.9	-	-	-
Descending colon (%)	12.9	6	13.5 (including sigmoid colon)	-	2
Sigmoid colon (%)	17.1	-	-	-	3
Rectosigmoid junction (%)	-	54.8	-	-	10
Rectum (%)	50	15.7	86.5	50	69
Most frequent site of CRC	Rectum	Rectosigmoid	Rectum	Rectum	Rectum

**Table 1:** Studies in East Africa depicting period of study, anatomical site of tumors, most frequent anatomical site and percentages. The symbol – denotes data was not available for these anatomical sites in these studies.

In many East African hospitals proctosigmoidoscopy is the only available method of obtaining biopsies in patients with lower gastrointestinal symptoms. There is poor access to colonoscopy facilities and CT scanning with many patients having a laparotomy for diagnosis of CRC in many East African hospitals (29).

A study from Kenya showed that only 23% of patients with colorectal carcinoma had colonoscopy during the years 1993-1998 which improved to 51% during the years 1999-2005 (17). Similarly a retrospective study from Tanzania also showed that from a total of 901 cases, rectal cancer (482) was more common compared to colon cancer (344) and (75) rectosigmoid junction cancer (30). Colon cancer was most often diagnosed in males compared to females (162:182) whilst rectum cancer was most often diagnosed in females compared to males (265:219) (30). Katalambula et al noted that the risk of colon cancer was significantly higher in patients below 35 years. The same study found that rectum cancer was more common in older patients 80+ years (78.13%) and in patients 50-54 years old. However, Dijxhoorn et al noted no significant difference between males and females, and no significant difference in age i.e. younger or older patients (<50 years or >50years) regarding tumor site (24).

The world age-standardized incidence rate is higher for colon carcinoma than for rectum carcinoma in both sexes in Uganda. In colon carcinoma it stands at 4.0 per 100,000 in males and 4.1 per 100,000 in females for the period years 2008 to 2012 (4). In rectum carcinoma the world age-standardized incidence rate is lower at 3.8 per 100,000 population in males and 3.5 per 100,000 population in females for the year period 2008 to 2012 (4). Between the years 1998 to 2007 the colon to rectum cancer rate ratio has been variable in Uganda between 0.5 - 2.5: 1 which contrasts with that seen in the US and UK where the ratio is steadily constant at 2:1 (4). Table 2 shows the few studies carried out from countries in East Africa which depict the period of study, number of patients studied, sex ratios, ages and colon: rectum ratios.

Authors	Period (Year)	Number of Patients	Mean Age	Male:Female	Colon:Rectum
Saidi H et al., 2011 (Kenya)	2005-2010	233	53	1.43:1	1:1.02
Chalya PL et al., 2013 (Tanzania)	2006-2011	332	46	1.6:1	5.38:1
Dijxhoorn DN et al., 2014 (Uganda)	2006-2010	81	55	1:1.15	1:4.57
Medhin LB et al., 2019 (Eritrea)	2011-2017	241	57.62	1:1.61	1:1
Deressa BT et al., 2019 (Ethiopia)	2016-2017	147	46	1.88:1	1:4.21

**Table 2 :** Studies in East Africa depicting period of study, number of patients studied, sex ratios, ages and colon:rectum ratio

Some studies have reported rectal bleeding as the commonest symptom reported from colorectal cancer patients in East Africa (16,17). Distinguishing a benign from a malignant lesion such as CRC may pose a diagnostic challenge in our environment. A study from the UK showed that the prevalence of rectal bleeding was 14% (31,32). Several studies has shown that in elderly patients rectal bleeding can predict colorectal cancer especially when in combination with abdominal pain or a change in bowel habit (33,34). It is therefore reasonable to investigate rectal bleeding by colonoscopy in all cases. Colonoscopy should be carried out for patients presenting with rectal bleeding, change in bowel habit and anaemia and may consequently improve survival in these patients. Patients presenting with right sided CRC tend to have significantly lower haemoglobin levels and these tumors tend to bleed easily in contrast with left sided tumors which present with large bowel obstruction (17). Therefore colonoscopy should be carried out for adult patients presenting with anaemia with suspected CRC (35-37). In another study by Saidi HS et al, the most common presenting symptom was abdominal pain (76.9%). The same study reported a change in bowel habit (71.4%), rectal bleeding (54.3%), swelling (34.8%), tenesmus (17.1%), weight loss (14.3%) and bowel obstruction (14.3%) (17). Abdominal pain has been found to be the presenting symptom in other studies, in fact in one account 54% of patients presented with this symptom (34).

Due to patient variability, a change in bowel habit may be difficult to document. A study in Kenya reported a change in bowel habit in 71.4% of CRC patients (17). In the UK, studies have shown that diarrhea and increased stool frequency are more predictive of CRC than constipation (38). In East Africa, since left sided colonic and rectal lesions are more common and they commonly present with large bowel obstruction then this symptoms should dictate early referral to hospital and this will inevitably have a positive effect on survival.

A study from Tanzania found that only 3.3% of patients present with early stage CRC (Stage I) compared to 90% presenting with advanced stage CRC (Stage II, III and IV) (16). In Eritrea, late presentation of CRC is also a common feature and a high proportion of patients (61.7%) present with advanced stages (stage 2 and above) (21). These findings are consistent with findings from other studies in Africa (17, 39, 40, 41,42). This is mainly due to the absence of screening facilities for colorectal carcinoma in the East African population. There is considerable evidence that screening with faecal occult blood test and colonoscopy of asymptomatic individuals at an average risk can detect CRC at an early and curable stage which results in

a reduction in mortality (43). Colonoscopy may detect cancer-precursor lesions, such as polyps which when removed reduce the incidence of CRC (44,45). The late presentation reported in all East African countries may be due to lack of screening programs, lack of accessibility to health-care facilities and lack of patient awareness with the disease.

### Treatment of Colorectal adenocarcinoma patients

The main aim of surgeons managing colorectal adenocarcinoma is curative resection. In East Africa many patients present at an advance stage and so a number of surgical interventions have been reported. The majority of surgical operations for rectal carcinoma are abdomino-perineal resections which is performed in up to between 54%-71% of patients (46-49) due to the advanced stage of presentation of rectal tumors. Anterior resections for rectal tumors have been reported in only up to between 22% - 26% of patients (46, 49) possibly because of the lack of facilities to provide chemotherapy and radiotherapy pre-operatively in rectal carcinoma in East Africa due to the advanced stage of presentation of rectal carcinoma.

The frequency of other operations documented in East African studies include right hemicolectomy in 22.1-43% of patients (16, 46), left hemicolectomy in 33% - 58.6% of patients (10, 40), transverse colectomy in 3% - 12% of patients and sigmoid colectomy in 3% of patients(16,46). Table 3 shows the frequency of operations documented in East African studies. A colostomy itself may lead to psychological distress and social stigma and has been reported to lead to suicide and this is actually not related to the location of the CRC tumor (50). Older patients (>70 years) and a marital status of single have been found to be associated with a higher risk of suicide (50-52). Erectile impotence in male patients may result from disruption of the pelvic nerves intra-operatively or psychologically due to the presence of a stoma (53,54). Since many rectal cancer operations occur in young Africans that are sexually active, the impact operations such as abdomino-perineal resection has on psychological wellbeing needs to be factored in when treatment options are being considered.

Surgical Procedure	Saidi H et al., 2011 Frequency	Saidi H et al., 2011 Percentage(%)	Chalya PL et al., 2013 Frequency	Chalya PL et al., 2013 Percentage(%)	Deressa BT et al., 2019 Frequency	Deressa BT et al., 2019 Percentage (%)
Rt.Hemi-colectomy	43	22.9	72	22.1	-	-
Transverse colectomy	3	1.6	12	3.7	-	-
Lt. Hemi-colectomy	33	17.6	191	58.6	-	-
Hemi-Colectomy (not specified)	-	-	-	-	18	31
Sigmoid colectomy	3	1.6	-	-	-	-
Anterior resection	22	11.7	8	2.5	8	26
Abdomino-perineal resection	54	28.7	28	8.6	22	71
Colostomy/ Biopsy only No surgery	30	15.9	15	4.6	-	-

**Table 3:** Studies in East Africa depicting period of study, frequency of the different types of operations and percentages together with the authors of the studies. The symbol – denotes data was not available for these operations in these studies.

Complete resection with adjacent involved lymph nodes in CRC in the early stage is the only chance for cure (23,54). However many patients with CRC present at a late stage in East Africa and therefore only palliative surgery in the form of a colostomy is possible. Only approximately 3.3% of CRC in East Africa is curable by complete resection in keeping with many studies from developing countries (17, 39, 40) due to many patients presenting at an advanced stage.

Chemotherapy and radiotherapy play an important role in locally advanced colorectal cancer (55). Unfortunately in resource poor countries in East Africa, limited access to chemotherapy poses a major challenge in the treatment of CRC. In most countries, drug side effects and financial difficulties are the main reasons for no adherence to this form of treatment (23). In locally advanced CRC, radiotherapy when applied preoperatively or postoperatively has been shown to improve local control rates. Studies in resource limited countries have shown that only a few CRC patients have access to this form of treatment and one study reported that only 7.2% of patients had access to radiotherapy (16, 23, 56).

#### **Outcome of management of colorectal adenocarcinoma**

The higher post-operative complication rate seen in patients with CRC at ~26% may be due to the late stage presentation and delayed effective treatment in East Africa (16). One study reported a high rate of wound sepsis at 41.9% possible due to the late presentation of CRC with large bowel obstruction (16). Perineal wounds especially perineal wound breakdown may result in the need for another operation which results in increased hospital stay and costs.

The mortality rates of CRC in East Africa have ranged from 10.5% to 29.4% which is higher than that reported in West Africa (6.1%) (16, 46, 57). This high mortality may be due to the fact that many CRC patients in East Africa present at a late stage. In Uganda the 5-year relative survival of CRC has been found to be as low as 8.3% which is in contrast with the 5-year relative survival of 54.2% for Black American patients (58). In Kenya the 3-year survival of CRC was 40% which contrasts with a better rate at a longer follow up of 65% for whites at five years. Lack of patients' awareness to present to hospital early with symptoms, lack of screening facilities, lack of adequate treatment facilities, patients presenting at a late stage and differing tumor biology may explain this discrepancy in survival in East Africa. However the determinants of survival of CRC in East Africa have not been carefully examined hence further studies are needed to address this issue.

#### **Histopathology**

Moderately differentiated adenocarcinoma is the predominant histopathological type of CRC reported in a few studies from Uganda and Tanzania as 88.9% and 98.9% respectively (16,24). The same study of 81 cases from Uganda showed that mucinous adenocarcinoma was more common in younger compared to older patients however the presence of signet ring cells was similar (24). Poorly differentiated adenocarcinoma is more common in younger patients than those diagnosed at an older age(24). Whilst the study from Tanzania showed that mucinous and signet ring carcinoma are significantly more common under 40 years (16).The proportions reported from this study were mucinous adenocarcinoma (11.6%), signet ring adenocarcinoma (4.6%) and anaplastic (undifferentiated) carcinoma (5.2%) (16). These two studies from Uganda and Tanzania showed that mucinous adenocarcinomas were more frequent in the right colon than the left colon. A poor prognosis was seen with mucinous and signet ring adenocarcinomas compared to non-mucinous adenocarcinomas (16,24,47). These findings in East Africa concur with other findings from West Africa (51). These same findings also suggest that there is a higher incidence of mucinous and anaplastic tumors in Africa compared to the Western world.

## **Hereditary non-polyposis colon cancer - Lynch syndrome**

There has been an increase in the incidence of a unique early-onset, non-polypoid type of CRC affecting the rural population of Uganda, whose diet consisting of high-fibre, vegetables and fruits and low in fat significantly differs from the high fat diet in high income countries (14). Interestingly a study in Uganda showed that 27.3% of patients identified with CRC had histological and demographic features suggestive of MSI tumors based on the Revised Bethesda Guidelines (24). Mucinous and poorly differentiated adenocarcinoma were more likely to be diagnosed on histology in patients under the age of 50 and this suggests that MSI tumors are more common in younger patients (24). The age distribution and morphological pattern of CRC in Uganda hence differs from published data reported in the UK and US (24).

In East Africa, the connection between family history and the occurrence of CRC is under investigated (21). Only a few authors have looked at the possibility of mismatch repair mutations in the aetiopathogenesis of colorectal adenocarcinoma in East Africa (24). Therefore recognizing that a contribution of the aetiology of CRC in East Africa may be due to hereditary genetic factors makes immunohistochemistry testing and MSI testing imperative to be carried out by researchers in the region in order to truly establish the molecular subtypes of colorectal adenocarcinoma and hence the prevalence of Lynch syndrome in our community.

### **Conclusions**

In East Africa there is a steady increase in the number of CRC patients being treated. This may be due to an increase in the number of cancer registries in the region reporting this cancer. The male: female ratio varies, whilst the median age is around 41-59 years with more left sided CRC than right sided CRC. Mucinous adenocarcinoma which carry a worse prognosis have a higher incidence than the Western world. The majority of authors have stated that familial polyposis syndromes are a rarity in East Africa however some authors have found similarities in the demographics, histopathological type and non-polypoid MSI-H CRC found principally in black Americans.

Colorectal cancer in East Africa requires more unravelling especially in the aetiopathogenesis and therefore immunohistochemical staining and MSI testing is being recommended to establish the molecular subtypes and hence the prevalence of Hereditary Non-Polyposis colorectal carcinoma (HNPCC). This will clarify the genetic aetiopathogenesis of colorectal adenocarcinoma in the East African region.

### **ETHICAL APPROVAL**

This narrative literature review was partly taken from the PhD proposal of the author which was approved by the Higher Degrees Research and Ethics Committee, School of Biomedical Sciences, College of Health Sciences, Makerere University, Kampala, Uganda and Uganda National Council for Science and Technology and both approvals are preserved by the author.

### **ACKNOWLEDGEMENTS**

The author declares that this narrative literature review was partly taken from his approved PhD proposal. The author wishes to express his sincere gratitude to Professor Michael Odida and Professor Henry Wabinga, School of Biomedical Sciences, College of Health Sciences, Makerere University for their supervisory contribution to his PhD proposal.

### **COMPETING INTERESTS**

The author declares that no competing interests exist.

### **REFERENCES**

1. Arnold M, Sierra MS, Laversanne M et al. Global patterns and trends in colorectal cancer incidence and mortality. Gut 2016; published online Jan 27. DOI: 10.1136/gutjnl-2015-310912.



2. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh JW, Comber H, Forman D, Bray F. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer*. 2013 Apr; 49(6): 1374-1403. DOI: 10.1016/j.ejca.2012.12.027. Epub 2013 Feb 26.ewasa
3. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *In J Cancer* 2015; 136: E359-86.
4. Center M, Jemal A, Smith R, Ward E. Worldwide variations in colorectal cancer. *CA Cancer J Clin* 2009; 59: 366-78.
5. Weitz J, Koch M, Debus J, Hohler T, Galle PR, Buchler MW. Colorectal cancer. *Lancet* 2005, 365: 153-165.
6. Siegel RL, Miller KD, Jemal A. Cancer Statistics. 2015. *CA: a cancer Journal for 2015; Clinicians*: 65(1): 5-29.
7. Katsidzira L, Gangaidzo I, Thompson S, Rusakaniko S, Matenga J, Ramesar R. The sifting epidemiology of colorectal cancer in sub-saharan Africa. *Lancet Gastroenterol Hepatol* 2017; 2: 377-83.
8. Okobia MN. Cancer care in Sub-Saharan Africa – urgent need for population-based cancer registries. *Ethiop. J. Health. Dev.* 2004; 17: 89-98.
9. Holcombe C & Babayo U. The pattern of malignant disease in north east Nigeria. *Tropical and geographical medicine*. 1991; 43(1-2), 189-192.
10. Korir A, Okerosi N, Ronah V, Mutuma G, Parkin M. Incidence of cancer in Nairobi, Kenya (2004-2008). *Int. J. Cancer*. 2015: 137; 2053-2059.
11. Wabinga HR, Namboozee S, Amulen PM, Okello C, Mbus L, Parkin DM. Trends in the incidence of cancer in Kampala, Uganda 1991-2010. *Int. J. Cancer*. 2014 Jul 15: 135(2): 432-9.
12. Ssali JC. Carcinoma of the large bowel in Uganda. *Ann. R. Coll. Surg. Eng.* 1977; 59(5): 402-2.
13. Zhang T. The effects of economic development and built environment of diabetes in China. *Popul Health Metr* 2017; 15:35.
14. Makobore P, Masiira N, Elobu E. Characterization of colorectal carcinoma in Uganda: Is Ugandan Tumor Unique? *Annals of Oncology*. Vol.23, Suppl. 4. June 2012.
15. Walker A, Segal I. Colorectal cancer in an African city population in transition. *Eur. J. Cancer Prev.* 2002; 11: 187-91.
16. Chalya PL, Mchembe MD, Mabula JB, Rambau PF, Jaka H, Kay M, Mkongo E, Masalu N. Clinicopathological patterns and challenges of management of colorectal cancer in a resource-limited setting: a Tanzanian experience. *World Journal of Surgical Oncology* 2013; 11:88; <http://www.wjso.com/content/11/1/88>.
17. Saidi HS, Karuri D, Nyaim EO. Correlation of clinical data, anatomical site and disease stage in colorectal cancer. *East African Medical Journal* Vol. 85; No.6; June 2008.
18. Ashenafi S. The frequency of large bowel cancer as seen in Addis Ababa University, Pathology Department, *Ethiop. Med. J.* 2000; 38: 277-282.
19. Siegel RL, Fedewa SA, Anderson WE, Miller KD, MaJ, Rosenberg FS, Jemal A. Colorectal cancer incidence patterns in the United States, 1974-2013. *J Natl Cancer Inst.* 2017 Aug 1; 109(8). DOI: 10.1093/jnci/djw322.
20. Siegel RL, Jemal A, Ward EM. Increase in incidence of colorectal cancer among young men and women in the United States. *Cancer Epidemiol Biomarkers Prev* 2009 Jun; 18(6): 1695-8 DOI: 10.1158/1055-9965.EPI-09-0186.
21. Medhin LB, Achila OO, Abraham AT, Efreem B, Hailu K, Abraha DM. Incidence of colorectal cancer in Eritrea: Data from the National Health Laboratory, 2011-2017; 2019; *PLOS ONE* 14(11): e0224045. <https://doi.org/10.1371/journal.pone.0224045>.
22. Sule AZ, Mandong BM: Malignant colorectal tumors in patients 30 years and below: a review of 35 cases. *Cont. Afr. J. Med.* 1999, 45: 209-212.
23. Yawe KT, Bakart AA, Pindiga KH, Mayun AA: Clinicopathological pattern and challenges in the management of colorectal cancer in Sub-Saharan Africa. *J. Chinese Clin. Med* 2007, 2: 688-695.
24. Dijkhoorn DN, Boutall A, Mulder CJ, Ssebuufu R, Mall A, Kalungi S, Baigrie C, Goldberg PA. Colorectal cancer in patients from Uganda: A histopathological study. *East Centr. Afr. J. Surg.* 2014; Vol 19: 112-119.

25. Takada H, Ohsawa T, Iwamoto S, Yoshida R, Nakano M, Imada S, Yoshioka K, Okuno M, Masuya Y, Hasegawa K, Kamano N, Hioki K, Muto T, Koyama Y. Changing site distribution of colorectal cancer in Japan. *Dis Colon Rectum* 2002 Sep; 45(9): 1249-54.
26. De Leon PM, Benatti P, Borghi F, Pedroni M, Searselli A, DiGregario C, Losi L, Viel A, Genuardi M, Abbati G, Rossi G, Menigatti M, Lamberti I, Ponti G, Roncucci L. Aetiology of colorectal cancer and reluctance of monogenetic inheritance. *Gut* 2004 Jan; 53(1): 115-122.
27. Minardi AJ, Sittig KM, Zibari GB, McDonald JC. Colorectal cancer in the young patient. *Am Surg*, 1998 Sep; 64(9): 849-53.
28. Kam MH, Seow-Choen F, Peng XH, Eu KW, Tong CL, Heah SM, Ooi BS. Minilaparotomy left iliac fossa skin crease incision versus midline incision for left sided colorectal cancer. *Techniques in Coloproctology*, 2004, 8, 85-88.
29. Laiyemo AO, Brawley O, Irabor D, Bautall A, Ramesar RS, Madiba TE. Towards colorectal cancer control in Africa. *Int. J. Cancer*. 2016; 138(4): 1033-1034.
30. Katalambula LK, Ntwenya JB, Ngoma T, Buza J, Mpolya E, Mtumwa AH, Petrucks P. Pattern and distribution of colorectal cancer in Tanzania: A retrospective cohort audit at two National hospitals. *J. Canc. Epidemiol.* 2016; 1-12.
31. Metcalf JV, Smith J, Jones R, Oswald RC. Record incidence and causes of rectal bleeding in general practice as detected by colonoscopy. *Br. J. Gen. Pract.* 1996; 46(404): 161-4.
32. Allan E, Nicolaidis C, Helfond M. The evaluation of rectal bleeding in adults. *J. Gen. Int. Med.* 2005; 20; 18-90.
33. Norrelund N, Norrelund H. Colorectal cancer and polyps in patients aged 40 years and over who consult a GP with rectal bleeding. *J. Fam. Pract.* 1996; 13(2): 160-5.
34. Selvachandran SN, Hodder RJ, Ballal MS, Jones P, Cade D. Prediction of colorectal cancer by a patient consultation questionnaire and scoring system: a prospective study. *Lancet* 2002; 360: 278-283.
35. Fijten G, Starmans R, Muris J, Schowten H, Blijham G, Knottnerus JA. Predictive value of signs and symptoms for colorectal cancer in patients with rectal bleeding in general practice. *Family Pract.* 1995; 12: 279-286.
36. Hamilton W, Round A, Sharp D, Peters TJ. Clinical features of colorectal cancer before diagnosis: a population-based case-control study. *Br. J. Cancer* 2005; 93(4): 399-405.
37. MacArthur C, Smith A. Factors associated with speed of diagnosis, referral and treatment in colorectal cancer. 1984. *J. Epidemiol. Community Health*, 38; 122-6.
38. Glass RL, Smith LE, Cochran RC. Subtotal colectomy for obstructing carcinoma of the left colon. *Am. J. Surg.* Vol. 145; Issue 3, March 1983; 335-336.
39. Fazeli MS, Adel MG, Lebaschi AH. Colorectal carcinoma: a retrospective descriptive study of age, gender, stage and differentiation in Iran from 1995 to 2001 as observed in Tehran University. *Dis Colon Rectum* 2007; 50: 990-995.
40. Sule AZ, Mandong BM, Iya D. Malignant colorectal tumors: a ten year review in Jas, Nigeria. *West Afr. J. Med.* 2001. Octo-Dec; 20(4): 251-5.
41. El-Bolkainy TN, Sakr MA, Nouh AA, El-Din NH. A comparative study of rectal and colonic carcinoma: demographic, pathologic and TNM staging analysis. *Journal of the Egyptian National Cancer Institute*, 01 Sep. 2006; 18(3): 258-263.
42. Elesha SO, Owonikoko TK. Colorectal neoplasms: a retrospective study. *East Afr. Med. J.* 1998; 75: 718-723.
43. Walsh JM, Terdiman JP. Colorectal cancer screening: scientific review. *JAMA.* 2003 Mar 12; 289(10): 1288-96.
44. Winawer SJ, Zauber AG, Ho MN, O'Brien MJ, Gottlieb LS, Sternberg SS, Wayne JD, Schapiro M, Bond JH, Panish JF. Prevention of colorectal cancer by colonoscopic polypectomy. The National Polyps study workgroup. *N. Eng. J. Med.* 1993 Dec.30; 329(27): 1977-81.
45. Mandel JS, Bond JH, Church TR. Reducing mortality from colorectal cancer screening for faecal occult blood. *N. Eng. J. Med.* 1993; 328; 1365-1371.

46. Saidi H, Abdihakim M, Nijihia B, Jumba G, Kiarie G, Githaiga J, Abinya NO. Clinical outcomes of colorectal cancer in Kenya. *Ann. Afr. Surg.* 2011 Vol(7); 42-45.
47. Abdulkareem FB, Abudu EK, Awolola NA. Colorectal carcinoma in Lagos and Sagamu, Southwest Nigeria: a histopathological review. *World J. Gastroenterol.* 2008; 14; (42): 6531-5.
48. Meyer JE, Narang T, Schnoll-Sussman FH et al., Increasing incidence of rectal cancer in patients aged younger than 40 years: an analysis of the surveillance, epidemiology and end results database. *Cancer* 2010 Sep. 15 (published online August 23 wileyonlinelibrary.com).
49. Deressa BT, Cihoric N, Tefesse E, Assefa M, Zemenfes D. Multidisciplinary cancer management of colorectal cancer in Tikur Anbessa Specialised Hospital, Ethiopia. *J. Glob. Oncol.* 2019 Oct; 5: 1-7.
50. Samawi HH, Shaheen AA, Tang PA, Heng DYC, Cheung WY, Vickers MM. Risk and predictors of suicide in colorectal cancer patients: a surveillance, epidemiology and end result analysis. *Curr. Oncol.* 2017 Dec; 24(6): e513-e517.
51. Adekunle OO & Ajao GO. "Colorectal cancer in adolescent Nigerians." *Scandinavian Journal of Gastroenterology. Supplement*, 1986. Vol. 21, no.124; pp.183 – 186.
52. Adekunle OO & Lawani JA. "Clinical aspects and management of carcinoma of the rectum in Nigerians." *East African Medical Journal.* 1982. Vol. 59, no.3, pp. 206-213.
53. Aghaji MAC & Obiekwe OM. Sexual function in males following abdominoperineal resection in Nigeria. 1991. *The Central African Journal of Medicine*; 1991. 37; 301-303.
54. Irabor D, Adedeji OA. Colorectal cancer in Nigeria: 40 years on. A review. *European Journal of Cancer Care.* 2009; 18; 110-115.
55. Cunningham D, Atkim W, Lenz HJ, Lynch HT, Minsley B, Nordlinger B, Starling N. Colorectal cancer. *Lancet.* 2010; 375: 1030-1047.
56. Overman MJ, Hoff PM. EGFR-Targeted Therapies in Colorectal cancer. *Dis. Colon & Rectum.* 2007. 50; 1259-1270.
57. Datubo JCB, Naaeder SB, Tettey, Gyosi RK. Colorectal carcinoma: an update of current trends in Accra. *WAJM.* 2010; 29: 178-183.
58. Gondos A, Brenner H, Wabinga H, Parkin DM. Cancer Survival in Kampala, Uganda. *Br. J. Cancer* 2005 9 May; 92(9): 1808-1812.