

**Valorization Of Agro-industrial Bio-waste From
Seed Cotton In The Restoration Of Degraded
Soils In The District Of Korhogo In Northern
Côte d'Ivoire**

ABSTRACT

The objective of this study was to restore degraded soils with organic manure from agro-industrial bio-waste (seed cotton) in the Department of Korhogo in Northern Côte d'Ivoire.

The research plots were set up in a Fisher block experimental design with four treatments, namely T0 (control soil), T1 (bio-waste at 1 month of biodegradation), T2 (bio-waste at 3 months of degradation) and T3 (bio-waste at 6 months of degradation).

On each of the treatments, bio-waste was applied using spreading technique and the variety F8128 of corn was sown to evaluate the agronomic performance of bio-waste. Four agronomic parameters were measured: plant height, collar diameter, ear weight and grain yield. Likewise, the physical, physico-chemical and chemical analyses of the soil were performed before sowing and after corn harvesting.

The work was carried out from 2018 to 2019 in the district of Korhogo in Northern Côte d'Ivoire.

Our study shows significant increase in corn grain yield in each of the treatments (T1 = 2.26 T/ha; T2 = 1.98 T/ha and T3 = 1.48 T/ha) compared to the control (T0 = 1 T/ha). Laboratory analyses of the soil and bio-waste indicate a very low level of organic matter (MO varies from 0.55 to 0.77 g.kg⁻¹) in the soil of the experimental plot and good mineralization of the organic matter in the bio-waste regardless of the decomposition time (C/N varies from 12 to 13). After application of the bio-waste, the organic matter content (T3 = 10.28 g.kg⁻¹; T1 = 23.61 g.kg⁻¹ and T2 = 23.63 g.kg⁻¹) and nitrogen content (T3 = 0.69 g.kg⁻¹; T1 = 1.16 g.kg⁻¹ and T2 = 1.21 g.kg⁻¹) of the various treatments significantly increased compared to the control soil (MO = 0.66 g.kg⁻¹ and N = 0.013 g.kg⁻¹). The pH level increased where the organic manure was applied becoming slightly acidic (5.7 to 6.4) compared to the control that had strong acidic reaction (4.8 to 5.1).

Based on our study, it is evident that organic fertilizer has positive effect on corn yield. The seed cotton bio-waste has had an improving effect on the degraded soil of in Northern Côte d'Ivoire.

Keywords: bio-waste, Côte d'Ivoire, corn, restoration and degraded soil.

1. INTRODUCTION

Land degradation is a phenomenon that leads to a permanent loss of the biological and economic productivity of ecosystems and agro-ecosystems. It concerns all types of environments exploited by human societies to ensure their livelihoods [1]. The soil, which is one of the main bases of agroforestry production in sub-Saharan Africa, faces major degradation problems. This natural resource is particularly affected by the loss of its multiple

21 potentialities [1]. High soil degradation, particularly in sub-Saharan Africa, is linked to
 22 climatic conditions aggravated by high land pressure and inappropriate agricultural practices
 23 (mining agriculture) [2]. Indeed, as soon as litter is cleared and disappears, there is a rapid
 24 decrease in soil organic matter (MOS) and the beginning of chemical, biological and physical
 25 degradation of surface horizons. The fire brutally mineralizes the litters, temporarily
 26 straightens the pH, but releases CO₂ and ash which are blown by the wind or washed away
 27 during the first storms. Plowing in turn introduces oxygen into the soil, accelerates organic
 28 matter mineralization, and mixes the underlying humus and mineral horizons. At the end, ,
 29 cultivated soils are becoming both less productive and less resistant to rainfall energy [3].
 30 Those in the Korhogo district (Northern Côte d'Ivoire) do not remain on the margins of this
 31 very advanced degradation because of their overexploitation, which is linked to population
 32 growth [4].


33 The general objective of this work is to improve the degraded soils with organic manure
 34 (seed cotton) from the waste come from the textile industry in the district of Korhogo.
 35 Specifically, to: (i) evaluate the effect of the organic manure on corn cultivation, a plant that
 36 is widely consumed in Northern Côte d'Ivoire (ii) examine the behavior of this organic
 37 fertilizer on the properties of some degraded soils in the Korhogo district and (iii) establish a
 38 correlation between the decomposition time of cotton fibers and/or seeds (organic manures)
 39 and the properties of degraded soils in relation to corn yield.

41 2. MATERIAL AND METHODS

43 2.1 Study site

44 The work was carried out in the department of Korhogo in northern Côte d'Ivoire. The relief
 45 of the investigated area is monotonous, with altitudes varying, on average, between 300 and
 46 400 m. However, granitic inselbergs sometimes reaching altitudes of more than 500 m exist
 47 in the landscape. The department is made up of lateritic plateaus that can sometimes reach
 48 three meters in height, evidence of old peneplain. These plateaus are affected by a very
 49 slight and regular slope. The average annual rainfall oscillates around 1200 mm of rain. The
 50 area consists of forest galleries, wooded and shrubby savannah vegetation [5]. The
 51 geological formations of the area consist of a succession of bands of shale rocks, migmatite
 52 rocks and plutonic rocks. These are mainly granites, granodiorites, undifferentiated shales
 53 and sericite shales [6], from which several types of soils are derived, namely Ferralsols,
 54 Cambisols, Fluvisols and Luvisols [7]. The chosen area is the site around the village of
 55 Natiokobadara in Korhogo district (Fig.1).

56



57 **Fig. 1. Geographical location of the studied sites**

58

59 **2.2. Material**

60

61 **2.2.1. Organic manure**

62 The organic manure (bio-waste) used is based on seed cotton from the waste from the
63 textile industry in the city of Korhogo. Three types of manure were used depending on the
64 duration of degradation: 1 month (T1), 3 months (T2) and 6 months (T3). In addition, soil
65 samples from horizons with thicknesses ranging from: 0-20 cm; 20-40 cm and 40-60 cm are
66 available (Fig. 2).



67

68 **Fig. 2. Soil and organic manure used on the experimental site**

69 **2.2.2. Plant material**

70 The plant material used for our experiments consists of corn grains of the F8128 variety
71 improved by the Centre National de Recherche Agronomique (CNRA) an agronomic
72 research institution of Côte d'Ivoire [8]. It is a variety with a long cycle of 100 to 120 days
73 (Fig. 3; Table 1).

74

75

76

77

78

79

80

81

82



83 **Fig. 3. Variety of corn coded F8128 used for experimentation**

84

85
86
87

Table 1. Some characteristics of the corn variety F8128 used for the experimentation

Long cycle	Yield t/ha		Particular Characteristic	Seed color	Origin
	Potential	Medium			
100-120 Days	7	3 à 5	Streaks tolerance	Yellow, Horny/toothed	CNRA, 2006

88

89 **2.3 Methods**

90

91 **2.3.1 Morpho-pedological characterization and soil sampling of the experimental site**

92 A morpho-pedological characterization was performed at the experimental site to determine
93 the soil type. Thus, two soil profiles (100 cm x 80 cm x 120 cm) were opened and described,
94 taking care to indicate the date, location, map sheet, vegetation, local slope, topographical
95 position, microrelief, following the "ORSTOM" approach based on the method of [9].

96

97 **2.3.2 Experimental design**

98 The experimental design used is a Fisher blocks. Total experimental site area is 338 m² (26
99 m x 13 m) divided into three blocks separated by 2 m buffer-lanes. The various treatments
100 were installed on elementary plots of 15 m² (5 m x 3 m). They are separated by 2 m buffer-
101 aisles from each other. Four processing operations have been set up and are as follows:

102 T0: Soil Control = no organic fertilizer applied,

103 T1: Bio-waste at 1 month of decomposition,

104 T2: Bio-waste at 3 months of decomposition,

105 T3: Bio-waste at 6 months of decomposition.

106 For each treatment, organic manure (bio-waste) was applied to each unit using the
107 spreading technique before the seedbed was prepared. Thus, according to the technical
108 data sheet of the variety used, 60 kg of manure was added to each elementary plot, i.e. 60
109 kg of manure (bio-waste) per plots (15 m²), which corresponds to 40 T/ha of organic manure
110 [8]. Manual ploughing to a depth of about 15 cm was carried out to bury this manure in the
111 first depths of the soil. Corn growth parameters measured were plant height (cm) and collar
112 diameter (cm). At the production level, the ear weight (kg/ha) and grain yield of corn in (T/ha)
113 were measured.

114

115 **2.3.3. Analysis of soil and organic fertilizer samples**

116 Soil samples were analyzed using standard methods [10] before sowing and after corn
117 harvesting. This is the granulometry, by the Robinson Kölen method, of the organic carbon
118 (C) of the soil determined by titrimetry (Walkley-Black method) after oxidation using a
119 mixture of sulfuric acid (H₂SO₄) and potassium dichromate (K₂Cr₂O₇). Total nitrogen was
120 determined by the Kjeldahl method based on wet oxidation. Total phosphorus was
121 determined by colorimetry after reaction with phosphoric acid in the presence of ammonium
122 molybdate and ascorbic acid. The assimilable phosphorus (Olsen-Dabin method) was
123 extracted using sodium bicarbonate (NaHCO₃) at pH 8.5.

124 The exchangeable bases (K, Ca and Mg) were extracted using ammonium acetate.
125 Potassium was determined using flame photometers, while Ca and Mg were determined by
126 the flame atomic absorption spectrophotometer. The pH (water) was determined using the
127 pH meter after adding 50 ml of ionized water to 20 g of soil followed by agitation and settling.
128 The bulk density (da) was determined by the cylinder method, using undisturbed and fresh
129 samples, knowing the constant dry weight of the samples at 105°C and the volume of the
130 cylinders of the samples used [11].

131

132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159

2.3.4. Statistical analysis of data

The results of the soil, bio-waste and corn agronomic parameters analyses obtained were statistically processed by an analysis of variance (ANOVA 1) with the SPSS version 20 (IBM Corp., USA) software, and in case of significant difference, TUKEY post-ANOVA test was performed to differentiate the groups. The normality of the data and the homogeneity of the variance were checked beforehand using the Kolmogorov-Smirnov and Shapiro-Wilk tests, respectively.


3. RESULTS

3.1 Corn grain yields

Statistical analysis applied to the data collected at the yield outcome level indicates a highly significant difference ($F_{cal} = 39.66^{**}$ and $P_{cal} = 0.00023$). Corn kernel yields vary from 0.99 to 2.45 T/ha depending on the different treatments and the control soil. The results indicate that the corn grain yield recorded at the control soil level is 1 T/ha compared to 2.26 T/ha for T1; 1.98 T/ha for T2 T/ha and 1.48 T/ha for T3 (Fig.4).

3.2 Physical, physico-chemical and chemical characteristics of the soil at the experimental site before sowing

The soil of the experimental site has a brown (10YR6/6) and spotted rusty ochre (7.5YR6/8) coloring, with a depth of 60 cm. The horizons of subangular polyhedral structure and sandy-clayey texture are apparently not humiferous. The drainage class is approximately 5.3. There is an indurated layer at medium depth. The charge in concretions and nodules is greater than 40%. It is a Pisoplinthic Ferralsol in the WRB classification. The soil content low organic matter (0.55 to 0.77 g.kg⁻¹) relatively and its decomposition is very slowly (Table 2). The soil is very active ($IB > 2$) with a high acid pH (4.8 to 5.1) and a low nutrient reserve ($CEC = 3.91$ to 4.85 cmol⁺/kg).



160
161
162

Fig. 4. Different yields (T / ha) in grain of corn depending on the treatments

163
164
165

Table 2. Physical, physicochemical and chemical characteristics of the soil of the experimental site

Analyzed soil parameters		Depth of the horizon (cm)		
		0-20	20-40	40-60
Granulometry (g.kg ⁻¹) (Clay = A; Silt = L; Sand = S; thin = f et Coarse = g)	A	7.01	8.18	9.02
	Lf	17.02	16.27	16.95
	Lg	20.28	19.15	19.75
	Sf	25.11	26.23	23.35
	Sg	30.58	30.17	30.93
Apparent density (g.cm ⁻³)	Da	2.01	2.08	2.11
Porosity (%)	P	24.15	21.51	20.38
Carbon (g.kg ⁻¹)	C	0.32	0.45	0.39
Organic matter (g.kg ⁻¹)	MO	0.55	0.77	0.67
Index battance	IB	3.26	-	-
Potential hydrogen	pH eau	5.1	4.9	4.8
Total nitrogen (g.kg ⁻¹)	N	0.012	0.014	0.015
Ratio Carbon/ nitrogen	C/N	26.67	32.14	26.00
Phosphore assimilable (mg.kg ⁻¹)	P ₂ O ₃ ass	39.8	41.37	40.59
Cation exchange capacity (cmol ⁺ /kg)	CEC	4.85	3.91	4.41
Sum of exchangeable bases (cmol ⁺ /kg)	S	2.73	2.74	2.67

166
167
168
169
170
171
172
173
174
175
176
177
178

3.3 Physico-chemical and chemical characteristics of the bio-waste used

The characteristics of the bio-waste used in our experiments are shown in Table 3. Thus, the nitrogen and carbon contents of bio-waste for the treatments T1 (N = 0.02 g.kg⁻¹; C = 0.26 g.kg⁻¹) and T2 (N = 0.17 g.kg⁻¹; C = 2.04 g.kg⁻¹) are low compared to T3 (N = 1.38 g.kg⁻¹; C = 17.02 g.kg⁻¹). However, the degree of mineralization of organic matter is good for the T2 (C/N = 12) and T3 (C/N = 12.33) treatment compared to the T1 (C/N = 13) treatment which is slightly slow. The iron content is high for T2 (Fe = 40 mg.kg⁻¹) and T3 (Fe = 33.21 mg.kg⁻¹) treatments and low for T1 (Fe = 5.33 mg.kg⁻¹) treatments.

Table 3. Chemical composition of bio-waste (fibers and cotton seeds)

Manures	Parameters											Level of soil mineralization (Ratio C/N)
	N (g.kg ⁻¹)	C (g.kg ⁻¹)	P (mg.kg ⁻¹)	K (mg.kg ⁻¹)	Ca (mg.kg ⁻¹)	Mg (mg.kg ⁻¹)	Fe (mg.kg ⁻¹)	Na (mg.kg ⁻¹)	Mn (mg.kg ⁻¹)	Pb (mg.kg ⁻¹)	Zn (mg.kg ⁻¹)	
1 month (T1)	0,02	0,26	0,12	0,02	0,02	0,03	5,33	0,09	0,11	0,06	0,06	13
3 month (T2)	0,17	2,04	0,96	0,17	0,19	0,28	39,99	0,66	0,81	0,47	0,48	12
6 month (T3)	1,38	17,02	8,03	1,38	1,54	2,30	33,21	5,53	6,71	3,94	4,03	12,33

179
180
181
182
183
184
185
186
187
188
189
190
191
192
193

3.4 Physical, physico-chemical and chemical characteristics of the soils of the different treatments after corn harvesting

The rooting depth of the crop determines the size of the soil compartment that can be exploited by the root system. It is the lower limit of the system considered in the balance sheet method. Thus, after the various experiments, the treated soils sampled from a depth of 30 cm and analyses gave the results indicated in Table 4 below. From granulometry, statistical analyses indicate a significant difference in clay content (F_{cal} = 134.14; P_{cal} = 0.028). There is an enrichment of clay at the treatment level (T1 = 18.18 g.kg⁻¹; T2 = 18.46 g.kg⁻¹ and T3 = 18.89 g.kg⁻¹) than in the control soil (T0 = 8.07 g.kg⁻¹). After application of the bio-waste, there is a decrease in coarse sand at the treatment level (T1 = 23.61 g.kg⁻¹; T2 = 23.63 g.kg⁻¹ and T3 = g.kg⁻¹) compared to the control soil (T0 = 30.56 g.kg⁻¹). The soils under the various treatments had their porosity increased from 41.13% to 58.11% compared to the control soil (24.53%), i.e. a coefficient of variation of 35.6%. The control soils have an average pH value that is strongly acidic (pH water = 4.9). Six months after application of

194 organic manures, the pH values obtained are moderately to slightly acidic (T1 = 6.2, T2 = 6.2
 195 and T3 = 5.9). The CEC values obtained after application of organic manures are relatively
 196 higher (T1 = 8.57 cmol+/kg; T2 = 7.8 cmol+/kg; T3 = 7.08 cmol+/kg) than those recorded at
 197 control soil level (T1 = 4.39 cmol+/kg).

198
 199

Table 4. Characteristics of telltale soil and treatments after agricultural use

Analyzed soil parameters (sampling depth = 30 cm)		Treatments				F _{cal}	P _{cal}	P _{théor}	CV (%)
		T0	T1	T2	T3				
Granulometry (g.kg ⁻¹) (Clay = A; Silt = L; Sand = S; thin = f et Coarse = g)	A	8.07 b	18.18 a	18.46 a	18.89 a	134.14*	0.028	< 0.05	32.7
	Lf	16.75 b	14.07 b	15.33 b	16.34 b	159.12ns	0.065	> 0.05	7.6
	Lg	19.73 b	21.53 b	21.73 b	20.62 b	129.35ns	0.074	> 0.05	5.7
	Sf	24.90 b	22.60 b	20.85 b	26.31 b	84.53ns	0.061	> 0.05	4.2
	Sg	30.56 a	23.61 b	23.63 b	17.84 b	113.01*	0.0134	< 0.05	21.6
Apparent density (g.cm ⁻³)	Da	2 a	1.56 b	1.21 b	1.11 b	122.57*	0.0142	< 0.05	34.8
Porosity (%)	P	24.53 b	41.13 a	54.34 a	58.11 a	145.16*	0.0341	< 0.05	35.6
Carbon (g.kg ⁻¹)	C	0.39 b	8.25 a	9.75 a	10.87 a	534.88**	0.0021	< 0.01	65.4
Organic matter (g.kg ⁻¹)	MO	0.67 b	14.19 a	16.77 a	18.70 a	919.90**	0.0053	< 0.01	67.1
Index battance	IB	2.71 a	0.23 b	0.21 b	0.19 b	816.34**	0.00312	< 0.01	71.5
Potential hydrogen eau	pH eau	4.9 b	6.2 a	6.2 a	5.9 a	6.69**	0.0057	< 0.01	25.71
Total nitrogen (g.kg ⁻¹)	N	0.01 b	1.24 a	1.34 a	1.41 a	913.42**	0.0045	< 0.01	62.2
Ratio Carbon/ nitrogen	C/N	28.29 a	6.65 b	7.28 b	7.71 b	853.02**	0.0027	< 0.01	61.61
Phosphore assimilable (mg.kg ⁻¹)	P ₂ O ₃ a ss	40.59 b	68.35 a	62.24 a	62.33 a	935.39**	0.0032	< 0.01	37.8
Cation exchange capacity (cmol ⁺ /kg)	CEC	4.39 b	8.57 a	7.80 a	7.08 a	179.18**	0.0012	< 0.01	25.43
Sum of exchangeable bases (cmol ⁺ /kg)	S	2.71 b	6.60 a	5.79 a	5.10 a	185.91**	0.0024	< 0.01	27.47

200 The averages followed by the same letter on the same line, are not significantly different at
 201 the Probability threshold < 0.05, according to the method of Tukey, * = significatif, ** = highly
 202 significant, ns = not significatif (F_{cal} = Calculated Fisher, P_{cal} = Calculated Probability, P_{Théo} =
 203 Theoretical Probability), CV = coefficient of variation.

204 3.5 Relationship between the characteristics, physical, chemical and grain 205 yields of corn

206 The various correlation tests carried out between the soil properties of the different
 207 treatments and the grain yield of corn have yielded several levels of significance. Thus, at
 208 the control treatment level (T0) significant correlations are observed between A and CEC (r
 209 = 0.90); P and IB (r = - 0.81) and P and P₂O₃ass (r = - 0.76). The total porosity is therefore
 210 negatively correlated with the beat index and available phosphorus (Table 5). At the T1
 211 processing level, the highest correlations are observed between P and MO (r = 0.94); P and
 212 pH water (r = 0.85); P and C/N (r = 0.95). The grain yield of corn is highly correlated to the
 213 variables P, MO, C/N and CEC with correlation coefficients in the order of: 0.9; 0.90; 0.86
 214 and 0.90 respectively (Table 5). The correlation matrix obtained after linking the data at the
 215 T2 processing level gave significant correlations between the variables: P and IB; P and pH
 216 water; P and C/N with correlation coefficients of 0.97; 0.81 and 0.86 respectively. In the
 217 same treatment, negative and significant correlations exist between: Rend and P; Rend and
 218 N; Rend and P₂O₃ass with r respectively of: -0.88; -0.98 and -0.91 (Table 5). The linking of
 219 different data obtained through the T3 processing gave significant correlations between: IB
 220 and P; IB and N with correlation coefficients of: 0.96; 0.97 and respectively between Rend
 221 and P; Rend and IB; Rend and N; Rend and P₂O₃ass for respectively r coefficients which
 222 are: 0.97; 0.99; 0.97 and 0.92.

223

224
225
226

Table 5. Correlation between soil properties of different treatments and corn grain yields

Treatments	Variable torque (X;Y)	r	r ²	t	Pr (> t)	Significativity
T0	A - CEC	0.9	0.81	5,02	0,0024	**
	P - IB	-0.81	0.66	-3.40	0.0145	*
	P - P _{ass}	-0.76	0.58	-2,86	0,0285	*
T1	P - MO	0.94	0.88	5.43	0.0056	**
	P - pHeau	0.85	0.73	3.28	0.0302	*
	pHeau - MO	0.82	0.67	2.82	0.0474	*
	C/N - P	0.95	0.91	6.29	0.0033	**
	Yields - P	0.91	0.82	4.29	0.0127	*
	Yields - MO	0.9	0.81	4.18	0.0139	*
	Yields - C/N	0.86	0.73	3.3	0.0297	*
	Yields - CEC	0.9	0.81	4.12	0.0145	*
T2	P - IB	0.97	0.94	7.59	0.0016	**
	P - pHeau	0.81	0.66	2.78	0.0516	*
	P - C/N	0.86	0.74	3.39	0.0275	*
	Yields - P	-0.88	0.78	-3.8	0.0192	*
	Yields - N	-0.98	0.95	-8.84	0.0009	**
	Yields - P ₂ O ₃ ass	-0.91	0.83	-4.42	0.0115	*
T3	IB - P	0.96	0.92	6.62	0.0027	**
	IB -N	0.97	0.95	8.37	0.0011	**
	Yields - P	0.97	0.94	8.02	0.0013	**
	Yields -IB	0.99	0.98	20.09	0.0001	**
	Yields - N	0.97	0.95	8.37	0.0011	**
	Yields - P ₂ O ₃ ass	0.92	0.85	4.69	0.0093	**

227
228
229

* = significant. ** = highly significant;

230

4. DISCUSSION

231

The different corn grain yields recorded show a significant difference. This yield is 1 T/ha at the level of the control plot (T0), 1.48 T/ha for plot T3, 2.26 for T1 and 1.98 T/ha for plot T2.

232

233

These results can be explained by the fact that the control soil of our experimental site is rich in aluminum oxide, unlike oxides. However, an aluminum-rich soil creates plant toxicity

234

235

resulting in the slowing down of plant growth, due to aluminum phosphates precipitation in the roots and the blocking of the cellular divisions of the terminal meristems of the roots and,

236

237

possibly, of the aerial buds. Exchangeable aluminum strongly inhibits the activity of soil microflora and particularly blocks the development of rhizobiums [12]. The improved yields of

238

239

the other treatments are explained by the fact that within the organic matter, the aluminum ion is complexed in particular by humic and fulvic acids. Organic matter reduces toxicities by

240

241

complexing ions and therefore aluminum, on the browned soils of Toumodi located in Centre of Côte d'Ivoire [13]. Impacts of this improved soil fertility on maize yields have been

242

243

established. A significant increase in soil fertility and maize yields due to manure is

244

245

observed. The morpho-pedological description of the soil (control, T0) of the experimental site revealed an indurated layer at medium depth. The drainage class is approximately 5.3. Physical

246

247

physico-chemical and chemical analyses of soil samples indicate that the relatively low organic matter (0.55 to 0.77 g.kg⁻¹) decomposes very slowly (C/N = 26.67 to 32.14). The soil

248

249 is very active ($IB > 2$) with a high acid pH (4.8 to 5.1) and a low nutrient reserve ($CEC = 3.91$
250 to $4.85 \text{ cmol}^+/\text{kg}$). The total porosity negatively correlated to the beat index is low (about
251 22%). From results of previous works carried out on degraded soils in Northern region of
252 Côte d'Ivoire the characteristics of soils (Pisoplinthic and Ferralsol) described above in
253 present study would be degraded [4]. This degradation is certainly due to overgrazing and
254 continuous cultivation, which have strongly contributed to reducing vegetation cover and
255 organic matter levels, thus promoting compaction.

256 After the application of organic manure, the clay content in the various treatments T1 (18.18
257 $\text{g}\cdot\text{kg}^{-1}$), T2 ($18.46 \text{ g}\cdot\text{kg}^{-1}$) and T3 ($18.89 \text{ g}\cdot\text{kg}^{-1}$), varied on average by 32.7% compared to
258 the control soil T0 ($8.07 \text{ g}\cdot\text{kg}^{-1}$). Indeed, sufficiently high concentrations of organic matter
259 significantly limits or even stop the soil leaching process [14].

260 The organic matter (OM) and nitrogen (N) content of the various treatments increased by
261 67% and 62.2% respectively compared to the control soil. At the water pH level, the
262 improvement is about 25.71%, because at the control treatment level (T0) the values
263 obtained indicate that the soil is strongly acidic ($\text{pH}=4.8$ to 5.1) while after application of
264 organic manure, the soils have become slightly acidic (5.9 to 6.2). Initially, the C/N ratio that
265 characterizes the state of organic matter mineralization is 28.29 (very slow mineralization).
266 After the application of organic manure, the C/N ratios at the treatment level ranged from
267 6.65 to 7.71.

268 There is a clear improvement in the mineralization of organic matter, especially in the T3
269 treatment. The results of the application of bio-waste mark the beginning of the restoration of
270 this degraded soil. The consequence of a good evolution of organic matter is that humus,
271 flocculating with clay, forms the clay-humic complex, a real reserve and nourishing base of
272 the earth. The formation and accumulation of humus allows the storage of most of the
273 elements (C, O, H, H, N, P, K, S, trace elements) essential to plant life [15].

274 The soils under the various treatments had their porosity increased from 41.13% to 58.11%
275 compared to the control soil ($P = 24.53\%$), i.e. a coefficient of variation of 35.6%. The
276 correlations established between porosity (P) and beat index (IB) are: negative at the control
277 soil level ($r = -0.81$) but positive and significant at the treatment levels (r ranges from 0.96 to
278 0.97). Indeed, humic compounds, by binding to clay, contribute to improving soil porosity and
279 water retention capacity. As a result, the soil is more aerated, less prone to compaction,
280 leaching and erosion by rain and watering, and it stores water better: all factors that make it
281 more fertile. In addition, the structural role of humus induces a good penetration of the soil
282 by the air and therefore oxygenation in depth to stimulate a good biological activity of the
283 macro and micro fauna and flora of the soil. Roots penetrate more easily at depth, which
284 contributes to improving their exploration of the soil and subsoil and thus their supply of
285 water and minerals [16].

286

287 5. CONCLUSION

288 The results of this study show that in the locality of Natiokobadara in Korhogo, the soils
289 investigated have, for the most part, a very low humiferous impregnation and a
290 predominantly sandy-clayey-silt texture, very compact, with a rare or total absence of
291 vegetation, especially on the upper slopes. They are highly acidic ($\text{pH}=4.8$ to 5.1) with very
292 slow mineralization ($C/N=26$ to 32.14) of organic matter. The nutrient supply is very low. Six
293 months after application of bio-waste at a rate of 60 kg per experimental unit or 33.33 tons
294 per hectare, the soils became slightly acidic ($\text{pH}=5.7$ to 6.4) compared to the baseline
295 (strongly acidic). The mineralization of soil organic matter has evolved towards normality
296 because the C/N ratio obtained at the treatment level varies from 6.65 to 7.71. Corn grain
297 yield is more significant at the treatment level (T1 = 2.26 T/ha; T2 = 1.98 T/ha and T3 = 1.48
298 T/ha) than that of the control soil (T0 = 1 T/ha). The seed cotton bio-waste has had an
299 improving effect on the degraded soil in Northern Côte d'Ivoire.

300

301

COMPETING INTERESTS

302

“Authors have declared that no competing interests exist.”.

303

304

REFERENCES

305

1. Bied-Charreton M. The fight against desertification, a global cause. The State of the World, Paris, Discovery; 2007. French.

306

307

2. Ouandaogo N, Ouattara B, Pouya MB, Gnankambary Z, Nacro HB, Sedogo PM. 2016. Effects of organo-mineral fertilization and crop rotations on soil quality. International Journal and Biological and Chemical Sciences. 2016 10 (2): 904-918. French.

308

309

310

311

3. Roose E, Barthès B. Soil carbon erosion and its selectivity at the plot scale in Tropical and Mediterranean regions. In: Roose E, Lal R, Feller C, Barthès BS, editors. “Soil erosion and carbon dynamics”, eds., Advances in Soil Sciences, CRC Press, Boca Raton, Florida; 2006.

312

313

314

315

4. N'Guessan A, Diarrassouba N, Konan A, Nangha KY, Inza JF, Kouamé AY. Indicators of physical soil degradation in northern Côte d'Ivoire: the case of Boundiali and Ferkéssédougou. Africa Science. 2015; 11 (3). French.

316

317

318

5. YT. Climate, socio-economic changes and landscapes in Côte d'Ivoire. Synthesis Brief of Scientific Activities presented for the purpose of obtaining the habilitation to direct research. University of Science and Technology of Lille, France; 2005. French.

319

320

6. Yace I. Initiation to geology. The example of Côte d'Ivoire and West Africa. Petrology, Regional Geology. CEDA, SODEMI; 2002. French.

321

322

323

7. Koné B, Diatta S, Sylvester O, Yoro G, Camara M, Dome DD, Assa A. Estimate of the potential fertility of ferralsols by color. Canadian Journal of Soil Science. 2009; 89: 331-

324

325

342. French.

326

327

8. Akanvou L, Akanvou R, Anguété K, Diarrassouba L. Cultivating maize well in Côte

328

d'Ivoire. National Center for Agronomic Research (CNRA). Corn factsheet; 2006.

329

330

French.

331

9. Boulet R, Chauvel A, Humbel FX, Lucas Y. Structural analysis and cartography in

332

pedology I Taking into account the two-dimensional organization of soil coverage:

333

toposequence studies and their main contributions to knowledge of the soil. Cah.

334

ORSTOM, Ser. Pédol.1982; 19 (4): 309-321. French.

335

10. McKeague JA. 1978. Manual on Soil Sampling and Methods of Analysis. 2nd edition.

336

Canadian Society of Soil Science, Ottawa; 1978.

337

11. Blake RG, Hartage KH. 1986. Bulk density. In: Klute A, edidor. Methods of soils

338

analysis. Part1. 2nd edition. Madison, WI, USA: American Society of Agronomy; 1986.

339

12. Boyer J. Exchangeable aluminum: agronomic implications, evaluation and correction

340

of its toxicity in tropical soils. ORSTOM notebook, Pedological Series. 1976; 19 (4): 259-

341

269. French.

342

13. Yoboué KE. Pedogeochemical and mineralogical studies of the browned soils

343

derived from the formations of the volcanosedimentary complex of anikro and kahankro

344

(Toumodi) in the central south of the Ivory Coast. PhD Thesis Es-Sciences, University of

345

Cocody, Abidjan, Cote d'Ivoire; 2010. French.

346

14. Sauzet O, Cammas C, Gilliot JM, Bajard M, Montagne D. Development of a novel

347

image analysis procedure to quantify biological porosity and illuvial clay in large soil thin

348

sections. Geoderma. 2017 292: 135-148. doi: 10.1016 / j.geoderma.2017.01.004

349

15. Ben HH, Karbout N, Kridan K, Sanaa M, Jedidi N. Characterization of the colloidal

mineral and organic fractions of the surface horizons of the toposequence soils in the

semi-arid zone of Tunisia. Study and Management of Soils. 2012; 19: 105-118. French.

350 16. Le Villo M, Arrouays D, Deslais W, Daroussin J, Le Bissonais Y, Clergeot D. Estimate of
351 the quantities of exogenous organic matter necessary to restore and maintain French loamy
352 soils at a given organic level. Study and Management of Soils. 2001; 8 (1): 47-63. French.

353

354 **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

355 Here is the Definitions section. This is an optional section.

356 **ANOVA:** Analysis of variance

357 **CNRA:** National Center for Agronomic Research

358 **MO:** Organic Matter

359 **Term:** Definition for the term

360

361

UNDER PEER REVIEW