

Original Research Article

PHYSICOCHEMICAL ATTRIBUTES IN MANGO FRUIT (*Mangifera indica*) AS INFLUENCED BY STORAGE TEMPERATURE AND HOT WATER TREATMENT

ABSTRACT

Background: *The temperature being the most important environmental factor that influences the deterioration of perishable commodities. It is often critical that fresh produce rapidly reach the optimal pulp temperature for short term storage if it is to maintain its highest visual quality, flavour, texture and nutritional content (Kader, 2013).*

Aims: *The effects of storage temperature and hot water at various temperature and duration on chemical and textural characteristics of the Keitt mango fruit were evaluated for the 2015/16 growing season in Botswana.*

Materials and method: *The treatments were fruits dipped in distilled water at room temperature (25 ± 2 °C- control), fruits dipped in hot water at 50 and 55 °C for a duration of 3, 5 and 10 minutes, and storage temperatures at 4, 7, 10, 13, or 25 ± 2 °C, plus 95% RH.*

Results: *The results showed that as the storage temperature and water temperature decreased, the proline content and electrolyte leakage increased significantly ($P \leq 0.0001$). The interactions of storage temperature and hot water temperature, and duration in which mango fruit was treated with hot water, significantly ($P \leq 0.01$) maintained vitamin C content, firmness and reduced fruit weight loss during storage and seven days after storage when the fruit was kept at room temperature.*

Conclusion: *Chemical and physical attributes of Keitt mango fruits were significantly improved by the interactions between storage temperature, hot water temperature and duration.*

Keywords: *membrane permeability, electrolyte leakage, chilling injury, electrolyte leakage, chilling injury, textural characteristics, proline content.*

1.0 INTRODUCTION

Mango (*Mangifera indica* L) belongs to the Anacardiaceae family, also known as the cashew family, with about 75 genera and 700 species, mostly tropical with subtropical and temperate species (Nakasone and Paull, 1998; Derese et al., 2017). Mango is one of the world's most popular fruit crops cultivated in the tropical and subtropical climate (FAO, 2004; Lauricella et al., 2017). Fresh mango has an irresistible combination of flavours and textures that brings new excitement to recipes. Mangoes are an excellent source of vitamins A and C and a good source of

fibre. Many studies have shown that free radicals in the living organisms cause oxidative damage to different molecules such as lipids, proteins, nucleic acids and these are involved in the interaction phases of many degenerative diseases. The mango peel extract exhibited good antioxidant activity in different systems and thus may be used in nutraceuticals (Asula *et al.*, 2007; Lourenço *et al.*, 2019). In addition to that, mangoes are best noted for their vibrant flesh colour, juicy texture and sweet flavour along with important nutrient combination for their phytochemicals. Fruits are living tissues and are diverse in morphology, structure, composition and general physiology. Due to high moisture content, active metabolisms, tender nature and rich in nutrients, they are vulnerable to dehydration, physiological disorders, environmental stress, mechanical injury and pathological breakdown therefore usually considered to be highly perishable (Kader, 2005). Therefore, these characteristics limit the storage life of the fruits and vegetables and cause significant deterioration following harvest. Most fruits that originated from the tropical or subtropical regions are chilling sensitive (Gross *et al.*, 2002; Pio *et al.*, 2019). Chilling injury is a storage disorder that occurs at temperatures below the critical threshold but non-freezing temperature (Hardenburg *et al.*, 1986; Mercer and Smittle, 1992; Sharom *et al.*, 1994). The problem limits the use of low storage temperature to manage postharvest ripening because the temperatures that are low enough to delay ripening, decay and senescence may also be damaging to the fruit. Chilling injury is known to significantly change the microstructure of the tissue which in severe cases may lead to tissue breakdown due to failure to carry normal metabolic processes (Han *et al.*, 2006). Various physiological, biochemical alteration and cellular dysfunction occur in chilling sensitive species in response to chilling stress (Wang, 1982). These alterations include increased membrane permeability and alteration of activities of membrane proteins. Temperature management plays a critical role in ensuring that high-quality mangoes. Avoiding high temperature and reducing temperature to the optimum reduces the rate of physiological and biochemical changes that occur in mango after harvest, minimizes water loss from the fruit and slows the growth of decay-causing micro-organism like anthracnose (Brencht and Cecilia, 2012). However, there is a limit to the low temperature that mango can tolerate due to their susceptibility to chilling injury, a disorder that results in flavour loss, surface blemishes and inhibition of ripening. Heat treatment induces heat shock proteins which protect the product from both heat and chilling injury, suppresses oxidative activity and maintain membrane stability. High levels of reducing sugars and proline were also found to correlate

positively with the resistance to chilling injury (Purvis, 1981). The high levels of proline could be associated with the heat shock proteins which assist in protection against stresses by controlling the proper folding and conformation of the cell membrane and enzymatic proteins. Since low temperature can alter the solubility and folding properties of many proteins, this chaperone activity plays an important role for protection against chilling injury (Vierling, 1991; Saltveit, 2005; Sevillano *et al.*, 2009). In addition to that, mango shelf-life is reduced by high temperature and as such, there is a need to be conscious with the temperature at which mango is being stored to prolong its postharvest shelf-life and maintain its quality. Postharvest losses resulting from chilling injury are higher than have been recognized. This dilemma results in tremendous postharvest losses for chilling injury-sensitive crops (tropical crops). It is, therefore, important to alleviate chilling injury in tropical crops but without compromising the chemical and textural characteristics of the fruit hence the significance of the current study. The objective of the study is to evaluate the effects of storage temperature and hot water treatment and duration on the chemical and textural characteristics of the mango fruit.

2.0 MATERIALS AND METHODS

2.1 Experimental site

A laboratory experiment was conducted from February and May 2015 at Botswana University of Agricultural and Natural Resources, Sebele. Sebele lies about 10km from the centre of Gaborone city on latitude 24°34'S and longitude 25°57'E elevated at 994 m above sea level.

2.2 Experimental design

A factorial experiment ($5 \times 3 \times 3$) laid down in randomized complete design was used with three replications. The treatments were storage temperature, hot water at different temperatures and time of exposure to hot water. The mango fruits were dipped in distilled water at room temperature (25 °C), mango fruits dipped in hot water at 50 and 55 °C; duration (time) in water treatments were 3, 5 and 10 minutes, and storage temperatures at 4, 7, 10, 13 and 25 ± 2 °C. Mango fruits dipped in water at various temperatures and durations were then stored in the temperatures indicated above. In each storage temperature, there were 135 fruits. Mango fruits were packed in perforated paper board cartons for each treatment. The mango cultivar Keitt was used for the study. The fruits were at physiological maturity with the flesh yellow in colour but peel still green and hard.

2.3 Variable assessment

Dependent variables analyzed were vitamin C (ascorbic acid), fruit weight loss, fruit firmness, electrolyte leakage and proline content. The above variables were determined both immediately after removal from cold storage and after seven days' storage at ambient temperature (25 ± 2 °C).

2.3.1. Fruit weight loss

Nine mango fruits were weighed before and after storage to calculate the per cent of the fresh weight loss. This was determined by subtracting the actual average weights of the fruits in each replication. The formula below was used to calculate per cent weight loss.

$$\% \text{ WL} = (\text{IAW} - \text{AAW}) \times 100/\text{IAW}$$

Where WL stands for weight loss, IAW for Initial average weight and AAW for actual average weight after storage

2.3.2 Fruit firmness

Mango fruit flesh firmness was determined using a hand-held Effegi penetrometer (Alfonsine, Italy) with 8 mm tip diameter. Nine fruit per replicate were used. Measurements were taken from two opposite sides of the fruit (red and green) with skin removed.

2.3.3 Electrolyte leakage

Electrolyte leakage was determined according to the method of Chan *et al.* (1985). Five discs taken with a 10 mm diameter cork borer from the peel and pulp tissue, then sample tissues were rinsed with deionized water to eliminate the electrolyte at the cut surface. The samples were placed in a flask containing 25 ml of 0.4 M mannitol. Incubation for 30 minutes at 25 °C was done where the electrical conductivity was measured in a suspending solution with an EC meter as an initial reading. The samples in a flask were heated at 98 °C for 15 minutes and the electrical conductivity was re-measured after cooling. Membrane permeability was calculated using the formula given below:

$$\% \text{ Electrolyte leakage} = \text{IL}_{\text{initial}} \times 100/\text{IL}_{\text{final}}$$

Where $\text{IL}_{\text{initial}}$ stands for initial ion leakage reading at initial temperature, and IL_{final} for final ion leakage reading at the final temperature.

2.3.4 Proline

Mango fruit pulp was cut and homogenized in 10 ml of 3% aqueous sulfosalicylic acid. The homogenate was filtered through Whatman filter paper (grade 1). Then 2 ml of the filtrate was reacted with 2 ml acid-ninhydrin and 2 ml glacial acetic acid in a test tube for an hour at 100 °C in a water bath to develop the colours. Soon after removal from the water bath, the test tube was cooled in an ice bath and proline extracted with 4 ml toluene, mixed vigorously with a test tube stirrer for 15-20 seconds. The chromophore containing toluene was aspirated from the aqueous phase, warmed to room temperature and the absorbance was read in a UV 160 IPC spectrophotometer (Parnomex Inc. New Dehli) at 520 nm using toluene as a blank. Proline content in mango fruit pulp was determined using the formula given below (Bates *et al.*, 1973):

$$\mu\text{mole proline/g of fresh weight} = (\mu\text{g proline/ml} \times \text{ml toluene} / 115.5 \mu\text{g/mole}) / (\text{g sample})$$

2.4 Data analysis

The data collected were subjected to analysis of variance (ANOVA) using the Statistical Analysis System (SAS). Treatment means were separated using the Least Significant Difference (LSD) at P=0.05.

3.0 RESULTS AND DISCUSSIONS

3.1 Vitamin C content

In the current study, storage temperature and water temperature interaction significantly ($P \leq 0.0001$) affected the vitamin C content of the mango fruits (Table 1 and 2), as the storage temperature and hot water temperature increased, the vitamin C content decreased significantly ($P \leq 0.0001$). Also as the storage temperature and duration in which mango fruit was held in hot water increased the vitamin C content significantly ($P \leq 0.0001$) decreased (Table 1). Loss of vitamin C can occur by irreversible conversion of dehydroascorbic acid to 2,3-dioxo-L-gluconic acid, which is then further metabolized. The reaction is pH-dependent, being slow in acid pH, rapid at neutral pH and extremely rapid at alkaline pH (Kays *et al.*, 2004). The decrease in vitamin C content with increase in storage and the water temperature was attributed to decrease in juice pH and titrable acidity observed in the current study since the loss in vitamin C is pH and temperature-dependent, having rapid loss of vitamin C at higher fruit pH and temperature.

Storage of fruits and vegetables at temperatures below 5 °C is reported to decrease the loss of vitamin C (Hardenburg *et al.*, 1986; Kays *et al.*, 2004). The rate of loss of vitamin C is higher with higher storage temperature, an effect associated with loss of acidity (Kays *et al.*, 2004). Mangoes stored at 24-26 °C for 6 days has been reported to result in a drop in vitamin C content from the initial 71 mg/100g to 63.9 mg/100g (Kays *et al.*, 2004). Lee and Kader (2000) reported that the loss of vitamin C in fresh commodities is enhanced by extended storage and high temperature. Yousef *et al.* (2012) reported that ascorbic acid content decreased gradually and significantly during storage at 8, 10, 13 °C as well as in mango fruits dipped in hot water at 48 and 52 °C for 10 minutes.

3.2 Proline content

There was a significant ($P \leq 0.0001$) effect of the storage temperature and water temperature interaction on mango proline content immediately after removal from cold storage and seven days after cold storage on fruit held at room temperature (Table 1 and 2). As the storage temperature and water temperature decreased, the proline content increased significantly ($P \leq 0.0001$). Mangoes dipped in water at 25 °C and stored at 4 °C, had the highest proline content of 25 $\mu\text{mole/g}$, while those dipped at 55 °C but stored at 13 °C had significantly ($P \leq 0.0049$) lower proline content of 0.23 $\mu\text{mole/g}$. As storage temperature decreased from 25 to 4 °C, irrespective of water temperature, the proline content significantly ($P \leq 0.0001$) increased from 0.71 to 18.82 $\mu\text{mole/g}$, respectively, accounting for the 23.7-fold increase in proline content. Mango fruits stored in the non-chilling temperatures of 13 and 25 °C and dipped in hot water at 50 or 55 °C had the lowest proline content in the range of 0.23-0.93 $\mu\text{mole/g}$. The high proline content in mango fruit stored at 4, 7 or 10 °C was attributed to chilling injury caused by the low temperatures. The high accumulation of proline in mango fruit stored below 13 °C in the current study was to enhance chilling injury tolerance to 'Keitt' mango fruits (Shang *et al.*, 2011; Li *et al.*, 2014). High proline content in chilled mango fruit in the current study was also attributed to either enhanced protein degradation at chilling temperatures and/or proline synthesis (Kumar *et al.*, 2003; Shang *et al.*, 2011; Cao *et al.*, 2012; Li *et al.*, 2014). As the hot water temperature increased from 25 °C to either 50 or 55 °C, the proline content decreased from 9.89 $\mu\text{mol/g}$ to either 5.87 or 4.32 $\mu\text{mol/g}$, respectively. The decrease in proline content at high water

temperatures higher than 25 ° C was attributed to by degradation of proline by the high temperatures (Shang *et al.*, 2011; Cao *et al.*, 2012).

Table 1.0 Effects of storage and hot water temperature, and hot water duration on mango chemical and textural characteristics immediately after removal from the storage regimes.

Treatments	Chemical and textural attributes				
	Vit C (mg/100 g)	Proline (μmole /g)	EL (%)	Firmness (N)	Weight loss (%)
Storage temperature °C					
4	39.1 ^a	16.82 ^a	57.08 ^a	67.34 ^a	10.57 ^e
7	32.76 ^b	12.31 ^b	48.77 ^b	45.91 ^b	12.49 ^d
10	28.31 ^{cc}	5.51 ^c	44.95 ^c	42.84 ^c	13.44 ^c
13	23.63 ^{dd}	1.13 ^d	35.31 ^d	34.69 ^d	14.80 ^b
25	21.31 ^d	0.72 ^d	21.36 ^e	34.16 ^d	15.68 ^a
Significance	**	**	**	**	**
Hot water °C					
25	24.27 ^{aa}	11.67 ^{aa}	55.12 ^a	45.62 ^a	14.22 ^a
50	23.23 ^a	11.08 ^a	47.67 ^b	41.43 ^b	14.96 ^b
55	21.35 ^b	10.89 ^{aa}	40.44 ^c	35.26 ^c	15.20 ^c
Significance	**	**	**	**	**
Hot water duration(minutes)					
3 minutes	23.19 ^a	8.05 ^{aa}	42.92 ^a	46.39 ^{aa}	13.11 ^c
5 minutes	22.98 ^a	7.11 ^{aa}	41.36 ^b	45.84 ^a	13.34 ^b
10 minutes	22.67 ^a	6.74 ^a	40.19 ^c	42.74 ^b	13.74 ^a
Significance	ns	ns	**	**	**
Interactions					
Storage temperature × hot water	**	**	**	**	**
Storage temperature × hot water duration	**	ns	**	ns	**
Hot water × hot water duration	ns	ns	ns	**	**
Storage temperature × hot water × hot water duration	ns	ns	*	*	**

** Highly significant at p<0.01, * significant at p<0.05 and ^{ns} non-significant at p>0.05. Means separated using Least Significant Difference (LSD) Test at p≤0.05, Means within columns followed by the same letters are not significantly different.

3.3 Fruit weight loss

There was a significant ($P \leq 0.0001$) effect on mango fruit weight loss due to interaction between the storage temperature and water temperature, storage temperature and water treatment duration as well as water temperature and water treatment duration (Table 1). As the storage temperature and water temperature increased, the percentage weight loss of the fruits increased significantly ($P \leq 0.0001$). Also, the interaction of storage temperature and water treatment duration significantly ($P \leq 0.0001$) increased the mango fruit weight loss. As the storage temperature increased from 4 to 25 °C and water treatment duration increased from 3 to 10 minutes, the percentage weight loss of the fruits significantly ($P \leq 0.001$) increased. The higher fresh weight loss of mango fruit at high storage and water temperatures, and longer duration in hot water at 50 or 55 °C and their interactions was attributed to higher evapotranspiration rate and respiration rate at the higher temperatures. Kumah *et al.* (2011) reported that there was a gradual increase in the cumulative weight loss in 'Keitt' mango fruit after the 4th day of storage and continued with the rapid increase in weight until 21 days after storage. Similarly to this finding, 'Keitt' mango fruits in the present study treated in hot water for 52 °C for 5 minutes, 50 °C for 5 and 10 minutes, and 48 °C for 10 minutes showed a rapid increase in fruit weight loss in comparison to control fruit. According to Kumah *et al.* (2011), the sharp rise in cumulative weight loss was due to high temperature and low relative humidity. Yousef *et al.* (2012) reported a progressive increase in fresh mass loss of mango fruit cultivar 'Copania' throughout the storage period and hot water treatments at 46 or 50 °C for 10 minutes. The results of the current study are further in line with Perez *et al.* (2004) who found that, in avocado fruits, the mass losses were 4.3% at 20 °C for 8 days and 3% at 10 °C for 22 days. A significant effect of storage ($P \leq 0.05$) on Dusheri cultivar of mango was observed and had an increase of weight loss of 36% after 15 days of storage (Rathore *et al.*, 2007). Roongruangsri *et al.* (2013) reported that the percentage of weight loss increased and moisture content of the peel decreased in two tangerine cultivars at a higher temperature and longer duration of storage. Low-temperature storage at 5 °C, reduced the losses of tangerine fruit weight loss and moisture content better than at 25 °C storage (Roongruangsri *et al.*, 2013). Wang (1993) found that weight loss was most severe in squash stored at 15 °C than squash kept at 5 °C. This weight loss was attributed to rapid ripening and senescence due to high temperature (Wang, 1993).

Table 2.0 Effects of storage and hot water temperature, and hot water duration on mango chemical and textural characteristics 7 days after storage at ambient temperature after immediate removal from different treatment temperatures.

Treatment	Vit C (mg/100 g)	Proline (μ mole /g)	EL (%)	Firmness (N)
	Storage temperature °C			
4	28.89 ^a	14.88 ^a	59.98 ^a	36.56 ^a
7	24.78 ^b	9.46 ^b	51.29 ^b	33.81 ^b
10	21.51 ^{cc}	1.47 ^c	46.75 ^c	28.38 ^{cc}
13	21.49 ^c	0.90 ^{dd}	42.58 ^d	27.48 ^c
25	18.09 ^d	0.69 ^d	27.09 ^e	23.08 ^d
Significance	**	**	**	**
Hot water °C				
25	20.36 ^a	9.02 ^a	52.87 ^a	35.88 ^a
50	18.01 ^b	4.46 ^b	44.33 ^b	28.60 ^b
55	17.21 ^{bb}	2.95 ^c	39.41 ^c	25.10 ^c
Significance	**	**	**	**
Hot water duration(minutes)				
3 minutes	18.41 ^{aa}	6.25 ^a	47.76 ^a	32.28 ^a
5 minutes	18.02 ^a	5.53 ^b	45.89 ^b	29.80 ^b
10 minutes	17.96 ^{aa}	4.65 ^c	42.96 ^c	27.50 ^c
Significance	ns	**	**	**
Interactions				
Storage temperature × hot water	**	**	**	**
Storage temperature × hot water duration	ns	**	ns	ns
Hot water × hot water duration	ns	ns	ns	**
Storage temperature × hot water × hot water duration	ns	*	ns	*

** Highly significant at $p < 0.01$, * significant at $p < 0.05$ and ^{ns} non-significant at $p > 0.05$. Means separated using Least Significant Difference (LSD) Test at $p \leq 0.05$, Means within columns followed by the same letters are not significantly different.

3.4 Electrolyte leakage

Membrane damage can be measured by the ion leakage, which in the present study the electrolyte leakage was significantly ($P \leq 0.0001$) higher in mango fruit stored in temperatures of 4, 7 or 10 °C than fruit stored at 13 or 25 °C. Also, fruit treated with hot water at 50 or 55 °C, irrespective of storage temperature, had lower electrolyte leakage and chilling injury incidence and severity than fruit treated with water at 25 °C. As the storage temperature and water temperature increased, the mango electrolyte leakage decreased on fruits assessed immediately after removal from storage temperatures and those that were stored at room temperature for seven days after removal. Fruits dipped in water at 25 °C and stored at 4 °C had the highest electrolyte leakage while those fruits treated with hot water at 50 and 55 °C and stored at 13 and 25 °C had the lowest electrolyte leakage, respectively. These results indicate the role of hot water treatment for up to 10 minutes in maintaining membrane integrity. As the duration in which mango fruit was held in hot water at 50 or 55 °C increased, the electrolyte leakage significantly ($P \leq 0.0001$) decreased in the current study. Junmatong *et al.* (2012) reported that the pulp and skin electrolyte leakage in 'Nam Dok Mai' mango cultivar commonly grown in Thailand, increased during storage at 5 °C and rapidly increased when fruits were transferred to room temperature. The increase in electrolyte leakage occurred before chilling injury symptoms (Junmatong *et al.*, 2012). The results of the current study suggests that oxidative stress is an early response of mango fruits to chilling injury as it initiates membrane degradation causing lipid peroxidation. These findings are consistent with those reported by several investigators who reported that electrolyte leakage increased in Tommy Atkins, Wacheng, Zill, Tainong and Nam DOK Mai mango fruit cultivars during storage at 2-7 °C for 7-30 days (Junmatong *et al.*, 2012; Shewfelt and del Rosario, 2000; González-Aguilar *et al.*, 2000; Zhao *et al.*, 2006; Ding *et al.*, 2007; Wang *et al.*, 2008). Similarly, Zhao *et al.* (2009) reported that the electrolyte leakage intensity reflected chilling injury development phase and a degree in tomato fruit. Although the electrolyte leakage increased with decrease in storage temperature from 25 to 4 °C, electrolyte leakage decreased in mango fruits treated with hot water for a longer duration of 10 minutes. It is also suggested that the reduction in the electrolyte leakage and chilling injury induced by hot water treatment in the current study was attributed to the role of hot water at 50 or 55 °C for a maximum duration of 10 minutes in maintaining cell membrane integrity and reducing lipid peroxidation of cell membranes.

3.5 Fruit firmness

The interaction between the storage temperature and water temperature, as well as interaction between the water temperature and water treatment duration significantly ($P \leq 0.0001$) influenced the fruit firmness immediately after removal from storage and seven days after storage at room temperature (Tables 1 and 2). As the storage temperature and the water temperature increased, fruit firmness decreased significantly ($P \leq 0.0001$) immediately after cold storage and seven days later at room temperature. There was also a significant ($P \leq 0.0016$) interaction between storage temperature and water treatment duration on mango firmness seven days after storage at room temperature (Table 2). As the storage temperature and water treatment duration increased, the fruit firmness significantly ($P \leq 0.0001$) decreased. Water temperature and duration of hot water treatment interaction had a significant ($P \leq 0.0005$) effect on the mango fruit firmness (Table 1 and 2). As the water temperature and duration of hot water treatment increased, the fruit firmness decreased significantly ($P \leq 0.0001$) immediately after cold storage and seven days after storage at room temperature. The decrease in mango fruit firmness caused by the interactions of storage temperature and water temperature, storage temperature and duration in which mango fruit was dipped, and hot water temperature and duration in which mango fruit was dipped was attributed to the role of temperature in fruit texture changes during ripening. The ripening phenomenon in fruits is associated with loss of firmness. Yousef *et al.* (2012) reported that the firmness of mango fruits showed a gradual and significant reduction during storage at 8, 10, 13 °C for four weeks compared with untreated fruits (control). However, storage at 8 °C, was more effective in keeping the fruits firmer after 28 days (Yousef *et al.*, 2012). They further reported a decline in mango fruit firmness in fruits stored at 10 °C after dipping in hot water at 48 and 52 °C for 10 minutes. In the current study, as the water temperature increased from 25 to 55 °C, the fruit firmness decreased significantly ($P \leq 0.0001$) from 46.5 N to 30.9 N after 28 days of storage. As storage temperature increased from 4 °C to 25 °C mango fruit firmness decreased from 45.9 N to 30.2 N, while fruit stored at 4°C but dipped in water at 25 °C or 55 °C had a firmness of 57.8 N and 38.1 N, respectively. This finding is consistent with that of Tian *et al.* (1996) who reported that increasing storage temperatures from 8 to 13 °C significantly decreased the firmness of mango fruits. Heat air treatment of mangoes at 38 °C has been reported to accelerate the softening of fruits due to changes in pectic components and activities of polygalacturonase, pectin methylesterase, and β -galactosidase in ‘Nam Dokmai’

mango fruit during storage at 25 °C (Ketsa *et al.*, 1998). Zhang *et al.* (2012) reported that the firmness in non-cold stored mango fruit declined rapidly from 74.8 n to 12.3 N after five days of storage at 20 °C. Jacobi and Gille (1997) suggested that the decrease in Kengston mango fruit firmness following heat treatment was attributed to an increase in activity of the enzymes pectin methylesterase, polygalacturonase, galactosidase and β -1, 4-gluconase.

4.0 CONCLUSION

This study showed that a combination of low-temperature storage in the range of 7-10 °C and hot water treatment of mango fruits at 50 or 55 °C for 10 minutes was effective in maintaining the chemical and physical attributes of Keitt mango during storage and seven days after storage when fruit was kept at room temperature. Chemical and physical attributes of Keitt mango fruits were significantly improved by the interactions between storage temperature, hot water temperature and duration.

Ethic: NA

Consent: NA

REFERENCES

1. AOAC. (1995). Official Methods of Analysis (15th ed). Association of Official Analytical Chemists, Washington, D.C.
2. Asula, C. M., Naidu, K. A., Rao, P. and Bhat S. G. (2007). Bioactive compounds and anti-oxidants potential of mango peel extract. 105: 982-988
3. Brecht J.K. and Cecilia, M. (2012). Time- temperature combination that Induce chilling injury of mangoes. Final Report: Center for food distribution and retailing, University of Florida, 20p.
4. Cao, S., Cai, Y., Yang, Z. and Zheng, Y. (2012). MeJA induces chilling tolerance in loquat fruit by regulating proline and γ -aminobutyric acid contents. Food Chemistry, 133: 1466-1470.
5. Chan, H.T.JR., Sanxter, S., and Couey, H.M. (1985). Electrolyte leakage and ethylene production induced by chilling injury of papayas. HortScience, 20: 1070-1072.
6. Ding, Z. S., Tian, S. P., Zheng, X. L., Zhou, Z. W. and Smith, Y. (2007). Response of reactive oxygen metabolism and quality in mango fruit to exogenous oxalic acid or salicylic acid under chilling temperature stress. Planta, 214: 895-901.
7. FAO (2004). FAO statistics retrieved from <http://www.fao.org>. 06/08/2015
8. González-Aguilar, G. A., Fortiz, J., Cruz, R., Baez, R., and Wang, C. Y. (2000). Methyl jasmonate reduces chilling injury and maintains postharvest quality of mango fruit. Journal of Agricultural and Food Chemistry, 48: 515-519.

9. Gross, K. C., Wang, C. Y and Saitveit, M. (2002). (eds). 2002. The commercial storage of fruits, vegetables and florist and nursery crops. <http://www.ba.ars.usda.gov/hb66/index.html>. 06/08/2015
10. Hardenburg, R. E., Watada, A. E. and Wang, C.W. (1986). The Commercial Storage of Fruits, Vegetables, and Florist and Nursery Stocks. U. S. Department of Agriculture, Agriculture Handbook 66: 1-133.
11. Jacobi, K.K. and Giles J. E. (1997). Quality of Kengstone mango (*Mangifera indica*) fruit following combined vapour heat disinfestation and hot water disease control treatment. *Postharvest Biol and Technol*, 12: 285-292.
12. Junmatong, C., Uthaibutra, J., Boonyakiat, D., Faiyue, B. and Saengnil, K. (2012). Reduction of chilling injury of 'Nam Dok Mai No.4' mango fruit by treatments with salicylic acid and methyl jasmonate. *Journal of Agricultural Science*, 4(10): 126-136.
13. Kader, A. A. (2005). Increasing food availability by reducing postharvest losses of fresh produce. *Acta Horticulturae*, 682: 2169 – 2175.
14. Kays, S. J. and Paull, R. E. (2004). *Postharvest Biology*. Published by Exon Press Athen, GA, 568p.
15. Ketsa, S., Chidtragool, S., Klein, J. D. and Lurie, S. (1998). Effect of heat treatment on changes in softening, pectic substances and activities of polygalacturonase, pectinesterase and β -galactosidases of ripening mango. *Journal of Plant Physiology*, 153: 457-461.
16. Kumah, P., Appiah, F. and Opoku-Debrah, J. K. (2011). Effect of hot water treatment on quality and shelf life of Keitt mango. *Agriculture and Biology Journal of North America*, 2(5): 806-817.
17. Kumar, S. G., Reddy, A. M., and Sudhakar, C. (2003). NaCl effects on proline metabolism in two high yielding genotypes of mulberry (*Morus alba* L.) with contrasting salt tolerance. *Plant Science*, 165: 1245-1251.
18. Lee, S. K. and Kader, A. A. (2000). Postharvest and Postharvest factors influencing vitamin C content of horticultural crops. *Postharvest Biology and Technology*, 20: 207-220.
19. Li, P., Zheng, X., Liu, Y. and Zhu, Y. (2014). Pre-storage application of oxalic acid alleviates chilling injury in mango fruit by modulating proline metabolism and energy status under chilling stress. *Food Chemistry*, 142: 72-78.
20. Mercer, M. D. and Smittle, D. A. (1992). Storage atmospheres influence chilling injury and chilling injury-induced changes in cell wall polysaccharides of cucumber. *Journal of the American Society for Horticultural Science*, 117(6): 930-933.
21. Nakasone H. Y and Paull R. E. (1998). *Tropical Fruits*, CAB international. United Kingdom.
22. Purvis, A. C. (1981). Free proline in peel of grapefruit and resistance to chilling injury during cold storage. *HortScience*, 16: 160-161.
23. Rathore, A. H., Masud, T., Shehla, S. and Soomro, A. H. (2007). Effect of storage on physio-chemical composition and sensory properties of mango (*Mangifera indica* L.) variety Dosehari. *Pakistan Journal of Nutrition*, 6(2): 143-148.

24. Roongruangsri, W., Rattanapanone, N. and Leksawasdi, N. (2013). Influence of storage condition on physio-chemical and biochemical of two tangerine cultivars. *Journal of Agricultural Science*, 5(2)
25. Saltveit, M. E. (2005). Influence of heat shocks on the kinetics of chilling induced ions leakage from tomato pericarp discs. *Postharvest Biology and Technology*, 36:87-92.
26. Sevillano, L., Scanchez-Baliesta, M.T., Romojano, F and Flores, F.B. (2009). Physiological hormonal and molecular mechanism's regulating chilling injury in horticultural species. *Postharvest technology applied to reduce its impact. Journal of the Science of food and Agriculture*. 89: 555-573
27. Shang, H. T., Cao, S. F., Yang, Z. F., Cai, Y. T. and Zheng, Y. H. (2003). Effect of exogenous γ -aminobutyric acid treatment on proline accumulation and chilling injury in peach fruit after long-term cold storage. *Journal of Agriculture and Food Chemistry*, 59: 1264-1268.
28. Sharom, M., Willemot, C. and Thompson, J. E. (1994). Chilling injury induces lipid phase changes in membranes of tomato fruit. *Plant Physiology*, 105: 305-308.
29. Shewfelt, R. L. and del Rosario, B. A. (2000). The role of lipid peroxidation in storage disorders of fresh fruits and vegetable. *HortScience*, 35: 575-579.
30. Tian, M. S., Woolf, A. B., Bowen, J. H. and Ferguson, I. B. (1996). Changes in colour and chlorophyll fluorescence of broccoli florets following hot water treatment. *Journal of the American Society for Horticultural Science*, 121(2): 310-313.
31. Vierling, E. (1991). The role of heat shock proteins in plants. *Annual review. Plant physiology Plant Molecular Biology*, 42:579-620.
32. Wang, C. Y. (1982). Physiological and biochemical responses of plants to chilling stress. *HortScience*, 17: 173-186.
33. Wang, C.Y. (1993). Approaches to reduce chilling injury of fruits and vegetables. *Horticultural Reviews*, 15: 63-95
34. Wang, L., Chen, S., Kong, W., Li, S. and Archbold, D. D. (2006). Salicylic acid pretreatment alleviates chilling injury and affects the antioxidant system and heat shock proteins of peaches during cold storage. *Postharvest Biology and Technology*, 41: 244-251.
35. Yousef, A. R. M., Emam, H. S. and Ahmed, D. M. M. (2012). Storage and hot water treatments on poststorage quality of mango fruit (*Mangifera indica* L.) variety Copania. *Australian Journal of Basic and Applied Sciences*, 6(13): 490-496.
36. Zhang, Z., Gao, Z., Li, M., Hu, M., Gao, H., Yang, D. and Yang, B. (2012). Hot water treatment maintains normal ripening and cell wall metabolism in mango (*Mangifera indica* L) fruit. *HortScience*, 47(10): 1466-1471.
37. Zhao, Z., Jiang, W., Cao, J., Zhao, Y., Gu, Y. (2006). Effect of cold-shock treatment on chilling injury in mango (*Mangifera indica* L cv 'Wacheng') fruit. *Journal of the Science of Food and Agriculture*, 86: 2458-2462.
38. Derese, S., Guantai, E. M., Souaibou, Y., & Kuete, V. (2017). *Mangifera indica* L.(Anacardiaceae). In *Medicinal Spices and Vegetables from Africa* (pp. 451-483). Academic Press.

39. Lauricella, M., Emanuele, S., Calvaruso, G., Giuliano, M., & D'Anneo, A. (2017). Multifaceted health benefits of *Mangifera indica* L.(Mango): the inestimable value of orchards recently planted in Sicilian rural areas. *Nutrients*, 9(5), 525.
40. Lourenço, S. C., Moldão-Martins, M., & Alves, V. D. (2019). Antioxidants of natural plant origins: From sources to food industry applications. *Molecules*, 24(22), 4132.
41. Pio, R., Souza, F. B. M. D., Kalcsits, L., Bisi, R. B., & Farias, D. D. H. (2019). Advances in the production of temperate fruits in the tropics. *Acta Scientiarum. Agronomy*, 41.

UNDER PEER REVIEW