

**Analysis of Soil Quality in a Deltaic Hydrocarbon Polluted Environment
Niger Delta Nigeria**

Abstract

Aim: The study was aimed to ascertain the damage done by an oil spill on the soils of Ogbia area in Bayelsa State, Niger Delta, Nigeria.

Study design: The study adopted the experimental research design which entailed the use of field measurements and a control site. Three communities which are Elebele, Imiringi and Otuasega were used for the study. At each of the sample locations, three random spots were augered at two depth-levels (Top Sample (T), 0 – 15cm; Bottom Sample (B), 15- 30 cm), with the aid of an auger to collect the samples for laboratory analysis. The parameters of interest to the study are TPH, THC, Organic Matter, THF, THB, pH, sand, silt clay and soil texture and these were analysed using standard techniques as recommended by DPR.

Results: The study revealed that there is a noticeable effect of oil exploitation activities on soil quality within the study area, In the case of THC, the result revealed that the level of THC in the sampled communities was higher than that of the non spill site, hence the presence of hydrocarbon which has caused the pollution of the soil. Organic matter content of the soil also reveals that the non-polluted site has more organic content than the sampled communities.

Conclusion: The study revealed that there is a statistically significant difference in soil quality of the selected communities and that of the non-spilt site. The study, therefore, recommended complete and total remediation of the soils in the area, as this will enhance the soil for increased food production.

Keywords: soil, hydrocarbon, polluted, deltaic, environment

Introduction

Over the years there have been a rising demand for oil and the dependence on oil revenue has greatly increased. This increase has brought an increase in the economic development of Nigeria, but the situation has led to an increase in environmental, degradation, pollution and contamination of the soils in the Niger Delta. This has arisen from the breakdown of oil pipelines and wells as well as petroleum distribution processes. Oil spills on land have brought about a

35 reduction in soil fertility which has affected food production (Sparrow & Sparrow, 1988; Racine,
36 1993).

37 While the benefit of the oil industry is not in doubt, it's explorative, developmental and
38 production processes generate a lot of waste amongst which are of "drilling cuttings, drilling
39 fluids, produced water, sludge, completion and workover fluids, trace metals, heat waste and
40 oxides of carbon, Sulphur and nitrogen" (Nwilo & Badejo, 2005). Fluids and cuttings arising
41 from oil drilling are the largest waste sources during exploration. It is speculated that at least two
42 barrels of fluids and cuttings are generated per foot of a typical well drilled (Nwilo & Badejo,
43 2005).

44 Several studies have revealed that oil pollution affects the quality of soils; in the study conducted
45 by Bada and Olarinre (2012), revealed that leaves had more heavy metal content in plants than
46 stems and roots.

47 Furthermore, Oyem and Oyem (2013), in Ugborodo community on oil spillage impacts on the
48 soil physico-chemical properties. Arising from oil spillage, it was observed that among the four
49 sampled communities parameters measured indicated high amount of hydrocarbon, hence a turn
50 out low fertility, which will bring about low food production and its attendant effects on the
51 sources of livelihood available to the people.

52 Ugboma, (2014) in a separate study on showed that soil physical and chemical characteristics
53 were affected by oil spill which resulted in soil fertility decrease and crop productivity decline as
54 well. This, therefore, provides evidence that oil spill affects soil quality which in turn affects the
55 quantity of food production in an area.

56 The record of oil exploitation activities can be likened to the record of oil poison. This is
57 because as oil exploration and exploitation commenced the next that happened "almost
58 immediately was the three major causes of oil pollution namely; the impact of the seismic
59 survey, gas flaring and oil spills" (Pyagbara, 2007). The consequences of oil exploitation on the
60 environment with an emphasis on the soil quality informed the need for this study.

61 Through a large number of literature research, the harm of oil-polluted soil mainly includes the
62 following aspects: Firstly, because of the small density, higher viscosity and lower emulsifying
63 ability of petroleum, it is easy to be absorbed in soil surface, affecting the permeability and

64 porosity of soil (Wang, 2009; He et al, 1999); petroleum is rich in carbon and a small number of
65 nitrogen compounds, so it can change the composition and structure of soil organic matter and
66 impact the C/N, C/P, salinity, pH, EH and conductivity of soil (Li et al, 2009).

67 The heavy metals (nickel and vanadium) in oil mixtures (Saadat et al, 2014) and high
68 concentrations of salt in oilfield output water can also damage the soil environment (Efsun et al,
69 2015). Secondly, microorganisms in the natural environment are quite abundant in healthy and
70 clean soil. In normal situation, the microorganisms which can resist the oil pollution stress are
71 not developed, while in contaminated soil, in order to adapt to this kind of environment, they can
72 produce certain enzyme system and gradually form a dominant population with symbiotic or
73 synergy effect (Chiara et al, 2009).

74 A number of studies have shown that the hydrocarbon pollution can change the microbial
75 population, the composition of the community structure and the enzyme system in soil, given
76 priority to the inhibitory action (Deng, 2014; Uzoije and Agunwamba et al, 2009). Thirdly, it can
77 impede the normal growth of crops such as reduce the germination rate and fertility and decline
78 the resistance to pests and diseases (Xu and Lu, 2010; Zhu, 2010; Shan et al, 2014). In addition,
79 the oil compounds could react with inorganic nitrogen and phosphorus, limiting the nitrification
80 and removal of phosphoric acid, so the effective nitrogen and phosphorus in the soil would
81 decrease and the absorption of crops will be affected (Liao et al, 2015; Pinchin et al, 2013; Shen,
82 2011).

83 Moreover, the polycyclic aromatic hydrocarbons in petroleum chemicals have carcinogenic,
84 mutagenic, teratogenic and other toxic effects. It can enter into the bodies of people and animals
85 through breathing, skin contact and diet, degrading the normal function of livers and kidney etc,
86 therefore causing a great threat to human's health. At last, the oil pollutants in the soil not only
87 impact the pedosphere, but also the atmosphere and water sphere.

88 **Materials and Method**

89 *The Study Area*


90 The area of study is Imiringi, Otuasega and Elebele communities which are located within the
91 Kolo Creek, which is located 4°47'0"N – 6°25'0"E within the lower Niger Delta. (See fig 1.1).

92 Ogbia is one amongst other local government areas in Bayelsa State, with the Head Quarters
93 situated in the town called Ogbia which is located on the South of the area within Latitude
94 $4^{\circ}45'00''\text{N } 6^{\circ}39'00''\text{E}$. It covers a total area of 695 km^2 . Kolo Creek oil and gas field as called by
95 SPDC is located in Imiringi town. The name Kolo creek as called by SPDC, and is “named after
96 Kolo Creek in Ogbia Local Government Area of Bayelsa State. The area is located about 10km
97 North-West of Yenagoa, which is the capital city of Bayelsa State. It is characterized by tropical
98 rain forest and fresh water swamps that are usually flooded in the raining season”.

99

100 The study was conducted in the Kolo Creek area as called by Shell Development Company of
101 Nigeria (SPDC) which is basically made up of three communities in Ogbia; there are Imiringi,
102 Otuasega having and Elebele communities which were selected for the study because of the
103 presence of oil wells in the communities. The study adopted the use of the experimental research
104 design which enabled the researchers achieve the purpose of the study.

105 Since the interest of the study is on soil quality, soil samples were collected using a systematic
106 sampling pattern (Tel and Hagarty, 1984). At each of the sample locations, three random spots
107 were augered at two depth-levels (Top Sample (T), 0 – 15cm; Bottom Sample (B), 15- 30 cm),
108 with the aid of an auger (EGASPIN, 2002). Also, at each of the sample locations and soil depth
109 levels (T or B), the soil samples were bulked together to give a composite sample. The soil
110 samples from different sample locations and soil depth levels were on each occasion collected in
111 polythene bags and labeled accordingly and sent to the laboratory for analysis. Two hypotheses
112 were also put forth to guide this study.



113

114 **Fig 1: Ogbia Area showing sampled communities**

115

116 *Analytical Procedure for Soil Quality Analysis*

117 *Total Heterotrophic Bacteria*

118 **A sample of water with a mass of 1 gram** is weighed into 9ml sterile diluents (0.85% NaCl)
 119 under aseptic condition. It is shaken vigorously to homogenize and serially diluted. Then 0.1ml
 120 aliquot of the inoculums is collected using a sterile pipette, inoculated on Nutrient Agar (NA)
 121 surface. The inoculums are spread evenly with sterile hockey stick (Bent rod). **Plates are**
 122 **incubated at 28^oC for 24 hours. Thereafter,** colonies are counted to obtain colony forming unit
 123 (cfu) value per ml of water sample. Distinct colonies are picked and streaked or subculture on
 124 freshly prepared Nutrient Agar medium to obtain pure culture after 24 hours incubation at 37^oC.
 125 The pure culture is gram stained for microscopic examination. It is also used to carry out
 126 biochemical tests for characterization and identification of the isolates.

127

128

129 **Total Petroleum Hydrocarbon**

130 This analysis shall be done using gas chromatography/flame ionization detector method.
131 Methods 3580A and 3550C describe the procedure used for extracting non-volatile and semi-
132 volatile organic compounds from solids such as soils, sludge and wastes, while method 8000
133 describes the determinative steps employed in the GC/FID of sample extracts obtained using the
134 methods described above.

136 **Soil pH Determination**

137 The ph of soil was done with the aid of a pH meter.

139 **Results**

140 The results of the laboratory analysis of the soil samples collected from the selected communities
141 are presented in the table below.

142 **Table 1: Result of Physico-Chemical/Microbiological Analysis of Soil**

Parameters	Imiringi	Elebele	Otuasega
TPH(mg/kg)	178.77	1,732.50	132.51
THC(mg/kg)	277.2	2246.0	223.7
Organic matter (%)	7.440	14.890	11.280
THF(cfu/g)x10 ³	1.2	0.2	0.2
THB(cfu/g)x10 ³	3.2	1.0	1.2
P ^H	6.30	6.50	5.30
Sand (%)	11.2	12.7	11.9
Silt (%)	31.6	33.4	32.8
Clay (%)	57.2	53.9	55.3
PSD/Texture	Silty clay	Silty clay	Silty clay

143 Source: Researchers field work (2015)

144

145

146

147

148 **Table 2: Comparism of soil quality Parameters of polluted sites with Non polluted soil.**

Parameters	Imiringi	Elebele	Otuasega	Non polluted site
TPH(mg/kg)	178.77	1,732.50	132.51	20.18
THC(mg/kg)	277.2	2246.0	223.7	36.0
Organic matter (%)	7.440	14.890	11.280	18.20
THF(cfu/g)x10 ³	1.2	0.2	0.2	0.5
THB(cfu/g)x10 ³	3.2	1.0	1.2	2.2
p ^H	6.30	6.50	5.30	7.10
Sand (%)	11.2	12.7	11.9	13.8
Silt (%)	31.6	33.4	32.8	35.2
Clay (%)	57.2	53.9	55.3	51
PSD/Texture	Silty clay	Silty clay	Silty clay	Silty clay

149 **Source Researchers field work (2018)**

150 **Discussions**

151 As seen in table 1, Total Petroleum Hydrocarbon in soils of the selected communities ranges
 152 between 132.51 and 1,732.50, with Otuasega community accounting for the lowest which is
 153 132.51mg/kg and Elebele community accounting for 1,732.50mg/kg which is the highest,
 154 although the result shows that total petroleum hydrocarbon value differs significantly from one
 155 location to another.

156 Total Hydrocarbon Content in soils of the selected communities ranges between 223.7mg/kg and
 157 2246.0mg/kg, with Otuasega community accounting for the lowest which is 223.7mg/kg and
 158 Elebele community accounting for 2246mg/kg which is the highest; although the result shows
 159 that total hydrocarbon content value in soils differs significantly from one location to another.

160 Organic matter in soils of the selected communities ranges between 7.440 and 14.890, with
 161 Elebele community accounting for the highest, with an organic content value of 14.890. This was
 162 followed by Otuasega community with an organic content value of 11.280 and Imiringi
 163 community with an organic content value of 7.440.

164 Total Heterotrophic Fungi in soils of the selected communities ranges between 0.2 and 1.2, with
 165 Imiringi community accounting for the highest, with a THF value of 1.2. The two other sampled
 166 community's Otuasega and Elebele had the same THF value of 0.2 each.

167 Total Heterotrophic Bacteria in soils of the selected communities ranges between 1.0 and 3.2,
168 with Imiringi community accounting for the highest, with a THB value of 3.2. This was followed
169 by Otuasega community with a THB value of 1.2 and Elebele community with a THB value of
170 1.2.

171 pH in soils of the selected communities ranges between 5.30 and 6.50, with Elebele community
172 accounting for the highest, with a pH value of 6.50. This was followed by Imiringi community
173 with a pH value of 6.30 and Imiringi community with a pH value of 5.30.

174 Sand level in soils of the selected communities ranges between 11.2% and 12.7% with Elebele
175 community accounting for the highest, with sand % value of 12.7%. This was followed by
176 Otuasega community with sand % value of 11.9% and Imiringi community with sand % value of
177 11.2%. Silt as observed to range between 31.6 and 33.4. The highest was recorded in Elebele with
178 a percentage value of 33.4% silt; Otuasega had 32.8% silt and Imiringi 31.6% silt.

179 Clay ranged between 53.9 and 57.2 with the highest percentage value of 57.2% recorded in
180 Imiringi. This was followed by Otuasega with clay percentage value of 55.3% and Elebele with a
181 clay percentage value of 53.9%. The texture of the soil as shown revealed that the three
182 community soils are silty clay soil.

183 Table 2 revealed that total petroleum hydrocarbon level in the three sampled communities is
184 above that of the non polluted site with Elebele community having more concentration of total
185 petroleum hydrocarbon than the other communities. The result here shows that there is a
186 noticeable effect of oil exploitation activities of soil quality within the study area.

187 In the case of THC, the result revealed that the level of THC in the sampled communities was
188 higher than that of the non-spill site, hence the presence of hydrocarbon which has caused the
189 pollution of the soil.

190 The organic matter content of the soil also reveals that the non-polluted site has more organic
191 content than the selected communities. This implies that the presence of oil in the soil due to
192 pollution has affected the organic content of the soil hence bringing a reduction in its amount
193 when compared with the non-polluted site.

194

195 *Hypothesis Testing*

196 *Table 3: Chi-Square analysis for the difference in soil quality between Imiringi (polluted site)*
 197 *and non-polluted site*

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
178.77	20.18	158.9	25150.79	1246.3
277.2	36.0	241.2	58177.4	1616.0
7.440	18.20	-10.76	115.78	6.36
1.2	0.5	0.7	0.49	0.98
3.2	2.2	1	1	0.45
6.30	7.10	-0.8	0.64	0.090
11.2	13.8	-2.6	45.69	3.31
31.6	35.2	-3.6	12.96	0.368
57.2	51	8.2	67.24	1.32
				$X^2 = 2875.12$

198 $df = (9-1) (2-1)$

199 $(9-1) (2-1)$

200 $8 \times 1 = 8$

201 Therefore we conclude that X^2 calculated value is 2875.12 while the critical value at 10 degree
 202 of freedom and 95% significant level is 15.51.

203 The calculated X^2 calculated value 2875.12 is greater than the critical value of 15.51 we,
 204 therefore, reject the null hypothesis which states that there is no statistically significant
 205 difference in soil quality of Imiringi and that of the non polluted site and accepts the alternate
 206 hypothesis which states that there is a statistically significant difference in soil quality in Imiringi
 207 and that of the non polluted site.

208

209

210

211

212 *Table 4:Chi-Square analysis for difference in soil quality between Elebele (polluted site) and*
 213 *non-polluted site*

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
1,732.50	20.18	1712.32	2932.01	145.3
2246.0	36.0	2210	4884100	135669.4
14.890	18.20	-3.31	10.96	0.602
0.2	0.5	-0.3	0.09	0.18
1.0	2.2	-1.2	1.44	0.65
6.50	7.10	-0.6	0.36	0.05
12.7	13.8	-1.1	1.21	0.088
33.4	35.2	-1.8	3.24	0.092
53.9	51	2.9	8.41	0.165
				X² = 135,816.5

214 $df = (9-1) (2-1)$

215 $(9-1) (2-1)$

216 $8 \times 1 = 8$

217 Therefore we conclude that X² calculated value is 135,816.5 while the critical value at 10 degree
 218 of freedom and 95% significant level is 15.51.

219 The calculated X² calculated value 135,816.5 is greater than the critical value of 15.51 we
 220 therefore reject the null hypothesis which states that there is no statistically significant difference
 221 in soil quality of Elebele and that of the non polluted site and accept the alternate hypothesis
 222 which states that there is a statistically significant difference in soil quality in Elebele and that of
 223 the non polluted site.

224

225

226

227

228

229

230 **Table 5: Chi-Square analysis for the difference in soil quality between Otuasega (polluted site)**
 231 **and non-polluted site**

Observed	Expected	O-E	(O-E) ²	$\frac{(O-E)^2}{E}$
132.51	20.18	112.33	12618.03	625.3
223.7	36.0	187.7	35231.3	978.65
11.280	18.20	-6.92	47.89	2.63
0.2	0.5	-0.3	0.09	0.18
1.2	2.2	-1	1	0.45
5.30	7.10	-1.8	3.24	0.456
11.9	13.8	-1.9	3.61	0.263
32.8	35.2	-2.4	5.76	0.164
55.3	51	4.3	18.49	0.363
				X² = 1608.456

232 $df = (9-1) (2-1)$

233 $(9-1) (2-1)$

234 $8 \times 1 = 8$

235 Therefore we conclude that X² calculated value is 1608.456 while the critical value at 10 degree
 236 of freedom and 95% significant level is 15.51.

237 The calculated X² calculated value 1608.456 is greater than the critical value of 15.51 we
 238 therefore reject the null hypothesis which states that there is no statistically significant difference
 239 in soil quality of Otuasega and that of the non polluted site and accept the alternate hypothesis
 240 which states that there is a statistically significant difference in soil quality in Otuasega and that
 241 of the non polluted site.

242 **Hypothesis Two**

243 The second hypothesis of the study, states that there is no statistically significant variation in the
 244 extent of soil damage amongst sampled communities in Ogbia.

245 The table below shows the variation in the extent of soil damage amongst sampled communities
 246 in Ogbia.

247

248

249 **Table 6: One way (ANOVA) for variation in soil quality in sampled communities in Ogbia**

ANOVA

<i>Source of Variation</i>	<i>SS</i>	<i>Df</i>	<i>MS</i>	<i>F</i>	<i>P-value</i>	<i>F crit</i>
Between Groups	948295.4	2	474147.7	1.804708	0.186144	3.402826
Within Groups	6305476	24	262728.2			
Total	7253771	26				

250

251 From the table above, it is revealed that calculated F value for the analysis is 1.804708 while the
 252 critical value is 3.402826. Since the calculated F value of 1.804708 is less than the critical value
 253 of 3.402826 at F^2_{26} degree of freedom, the implication of this is that the null hypothesis H_0 of no
 254 significant difference is accepted while rejecting the alternate H_1 which states that there is a
 255 statistically significant variation in the extent of soil damage amongst selected communities.
 256 Arising from the above it is pertinent to state that the study has revealed that there is no
 257 statistically significant variation in soil quality of the selected communities.

258

259 **Conclusion**

260 The study concludes that the level of damage done by oil spill on the soils and its spatial
 261 variation in Ogbia, the study therefore revealed that total petroleum hydrocarbon level in the
 262 three sampled communities is above that of the non polluted site with Elebele community having
 263 more concentration of total petroleum hydrocarbon than the other communities. The result here
 264 shows that there is a noticeable effect of oil exploitation activities on soil quality within the study
 265 area. For THC, the result revealed that the level of THC in the sampled communities was higher
 266 than that of the non spill site, hence the presence of hydrocarbon which have caused the pollution
 267 of the soil. The organic matter content of the soil also reveals that the non polluted site has more
 268 organic content than the sampled communities. This implies that the presence of oil in the soil
 269 due to pollution has affected the organic content of the soil hence bringing a reduction in its
 270 amount when compared with the no polluted site.

271 Statistically, the study revealed that there is a statistically significant difference in soil quality of
272 the sampled communities and that of the non spilled site, but on specific parameter, it was found
273 that total petroleum hydrocarbon and total hydrocarbon content were higher in the selected
274 communities than the non oil spilled site. The study further revealed that;

275 1. There is a noticeable effect of oil exploitation activities on soil quality within the study
276 area.

277 2. There is also a noticeable decline in the crop production arising from soil pollution
278 occasioned by oil exploitation.

279 3. There is a clear cut evidence of the state of the soil in the polluted sites as compared to
280 the non polluted which is an indication of the state of the soil having been ravaged by the
281 detrimental effect of oil exploratory activities in the area.

282 4. Total petroleum hydrocarbon level in the three sampled communities is above that of the
283 non polluted site with Elebele community having more concentration of total petroleum
284 hydrocarbon than the other communities.

285 5. The study revealed that there is a statistically significant difference in soil quality of the
286 selected communities and that of the non spilled site

287 On the basis of the above, the study recommended a complete and total remediation of the soils
288 in the area, as this will enhance the soil for increased food production.

289 ***Informed consent:*** Before the commencement of the research the communities were duly
290 informed of the research which they obliged and provided a guard to assist the researchers during
291 the period

292

293 ***Competing interests:*** there are no conflicting interests in this work

294

295

296 **References**

297 Bada B.S and Olarinre T.A (2012) Characteristics of Soils and Heavy Metal Content
298 of Vegetation in Oil Spill Impacted Land in Nigeria. Proceedings of the
299 Annual International Conference on Soils, Sediments, Water and Energy,

300 Vol 17 Article 2. http://scholarworks.umass.edu/soils_proceedings.
301
302 Chiara A, Rosario M, Flavia T, (2009) Bioremediation of diesel oil in a co-contaminated soil by
303 bioaugmentation with a microbial formula tailored with native strains elected for heavy
304 metals resistance. *Science of the Total Environment*, 407(8): 3024-3032.
305
306 Deng R Y. (2014) Microbiological monitoring and evaluation of compound pollution of
307 petroleum and heavy metal in slated soils. Shandong: Shan Dong University, China
308
309
310 Efsun D F, Olcay, Hüseyin S. (2015) Variations of soil enzyme activities in petroleum-
311 hydrocarbon contaminated soil *International Biodeterioration & Biodegradation*: 268-275
312
313 He L J, Wei D Z, Zhang W Q. (1999) Research of microbial treatment of petroleum
314 contaminated soil. *Advances in Environmental Science*, 7(3): 110-111 (in Chinese)
315
316
317 Li L, Liu M, Zhao J L. (2011) Current status and prospects of microbial remediation of soil
318 pollution in oil fields in northern Shaanxi. *Chinese Journal of Soil Science*. 2011,
319 42(4):1011-1013.
320
321 Nwilo, P. C, Badejo, O. T. (2005), 'Oil Spills Problem and Management in the Niger Delta
322 *International Oil Spills, Conference Monitoring*, p .2.
323
324 Oyem, Isama Lawrence Rank and Oyem, Isama Lawrence (2013) Effects of Crude
325 Oil Spillage on Soil Physico-Chemical Properties in Ugborodo Community.
326 *International Journal of Modern Engineering Research (IJMER)*
327 www.ijmer.com Vol. 3, Issue. 6, Nov - Dec. 2013 pp-3336-3342 ISSN: 2249-
6645 www.ijmer.com
328 Pyagbara, L.S. (2007). The adverse impacts of oil pollution on the environment and wellbeing of
329 a local indigenous community: The experience of the Ogoni. New York: Department
330 of Economic and social Affairs, United Nations.
331 Racine, Ch. H. (1993). Long-term recovery of vegetation on two experimental crude
332 oil spills in interior Alaska black spruce taiga. *Canadian Journal of Botany* 72:
333 1171 – 1177.
334
335
336 Saadat S, Mirkhani R, Mohebi A, (2014) Study on phytoremediation of soils polluted with heavy
337 metals and oil pollutants in agricultural lands affected by Persian Gulf War (Khouzestan,
338 fars, kohgiluyeh & boyrahmad and boushehr provinces).
339
340 Shan B Q, Zhang Y T, Cao Q L, (2014) Growth responses of six leguminous plants adaptable in

341 Northern Shaanxi to petroleum contaminated soil. *Environmental Science*, (35):1125-
342 1130
343
344
345 Sparrow, S. D. and Sparrow, E. B. (1988). Microbial biomass and activity in a subarctic
346 soil ten years after crude oil spills. *Journal of Environmental Quality* 17: 304 –
347 309.
348
349 Ugboma, P. P (2014) Effects of Oil Spillage on Soil Fertility in Udu Local Government
350 Area in Delta State. *International Journal of science and Technology, Ethiopia*,
351 Vol. 3(3), S/No 8, September, 2014: 47-56 ISSN: 2225-8590 (Print) ISSN 2227-
352 5452 (Online) DOI: <http://dx.doi.org/10.4314/stech.v3i3.5>
353 Uzoije A P and Agunwamba J C. (2011) Physiochemical properties of soil in relation to varying
354 rates of crude oil pollution. *Journal of Environmental Science and Technology.*, 4(3):
355 313-323.
356
357 Wang B. (2014) Mechanism of growth and physiological response of three arbor species to oil
358 contamination. Shaanxi: Northwest A&F University.
359
360 Zhu H, Liu K, Yang X, (2013) Sedimentary controls on the sequence stratigraphic architecture in
361 intracratonic basins: an example from the Lower Permian Shanxi Formation, Ordos
362 Basin, Northern China. *Marine and Petroleum Geology*. 45: 42-54
363
364 Xu Y H and Lu M.(2010) Bioremediation of crude oil-contaminated soil: Comparison of
365 different biostimulation and bioaugmentation treatments. *Hazardous Materials*, 183: 398-
366 401.
367