

THE ETHANOLIC EXTRACTS OF THE ROOT, STEM AND LEAVES OF *BALANITES AEGYPTIACA* AGAINST SELECTED MICROBES

ABSTRACT

Balanites aegyptiaca has been used in many traditional treatments of microbial infections. This work was carried out with the aim of determining the phytochemical compounds present in the ethanolic extracts of the root, stem and leaves of *Balanites aegyptiaca* and their effects against selected microorganisms. Phytochemical screening was undertaken to determine presence of secondary metabolites in the stem-bark, roots and leaves of the test plant. Well diffusion method was used to determine sensitivity of test organisms to the test plant. Result indicated the presence of alkaloids, tannins, flavonoids, saponins, steroids, and anthraquinones, although anthraquinones were absent in ethanolic extracts of root and stem bark of *Balanites aegyptiaca* but present in the ethanolic extract of the leaf. Antibacterial effects were tested against *Staphylococcus aureus* and *Klebsiella pneumoniae*. Results obtained from the ethanolic extract of the root stem and leaves showed significant zones of inhibition against *Klebsiella pneumoniae* and *Staphylococcus aureus* at all concentrations of 100mg/ml, 50mg/ml, 25mg/ml, 12.5mg/ml and 6.125mg/ml. The positive control performed better than the extracts in terms of zone of inhibition. The results of the minimum inhibitory concentration (MIC) indicated that the leaf ethanolic extract had the greatest activity against *K. pneumoniae* with MIC value of 12.5mg/ml, while the results from the minimum bactericidal concentration (MBC) showed that the leaf ethanolic extract had the greatest activity against *K. pneumoniae* with an MBC value of 12.5mg/ml.

Key words: *Balanites aegyptiaca*, phytochemical compounds, microorganisms, plant extracts

1. INTRODUCTION

The antimicrobial activities of plant oils and extracts have formed the basis of many applications, including raw and processed food preservation, pharmaceuticals, alternative medicine and natural therapies. Moreover, the increasing use of plant extracts in the food, cosmetics and pharmaceutical industries suggests that in order to find active compounds, a systematic study of medicinal plants is very important

The plant (*Balanites aegyptiaca* Delile) has long history of traditional uses for wide ranges of diseases [1].

The kernel oil is used for the treatment of wounds in Nigeria [2]. Also, in Nigeria, a mixture of dried leaves powder of *Balanites aegyptiaca* Delile and *Ricinus communis* in water is used for contraception [3]

35 [4]. Traditional medicine has remained the most affordable and easily accessible source of treatment in
36 the primary health care system of resource poor communities and local therapy is the only means of
37 medical treatment for such communities [5]. According to Alaribe [6] about 80% of Nigerian homes,
38 maintain some sort of private family traditional medicine practitioner. Existing data and contemporary
39 researchers seem to authenticate the assumption for general health improvement of the masses by
40 traditional healers. Plants have broader uses than just food and genetic reservoirs. Medicinal plants
41 have been used for centuries to treat a wide variety of ailments [7]. The presence of secondary
42 metabolites in plants has been associated in most of their therapeutic activities [8]. Herbal medicines are
43 now considered a part of Complementary and Alternative Medicine (CAM) and are gaining popularity due
44 to their potent antioxidant activity, minimal side effects and economic viability [9]. Many research efforts
45 have been directed towards the provision of empirical proofs to back up the use of plants species in trade
46 and medicinal practices in recent years [10]. This work intends to study specific microorganisms'
47 response to extracts of *B. aegyptiaca*

48 **2. MATERIALS AND METHODS**

49 **Source and preparation of plant materials**

50 The plant materials were collected from Gombe State University botanical garden Biological Science
51 Department; Gombe State University, it located in Gombe between the latitude 10°00N to 10°20N and
52 longitude 11°0E to 11° E. These were brought and identified by a taxonomist with voucher number
53 900191 at the Herbarium unit of the Department of Biological Sciences, Gombe State University. The
54 plant parts were air-dried for two weeks at room temperature (25°C) in the laboratory and then ground to
55 powder.

56 **2.1 Extraction procedures**

57 The ground plant parts were extracted at the Department of Pharmacognosy, Faculty of
58 Pharmaceutical Sciences, Gombe State University, following the methods of Sofowora [11].

59 **2.2 Preparation of Ethanolic Extraction of *Balanites aegyptiaca***

60 Approximately 800 g of the dried leaves and roots of *B. aegyptiaca* were extracted with 10 litres of 80%
61 (v/v) ethanol by maceration at (25°C) for 3days. The total mixture then was strained and filtered. The
62 filtrate was concentrated to dryness on a water bath at 100° C so as to obtain the dry extract which was
63 stored at -20°C for further studies.

64

65

66 **2.3 Phytochemical analysis**

67 The method of Sofowora [12] was employed for the test of the presence of the phytochemical properties.

68 **2.4 Source and preparation of test microorganisms**

69 The stock cultures of the test microorganisms were obtained from the Department of Microbiology,
70 Gombe State University. Their validity was determined by sub culturing onto nutrient agar and confirmed
71 by standard cultural, morphological and biochemical techniques as described by [13]. The inocula of the
72 test organisms were standardized by the method of Bauer and Thornsberry [14]. This was done by
73 suspending each test organism in 5ml of nutrient broth and the turbidity was compared with that of 0.5
74 McFarland standard. McFarland standard was prepared by adding 0.6ml of 1% barium chloride (BaCl₂)
75 to 99.4ml of 1% sulphuric acid (H₂SO₄) solution. The turbidity of the 0.5 McFarl and standards was used
76 for estimation the number of bacteria in broth culture (culture for 24 hours at 37°C) to pour into 5ml of
77 distilled water in order to obtain a standard bacterial suspension of 1 x 10⁵ cfu/ml [15].

78 **2.4 Preparation of concentration of extracts**

79 Approximately, **one**g of each extract was dissolved in 5mls of distilled water to yield 200mg/ml. **One** ml of
80 the 200mg/ml was taken and added to 1ml of distilled water to give a concentration of 100mg/ml. **One** ml
81 of the 100mg/ml extract concentration was also taken and added to 1ml of distilled water to get a
82 concentration of 50mg/ml. The procedure was repeated twice to give concentrations of 25mg/ml and
83 12.5mg/ml.

84 **2.6 Antibacterial susceptibility testing**

85 The antibacterial activity of the fractions of *B. aegyptiaca* was determined using the well method (Kirby-
86 Bauer Methods) as described by Abalaka [16]. Standard aseptic microbiological methods were followed
87 throughout this antibacterial study.

88

89 **2.7 Well method for antibacterial activity**

90 The well method was employed to assay the plant fractions for antibacterial activity. Petri dishes were
91 poured with nutrient agar and allowed for 30 minutes to solidify (This was done in duplicate for each
92 fraction and tested organism). The tested organisms were then inoculated by spreading on the inocula on
93 the surface of the medium using a sterile swab stick. A sterile Cork borer (size 3) was used to bore 4
94 wells in the medium. The different concentration of the plant fractions were placed in the wells using a
95 sterile syringe and needle (different for each sample and tested organism). These were then allowed a
96 diffusion time of 1 hour after which the plates were incubated at 37 °C for 24 hours. The positive control
97 was Ceftriaxone (100mg/ml). The potency of the extracts was determined by the clear zones of inhibition
98 around the wells and was respectively measured as the diameter zones of inhibition. The MIC was
99 determined using the method of Doughari [17], while MBC was determined using the method of Rotimi
100 [18].

101 **2.8 Data analysis**

102 One way Analysis of Variance (ANOVA) of was used to assess the efficacy of the plant parts in terms
103 of the activity as was shown by the zones of inhibition. Since there was no significant difference,
104 Duncan's multiple range test (DMRT) and Student t-test was not carried.

105 **2.9 Result and Discussion**

106 The result of preliminary phytochemical screening (Table 1) revealed the presence of alkaloids,
107 flavonoids, tannins, saponins and steroids in the roots and stem bark, while anthraquinones were present
108 in the leaf only. This is similar to that of Adebayo and Ishola [19] who investigated the activity of root,
109 stem-bark and leaves extracts of *Terminalia glaucescens*' against some pathogenic organisms.

110 Phytochemical screening of the fractions of *Terminalia glaucescens*' also revealed the presence of
111 alkaloids, tannins, saponins, steroids, flavonoids, anthraquinones and phlobatannins (mostly in root and
112 stem-bark). Also, Igbiosa [20] in the screening of the methanolic and water extracts of the stem bark of
113 *Jatropha curcas* revealed the presence of saponin, steroids, tannin, glycosides, alkaloids and flavonoids
114 in the extracts. Nishaa *et al*, [21] found tannins and anthraquinones (the largest group of quinones) to
115 possess antibacterial effects by inhibiting nucleic acid synthesis. Anthraquinones were absent in the root
116 and stem-bark extracts. This is similar to the work of Nishaa *et al* [21] who investigated the ethanol
117 extract of *Maranta arundinacea* rhizomes and found alkaloids to be present in the ethanolic extract, but
118 absent in the ethyl-acetate fraction. The presence of these phytochemicals has been reported to account
119 for the exertion of antimicrobial activity by plants [23].

120 The result of antibacterial activity (Table 2, 3 and 4) of the ethanolic extracts of root, stem bark and leaf of
121 *Balanites aegyptiaca* revealed the potentiality of the plant in treating and curing diseases cause by the
122 tested bacteria and other microorganisms. The ethanolic extract of the root, stem bark, and leaves has
123 inhibited the growth of *Klebsiella pneumoniae* and *Staphylococcus aureus*, with MIC values with low
124 concentration of 25mg/ml for *S. aureus* and 12mg/ml for *K. pneumonia*, respectively.

125
126 It was shown from previous studies that the root extract of the plant has more and lethal effect on
127 microorganisms, followed by the bark among the various part tested. But, based on the present work and
128 the statistical analysis, the leaf extract showed MIC value with low concentration of 12.5mg/ml for *S.*
129 *aureus* sand 50mg/ml for *K. Pneumoniae*, than the root and stem bark extract on both the tested bacteria
130 (Table 5). The root showed MIC value with concentration of 50mg/ml for both the tested organisms. The
131 result of the MIC and MBC also showed that *Balanites aegyptiaca* is bactericidal (Table 6).

132

133 **Table 1:** Qualitative phytochemical screening of *Balanites aegyptiaca*

Root	Stem Bark	Leaf
Ethanol	Ethanol	Ethanol

Alkaloids	+	+	+
Tannins	+	+	+
Flavonoids	+	+	+
Saponins	+	+	+
Steroids	+	+	+
Anthraquinones	-	-	+

+ = Present, - = Absent.

134
135

136
137 **Table2:** The mean of the sensitivity test of the microorganisms to the ethanolic root extract of *Balanites*
138 *aegyptiaca*

Microorganisms	Zones of inhibition (mm)					
	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	6.125mg/ml	control
<i>K. pneumoniae</i>	10	5	2	2	26	
<i>S. aureus</i>	16	13	9	7	4	24

143

144 Control: Ceftriaxone

145

146

147 **Table 3:** The mean of the sensitivity test of the microorganisms to the ethanolic stem bark extract of
148 *Balanites aegyptiaca*

Microorganisms	Zones of inhibition (mm)					
	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	6.125mg/ml	control
<i>K. pneumoniae</i>	14	11	9	5	3	24
<i>S. aureus</i>	8	12	9	7	5	25

153 Control: Ceftriaxone

154

155 **Table 4:** The mean of the sensitivity test of the microorganisms to the ethanolic leaf extract of *Balanites*
 156 *aegyptiaca*

Microorganisms	Zones of inhibition (mm)					control
	100mg/ml	50mg/ml	25mg/ml	12.5mg/ml	6.125mg/ml	
<i>K.pneumoniae</i>	16	12	9	7	5	26
<i>S. aureus</i>	20	16	13	10	6	24
Control: Ceftriaxone						



163 **Table 5:** Minimum inhibitory concentration (MIC) for microorganism of different extracts of *Balanites*
 164 *aegyptiaca* in mg/ml

Microorganisms	Root	Stem bark	Leaf
<i>K. pneumoniae</i>	25	25	12.5
<i>S. aureus</i>	50	25	25

169 **Table 6:** Minimum Bactericidal Concentration (MBC) for microorganism of different extracts of *Balanites*
 170 *aegyptiaca* in mg/ml

Microorganisms	Root	Stem bark	Leaf
<i>K. pneumoniae</i>	50	25	12.5
<i>S. aureus</i>	50	25	50

176 **Plate 1:** Inhibition zone of the root extract at different concentrations against *Klebsiella pneumoniae* and
 177 *Staphylococcus aureus*





183
184
185
186
187
188
189
190
191
192
193
194

Klebsiella pneumoniae

Staphylococcus aureus

Plate 2: inhibition zone of the stem bark extract at different concentrations against *Klebsiella pneumoniae* and *Staphylococcus aureus*




203 *Klebsiella pneumoniae*

Staphylococcus aureus


204
205
206

Plate 3: Inhibition zone of the leaf extract at different concentrations against *Klebsiella pneumoniae* and *Staphylococcus aureus*

207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234



Klebsiella pneumoniae



Staphylococcus aureus

3.0 Conclusion

Balanites aegyptiaca is traditionally used in treatment of various ailments. The phytochemical screening of the ethanol extract of root, stem bark, and leaves revealed the presence of; alkaloids, flavonoids, saponins, tannins, and steroids, in both and anthraquinones in the leaf only.

The result of the antibacterial activity of the ethanol extract of the root, bark and leaf of *Balanites aegyptiaca* inhibited the growth of *Klebsiella pneumoniae* and *Staphylococcus aureus* with MIC value of 12.5mg/ml for *K. pneumoniae* and 25mg/ml for *S. aureus*.

The demonstration of antibacterial activity of the *Balanites aegyptiaca* is indeed a development that will help to discover new antibiotics that can serve for treatment of infections that are caused by bacteria that are becoming resistant to most of the antibiotics used for treatment of diseases caused by the microorganisms. The fact that plants are very common makes it a cheaper alternative for drugs development for human consumption and use.

235 **References**

- 236 1. Abdallah EM. Plants: An alternative source for antimicrobials. *J. Appl. Pharma. Sci.* 2011 1(6):
237 16-20.
- 238 2. Abdel-Rahim EA, El-Sandany SS, Wasif MM. Biochemical dynamics of the Hypocholestrolemic
239 action of *B. aegyptiaca* fruit. *Food Chem.* 1986 20: 69-87.
240
- 241 3. Chothani DL, Vaghasiya HU. A review on *Balanites aegyptiaca* Del (desert date): phytochemical
242 constituents, traditional uses, and pharmacological activity. *Phcog. Rev.* 2011 5 (9): 55-62.
243
- 244 4. Misrak K, Amare A, Mohammed Y, Efficacy of plant extracts, traditional materials and
245 antibacterial chemicals against *Xanthomonas campestris* pv. *vesicatoria* on tomato seed. *African*
246 *Journal of Microbiology Research*, 2013; 7: 20: 2395-2400
247
- 248 5. Yinger H, Yewhalaw D, Traditional medicinal plant knowledge and use by local healers in Sekoru
249 district, Jimmazon, Southwestern Ethiopia. *Journal of Ethnobiology and Ethnomedicine.* 2007 3:
250 24- 30
251
- 252 6. Alaribe SI, A Survey of the Importance and Problems of Traditional Health care medicine, A case
253 study of Ezinihitte Mbaise L.G.A. Imo State 2008. Unpublished B.Sc. project, A.I.F.C.E. Owerri,
254 Imo State.
255
- 256 7. Vaidya AD, Devasagayam TP. Current status of herbal drugs in India: an overview. *Journal of*
257 *Clinical Biochemical Nutrition*, 2007 41:1-11.
258
- 259 8. Ogunleye DS, Ibitoye SF, Studies of antimicrobial activity and chemical constituents of *Ximema*
260 *Americana*. *Tropical Journal of Pharmacology* 2003 2:239-241.
261
- 262 9. Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC, Seal T, Mukherjee
263 B.. Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the
264 management of neurodegenerative diseases. *Journal of Ethno-Pharmacology* 2003 84:131-138.
265
- 266 10. Parker JD, Burkepile DE. Recent advances in plant-herbivore interactions. *Public Medicine* 2017
267 1(6): 1-3
268
- 269 11. Sofowora A. Medical Plants and Traditional Medicine in Africa. (2ndedn). Spectrum books Ltd,
270 Ibadan. Nigeria 2006. PP. 150 –153.
271
- 272 12. Cowan ST, Steel KJ. (2004). Manual for the identification of medical bacteria. Cambridge
273 University Press, New York. PP. 45 – 63.
274
- 275 13. Barry AL, Thornsberry C, Susceptibility tests: diffusion test procedures. In Balows, A. (ed.):
276 *Manual of Clinical Microbiology*, 1991 40: 213-219.
277
- 278 14. Bauer AW, Kirby MDK, Sherras JC, Trick M. Antibiotic susceptibility testing by standard single
279 disc diffusion method. *American Journal of clinical Pathology* 2003 45:490-496.
280
- 281 15. Abalaka ME, Mann A, Adeyemo SO. Studies on in-vitro antioxidant and free radical scavenging
282 potential and phytochemical screening of leaves of *Ziziphus mauritiana* L and *Ziziphus*
283 *spinachristi* L compared with Ascorbic acid. *Journal of Medical Genetics and Genomics*, 2011
284 3(2):28-34.
285

- 286 16. Doughari JH, Pukuma MS, De N. Antibacterial effects of *Balanites aegyptiaca* L. Drel. And
287 *Moringa oleifera* Lam. on *Salmonella* Typhi. *African Journal of Biotechnology*. 2007 6 (19), pp.
288 2212-2215.
289
- 290 17. Rotimi VO, Laughon BE, Bartlett JG, Mosadami HA. Activities of Nigerian chewing stick extracts
291 against *Bacterioides melaninogenicus*. *Antimicrobial Agents of Chemotherapy* 1988, 32: 598-
292 600.
293
- 294 18. Adebayo EA, Ishola OR. Phytochemical and antimicrobial screening of crude extracts from the
295 root, stem bark and leaves of *Terminalia glaucescens*. *African Journal of Pharmacy and*
296 *Pharmacology* 2009 3 (5): 217-222.
297
- 298 19. Igbinsola OO, Igbinsola EO, Aiyegoro O. A (2009). Antimicrobial activity and phytochemical
299 screening of stem bark extracts from *Jatropha curcas* (Linn). *African Journal of Pharmacy and*
300 *Pharmacology*. 3 (2): 058-062
301
- 302 20. Harbone, JB. Phytochemical methods. Chapman and Hall Limited, London. 1999 PP. 149 – 188
303
- 304 21. Nishaa S, Vishnupriya M, Sasikumar JM, Gopalakrishnan VK. Phytochemical Screening and GC-
305 MS Analysis of Ethanolic Extract of Rhizomes of *Maranta arundinacea*. *Research Journal of*
306 *Pharmaceutical, Biological and Chemical Sciences*. 2013 4 (2): 52
307
- 308 22. Lutterodt GD, Ismail A, Basheer RH, Baharudin, HM. Antimicrobial effects of *Psidium guajava*
309 extracts as one mechanism of its antidiarrheal action. *Malaysia Journal of Medical Science*. 1999
310 6(2): 17-20
311
- 312 23. Chapaga B, Wiesma Z. Larvicidal effects of aqueous extracts of *Balanites aegyptiaca* (desert
313 date) against the larva of *Culex pipiens* mosquitoes. *African Journal of Biotechnology*, 2005
314 4:1351-1354
315

316
317
318
319
320
321