

**Effect of Diethylcarbamazine (DEC) in Relapse cases of Nephrotic Syndrome in Filarial Endemic Region: A case Series**

Dr Jayendra R Gohil<sup>1</sup>, Dr Tanmay P Vagh<sup>\*2</sup>, Dr Dimpal Patel<sup>2</sup>, Dr Asha Patel<sup>2</sup>

1 Professor

2 Resident

\*corresponding author

From- Department of Paediatrics, Govt. Medical College, Bhavnagar, Gujarat.364001

**Author Contribution**

This work was carried out in collaboration among all authors. Author JG designed the study and author TV wrote the first draft of the manuscript. Authors DP and AP managed the literature searches. All authors read and approved the final manuscript.

**Keywords**

Nephrotic syndrome; Diethylcarbamazine DEC; Filariasis; Levamisole; Relapse; steroid sensitive;

**Comment [UP1]:** Put this immediately after the summary

**Abbreviations**

UTI; urinary tract infection; DEC; diethyl carbamazine citrate; T temperature; AG abdominal girth; URTI upper respiratory tract infection; D.O.A day of admission; CP crystalline penicillin

**Comment [UP2]:** The full words must written in the text (manuscript) where they appear first and the abbreviations can follow thereafter

## Abstract

**INTRODUCTION-** Incidence of nephrotic syndrome and filariasis both is high in China, Japan & India. Studies have shown association of filariasis in NS. In coastal belt of Gujarat, Filariasis occurs as the mosquito responsible is still prevalent. Therefore, filariasis association may be causing persistence of edema in NS relapse cases. After a 7<sup>th</sup> relapse patient was treated successfully with diethyl carbamazine citrate (DEC), we decided to study, the effect of DEC on weight loss and urine protein, in relapse patients of NS.

**MATERIAL METHOD-** In relapse patients of NS, DEC was given by oral route in dose of 72 mg/kg/cycle for 7 days. Weight record and urine protein was measured daily. Steroid as Tab prednisolone was administered at 2 mg/k/d.

**RESULTS-** ~~The 1<sup>st</sup> case was 7<sup>th</sup> relapse a~~ steroid dependant nephrotic syndrome with the 7<sup>th</sup> relapse; He had been on prednisolone and levamisole for 3 years. DEC was started on day 3 of admission and response was seen on the 5<sup>th</sup> day. ~~Urinary protein became nil on day 10, and patient has been relapse free since for 1 year. In each of the other 4 cases of with 1<sup>st</sup>, 2<sup>nd</sup>, 2<sup>nd</sup> and 4<sup>th</sup> relapse respectively, response of DEC was seen within 2 days. Thus, after starting DEC weight and urine protein reduced within 2 days in 5 relapse cases. Filaria was not detected in blood film of any patient and Elisa tests sent done in 2 were as negative.~~

Formatted: Superscript

**CONCLUSION-** ~~After starting steroid therapy, no satisfactory response, but after starting DEC, decrease in weight was seen and urine protein became nil within a week. DEC works well in filariasis and has also effects on immune system. Levamisole used as immunomodulator, also acts as an antifilarial agent. DEC~~ may be considered in relapse cases of NS in Filaria endemic regions as it is faster acting, ~~cost~~ effective and safe.

## INTRODUCTION

Nephrotic syndrome is characterized by heavy (nephrotic range) proteinuria as  $>3.5\text{gm}/24\text{hr}$  or a urine protein creatinine ratio  $>2$ , hypoalbuminemia ( $< 2.5\text{ g/dl}$ ), hyperlipidaemia (cholesterol  $>200\text{mg/dl}$ ), and edema.[1]

Prednisolone suppresses autoimmunity in nephrotic syndrome & so induces remission early. But long term use of prednisolone can cause steroid toxicity in frequent relapsers, steroid resistant and steroid dependant nephrotic syndrome.[2]

Diethyl carbamazine citrate (DEC) is an anti-parasitic agent used in the treatment of lymphatic filariasis, tropical pulmonary eosinophilia, and loiasis & acts by inhibiting arachidonic acid metabolism. DEC is administered by oral route in the dose of  $72\text{mg/kg/cycle}$  spread over 5-15 days (in our study we were giving for  $10\text{ m/k/d}$  for 7 days). DEC alters organelle membrane of Mf (microfilariae) and affects muscular activity of Mf. Side effects of DEC due to release of antigens from dying filariae are fever, skin rash, and swollen tender glands in neck, visual disturbance, nausea, dizziness, and itching.[3] It is safe in pregnancy.

Levamisole as immunomodulator for nephrotic syndrome is more effective in filarial endemic countries and beneficial effect of levamisole are reported only from south east asia countries.[4] As we see glomerular pathology occurs because of parasitic infections, instead of using levamisole, we can use better a more specific antifilarial agent, DEC, which has lower side effects.

Comment [UP3]: Delete from introduction

Comment [UP4]: Write in full

## Method

Here we are discussing the effect of DEC in minimal change nephrotic syndrome patients who had responded to steroids and who presented again with relapse. DEC is not a 1<sup>st</sup> line treatment of nephrotic syndrome here, we were using it when they such patients came with relapse.

In these 5 cases of minimal change nephrotic syndrome; DEC was administered by oral route in the dose of 72 mg/kg/cycle for 7 days (Table 1). After starting it, daily-weight was recorded and urine protein was measured daily (Figure 3-7). Steroid as Tab prednisolone was administered at 2mg/kg/d. Investigations are mentioned in Table 2.

**Protocol:** Each patient was treated as follows in relapse case of nephrotic syndrome: Prednisolone 2mg/kg/day till remission and then switch over to alternate day therapy. Immunomodulators are continued or added as required. DEC was started in all patients at 72mg/kg/cycle total dose over a period of 7 days.

## Course during treatment and Outcome

In the index case A, with 7<sup>th</sup> relapse, foci of infection was U.T.I. so treatment for UTI was started. Till 4<sup>th</sup> day no recovery was seen so considering relapse and no any recovery seen even after starting antibiotics so started DEC on 3<sup>rd</sup> D.O.A. Edema started to reduce on 5<sup>th</sup> D.O.A and slowly it disappeared on 8<sup>th</sup> D.O.A, urinary protein became clear on 9<sup>th</sup> D.O.A (figure 3).

In the 2<sup>nd</sup> case (B) with the 4<sup>th</sup> relapse, foci of infection was chronic tonsillitis so considering bacterial infection we started antibiotics, but there was no response came so DEC was started on 4<sup>th</sup> D.O.A and after that edema gradually started to reduce on 6<sup>th</sup> day. (figure 4).

~~In The 3<sup>rd</sup> case (C) was seen during the 2<sup>nd</sup> relapse, The~~ foci of infection was UTI and treatment for UTI started. Even after 7 days no any signs of recovery was seen so we commenced DEC started on the 7<sup>th</sup> D.O.A and after that edema REDUCED reduced on 9<sup>th</sup> D.O.A and Urinary Protein became clear on 10<sup>th</sup> D.O.A (figure 5).

In the 4<sup>th</sup> case (D) with 2<sup>nd</sup> relapse, foci of infection was due to URTI and antibiotic was started. Even after 6 days on antibiotics no changes in edema and proteinuria was seen so DEC started was started on 6<sup>th</sup> D.O.A and after that edema started to resolve slowly from the 7<sup>th</sup> D.O.A and Urinary Protein became clear on 9<sup>th</sup> D.O.A (figure 6).

~~In 5<sup>th</sup> case E, admission done and The~~ foci of infection in the 5<sup>th</sup> case (E), was URTI and symptomatic and specific treatment was started. But no response till 6<sup>th</sup> day so DEC started on 6<sup>th</sup> D.O.A and after that edema started to reduce on 9<sup>th</sup> D.O.A & Urinary Protein became clear on 4<sup>th</sup> D.O.A (figure 7).

No adverse events were detected by using DEC & Filaria was not detected in midnight thick blood film of any patient. Elisa test done in 2 patients was negative.

Comment [UP5]: Why capital letters?

Comment [UP6]: What is this? 6<sup>th</sup>

Comment [UP7]: Sure of this? DEC was started on 6<sup>th</sup> day of admission

Table 1: Case Details

History	Examination	Treatment
A.4 Y/F: Eyelid and pedal swelling since 7 day abdominal distension since 2-3 days. decreased urine output since 2 days UTI (Ecoli +ve) 7 <sup>th</sup> relapse, steroid sensitive	T Normal Eyelids swelling Pedal edema (pitting), transverse umbilicus AG-64cm BP-94/68mmHg	Inj.ciprofloxacin(7day) Tab.levamisole (4day) Tab.prednisolone(2mg/kg/dy)(20mg)(8day) DEC <del>for 7days</del> Tab zinc <del>for 12days</del>
B.2 Y/M: Swelling of eyelid since 3days cough and cold since 2days Tonsillitis 4 <sup>th</sup> relapse, steroid dependent	T-Normal AG-64cm BP-96/70mmHg Throat-congested, Tonsil enlarged Eyelids swelling Pedal edema +(pitting)	Inj. Amoxiclav <del>for (5 days)</del> Tab. Prednisolone <del>for (6days)</del> DEC for 7days
C.5.5 Y/M: fever 4day, Eyelid swelling 4 day, Swelling over abdomen 2 days, UTI (urine pus cells) 2 <sup>nd</sup> relapse, steroid sensitive	T-Normal AG-64cm BP-96/70mmHg Pedal edema + transverse umbilicus	Inj.ciprofloxacin (10 d) Tab.prednisolone(10 d) Tab.amoxycillin(1day) DEC <del>for 7days</del> Tab zinc <del>for(11 days)</del>
D.6 Y/F: eyelid swelling-2 days Pedal swelling-2 days Cough-2 days URTI 2 <sup>nd</sup> relapse, steroid sensitive	T-Normal AG-60cm BP- 100/70mmHg Throat- congested and Tonsil enlarged Eyelid swelling Pedal edema (pitting)	Inj.amoxiclav <del>for 4 days</del> Tab.prednisolone <del>for (4 days)</del> DEC for 7 days
E.6 Y/M: eyelid swelling since 3 days, swelling over abdomen since 3 days tonsillitis 1 <sup>st</sup> relapse, steroid sensitive	T-Normal AG-62cm, BP-98/66mmHg Throat-congested PA-visible veins, edema over abdomen, liver+2cm	Inj.CP <del>for 5 days</del> Inj.amoxyclav- <del>for 4 days</del> Tab.prednisolone <del>for (9 days)</del> DEC <del>for 7days</del>

Comment [UP8]: What does T stand for?

Doses- Inj.ciprofloxacin- 10mg/kg/d; Tab. Levamisole- 2mg/kg/d, 3d in a week for 6 week; Tab. Prednisolone- 2mg/kg/d; DEC- 72mg/kg/cycle; Tab zinc-10mg/kg/d; Inj.CP-50,000IU/kg/dose


Table 2: Investigations on admission

<b>Investigations</b>	<b>Case 1</b>	<b>Case 2</b>	<b>Case 3</b>	<b>Case 4</b>	<b>Case 5</b>
Total Protein (g/dl)	3.9	4	4.02	3.92	3.2
S. Albumin (g/dl)	1.42	1.5	1.42	1.63	2.1
S. Globulin (g/dl)	2.48	2.5	2.60	2.29	2.8
A/ G ratio	0.57	2.5	0.55	0.71	0.75
S. Cholesterol (mg/dl)	525	385	467	321	320
24hr Urine Protein (mg/24hr)	>2000	670	452	395	410
Urine Protein Creatinine ratio (mg/mg)	7.9	9.19	12.85	12.5	8
<b>Weight (kg) Maximum/ on discharge</b>	<b>21.94/18.1</b>	<b>20/18.8</b>	<b>21.3/19.3</b>	<b>20.2/18.2</b>	<b>20.63/18.2</b>

**S serum; A/G serum albumin/ globulin ratio**


Normal values: T. protein 6-8g/dl; S. Albumin 3.5-5g/dl; S.Globulin-0.5-3g/dl; A/G ratio-0.8-2.0; S. cholesterol <200mg/dl; 24hr urine protein 30-150 mg/24hr; Urine protein creatinine ratio <0.2mg/mg

Figure 3: Case A-4yr female- 7<sup>th</sup> Relapse



★ ----Shows urine Urinary Protein becomes nil


Figure 4: Case B 2yr male 4<sup>th</sup> Relapse



★---Shows urine Urinary Protein becomes nil




Figure 5: Case C. 5.5yr male -2<sup>nd</sup> relapse



★----Shows urine Urinary Protein becomes nil


Figure 6: Case D -6yr Female – 2<sup>nd</sup> relapse

**Comment [UP9]:** Title should be under not above the figure



★ ----Shows urine Urinary Protein becomes nil

Figure 7: Case E .6yr Male -1<sup>st</sup> Relapse



Comment [UP10]: This title should be below the figure

★ - Shows urine Urinary Protein becomes nil

## Discussion

### Discussion:

It is a denovo study in nephrotic syndrome patients. Initially reports were done to find out any foci of infection and as per report symptomatic management was done. Still in all these cases edema and heavy range proteinuria, did not disappear so considering connection of nephrotic syndrome with filariasis we started DEC. After starting DEC as effect occur initially weight loss starts slowly and urine heat coagulation becomes clear day by day.

In 1966, a new drug levamisole was discovered [5] and due to its immunomodulator property [6], its use had been- increased in frequent relapsers, steroid resistant & steroid dependant nephrotic syndrome. But its side effects like leucopenia, rash, seziures limits its use in nephrotic syndrome [7].


As per figure 1 Levamisole attenuates podocyte apoptosis and increase in RhoA (ras homolog family member A) activity and decrease in degradation of synaptopodin protein; and so prolonging podocyte survival [8].

**Comment [UP11]:**

**Comment [UP12]:** Please correct this sentence.  
It is not clear

**Figure 1** Mechanism of action of Drugs used in Nephrotic Syndrome (Figure reproduced from 'Nonimmunologic targets of immunosuppressive agents in podocytes' by Yoo TH [8])

**Comment [UP13]:** This figure is not necessary in the discussion



GCR=Glucocorticoid receptor ; Bcl2=B cell lymphoma 2; p53= tumor protein 53; b7-1=membrane protein; RhoA=ras homolog family member A; SMPD-3B=Sphingomyelin phosphodiesterase acid like 3B; Rac1=Rac family small GTPase 1

Incidence of nephrotic syndrome was quite high in China, Japan, India and many Asian countries in 1950s[9] &and incidence of lymphatic filariasis was also quite high in those countries.[10] As per figure 2 There is a clear association of filariasis and nephrotic syndrome.[11] In the Coastal belt of Gujarat, filariasis still occurs as the mosquito responsible for it , therefore filariasis association causing persistence of edema in nephrotic syndrome is possible.


**Comment [UP14]:** This phrase is not clear. Please clarify

After filarial infection of human body the generated toxins affect~~ts~~ several compartments of the immune system. And play a role in fighting parasitic infections & in the pathogenesis of associated glomerulopathies. During infection, released inflammatory mediators directly damage the different glomerular cell types & participate in the activation of specific subsets of T and B cells, resulting in different levels of antigen, antibody & immune complexes. Depending on the site of the immune-complex deposition and on the type of primary damaged glomerular cells, different glomerular lesions develop.[12]

**Comment [UP15]:** Merge these 2 sentences. What plays a role in fighting parasitic infection, the toxins or the immune system?

**Figure 2** Effect of Parasitic Infection (Figure reproduced from 'The nephrotic syndrome associated with filariasis' study done by Yap HK[11])

**Comment [UP16]:** This figure is not necessary under the discussion of the patients reviewed in this manuscript. What is shown in the figure has already been described in the text



NK cell=Natural killer cell; GN=glomerulonephritis; Ag=antigen; Ab=Antibody; GP-glomerulopathy

DEC works well in filariasis and other parasitic infections and due to serious side effects of levamisole & due to possible association of filariasis with nephrotic syndrome, it can be used in nephrotic syndrome as it is cost effective and safety is better than levamisole.

**Comment [UP17]:** Is it possible to postulate as to why DEC was of therapeutic benefit in these patients? Could it be due to its immune modulatory effect or anti microfilarial effect?

### Conclusion

As per above study, After Starting DEC, weight and urine protein started to reduce within 2 days.

DEC works well in filariasis and has also effects on immune system. Levamisole used as immunomodulator, also acts as an antifilarial agent. DEC may be considered in relapse cases of NS in Filaria endemic regions.

**Comment [UP18]:** Delete

It is cost-effective and safe. Further trials with controls and efforts to detect antigen is recommended.

### *What is already known?*

Levamisole is used for steroid dependant nephrotic syndrome. It is also a wormicidal drug acting on filarial worms. Filaria has been linked with minimal change nephrotic syndrome and filarial antigen has been observed in kidney biopsy. Diethylcarbamazine (DEC) is the drug of choice for Filarial infection.

### *What this case series adds?*

The effect of DEC (when added to Prednisolone) in relapse cases of nephrotic syndrome in filarial endemic region appears to reduce edema and proteinuria within 3-4 days of administration. DEC need to be studied in relapse cases in place of Levamisole.

### *Limitations of study*

This is a denovo case series and it is a proof of concept case series. A study with control is recommended which is lacking in this study.

### **Disclaimer regarding Consent/Ethical Approval:**

As per university standard guideline participant consent and ethical approval has been collected and preserved by the authors.



## References

1. Pais P, Eliis DA. Nephrotic syndrome, in Nelson Textbook of Pediatrics, Kleigman RM, Stanton BF, Geme JW, Schor NF, Behrman RE, Saunders Elsevier, 2012;19:1801-06.
2. Hahn D, Hodson EM, Willis NS, Craig JC. Corticosteroid therapy for nephrotic syndrome in children. Cochrane Database of Systematic Reviews 2015, Issue 3: CD001533.
3. WHO Model Prescribing Information: Drugs Used in Parasitic Diseases – 2/e: Helminths: Lymphatic filariasis: Diethylcarbamazine. *who.int*. 1995. p. 152
4. Kuhlmann FM, James MF. Antiparasitic Agents. *Infectious Diseases*. 2017;4(2)1345-1372.
5. Narasimham MV, Roychowdhury SP, Das M, Rao CK. Levamisole and mebendazole in the treatment of bancroftian infection. *Southeast Asian J Trop Med Public Health*. 1978 Dec;9(4):571-5.
6. Gupta M. Levamisole: A multi-faceted drug in dermatology. *Indian J Dermatol Venereol Leprol* 2016;82:230-6.
7. Symoens J, Veys E, Mielants M, Pinals R. *Cancer Treat Rep* 1978 Nov;62(11):1721-30.
8. Yoo TH, Fornoni A. Nonimmunologic targets of immunosuppressive agents in podocytes. *Kidney Res Clin Pract*. 2015;34(2):69–75.
9. Pal A, Kaskel F. History of nephrotic syndrome and Evolution of its Treatment. *Front Pediatr*. 2016;4:56
10. SudomoM, Chayabegara S, Duong S, Hernandez L, Wu WP, Bergquist R. Elimination of Lymphatic Filariasis in Southeast Asia. *Advances in Parasitology* 2010;72(C)205-233
11. Yap HK, Yeo PP, Chiang GS, Singh M, Lim CH. The nephrotic syndrome associated with filariasis. *Ann Acad Med Singapore* 1982;11:60-3.
12. van Velthuisen MLF. Glomerulopathy associated with parasitic infections. *Parasitol* 1996;12:102–107.