

# Control of aflatoxin production in cassava produced by dry fermentation in North Kivu, Democratic Republic of Congo

## ABSTRACT

Traditionally, cassava (*Manihot esculenta* Crantz) is transformed by fermentation in water (retting) or in the open air (dry fermentation) in the DRC. In the east of the country (North Kivu), dry fermentation is the main technique for processing cassava for its detoxification and conservation. The Congolese farmers ferment the cassava to the open air using a preselected microferment contained in the scrapings of the fermented cassava previously called "MUSIYIRO". These fermentations are spontaneously directed by the microorganisms of the uncontrolled autochthonous flora. Unfortunately, toxinogenic molds are often more active in the fermentation process during which they also produce aflatoxins. This study was undertaken to help prevent the production of aflatoxins in cassava during this process. To do this, we substituted the traditional ferment with a strain of *Rhizopus oryzae* used as starter (microferment). Six successive replications, in controlled fermentation and uncontrolled fermentation, in a peasant environment (Beni, North Kivu) and fermentation directed by the strain of *Rhizopus oryzae* were carried out. Aflatoxins were then dosed in cassava flour. The results of the assay revealed an absence of aflatoxins in cassava fermented by scrapings from fermentation led by *R. oryzae*, while the non-directed fermentation controls were all contaminated with aflatoxins. These results show that it is possible to prevent the production of aflatoxins in cassava during fermentation when an aflatoxin-inhibiting microbial biomass is used which can progressively invade and colonize the fermentation site and thereby control the fermentation activities of cassava.

*Keywords: Aflatoxins inhibition; Rhizopus oryzae; Dry fermentation; Cassava.*

## 1. INTRODUCTION

Cassava (*Manihot esculanta* Crantz) is the staple food and source of nourishment for more than one billion people worldwide [1] especially in Africa, Asia and South America. Because of its high water content, cassava must be transformed into various derivatives to ensure its availability outside harvest periods and to reduce post-harvest losses. The cassava products can be fermented, dried or roasted, and the most common form is chips. This product serves as food for both human and animals and can be maintained up to one year [2].

However, tropical climate of some geographical areas of cassava production may contribute to fungal development of many species and subsequent toxinogenesis on such raw material [3]. Moreover, the processing conditions and storage premises are not always well adapted to protect cassava products from secondary contamination and/or fungal development.

The FAO estimates that 25% of the world food crops are contaminated by mycotoxin, of which the most notorious are the aflatoxins (AFTs) [4-6]. AFTs are metabolites produced primarily by the fungi *Aspergillus flavus* and *Aspergillus parasiticus*. There are four major naturally produced AFTs, referred to as B1, B2, G1 and G2 [7, 8]. The B1 is the most toxic of the AFTs and potent naturally occurring liver carcinogen [10]. Reports estimated that more than 5 billion people in developing countries world-wide are at risk of chronic exposure to AFTs through contaminated foods [9, 10]. AFTs affect livestock and poultry causing

28 reduced feed efficiency, subtle immunosuppression, growth rate and death of animals [6, 11]. Other  
29 economic adverse effects of AFTs include low yields of food and fiber crops [12].

30 Considering the importance of this crop in Democratic Republic of Congo (DRC) and subsequent possible  
31 fungal toxin production, various studies have attempted to evaluate cassava products contamination with  
32 moulds and mycotoxins. A number of potentially mycotoxigenic fungi have been isolated from cassava  
33 products and mycotoxins contamination of cassava has been documented but the potential sanitary risk  
34 of such contamination was not fully assessed [13, 14].

35 The populations of Ituri, North and South Kivu in the north-east of the Democratic Republic of Congo  
36 (DRC), mainly consume a dry fermentation product of cassava tubers (*Manihot esculenta* Crantz) called  
37 "Kithabiro". Unfortunately, to date, these cassava transformations are still artisanal, rudimentary and  
38 unhygienic; fermentations are spontaneously directed by the microorganisms of the uncontrolled  
39 autochthonous flora.

40 Among the microorganisms involved in these cassava fermentations, the following strains were isolated:  
41 *Aspergillus flavus* (LINK), *A. flavus oryzae*, *A. niger*, *A. fumigatus* and *A. glaucus*, *Rhizopus oryzae* and  
42 *Mucor mucedo* [14].

43 The toxigenic molds of which *Aspergillus flavus*, *A. fumigatus* and *A. niger* responsible for the production  
44 of aflatoxins are almost permanent and actively participate in the softening of cassava and the  
45 development of characteristic aromas of traditional fermented cassava [15].

46 In fact, farmers conduct cassava fermentation in the open air using a preselected microferment contained  
47 in the fermented cassava scrapings previously called "MUSIYIRO". This "Musiyiro" is transplanted each  
48 time into the cassava batch to be fermented, basted and then covered with banana leaves and cotton  
49 bags to create the moisture needed for microbial growth and cassava fermentation without water.

50 After several successive subcultures, the various microorganisms of the site stabilize and constitute the  
51 STARTERS (Microferment) for the fermentation and the softening of cassava without water.

52 The studies carried out by Yandju [16] and Djoulde [17] in the raking of cassava called wet fermentation,  
53 have shown that the use of a non-toxinogenic efficient microferment has made it possible to obtain stable  
54 fermented cassava products and to guarantee the sanitary quality of these products.

55 Considering the high dependence of populations on cassava Fufu from dry fermentation, it is essential to  
56 consider improving cassava processing techniques in the farmer sector. Within this context, the quality  
57 and safety of agricultural products and food are surveyed to limit consumer exposure to aflatoxins.

58 The main objective of this study is to inhibit the production of aflatoxins during the dry fermentation of  
59 cassava called "fufu kithabiro" of North Kivu by the substitution of the traditional "Musiyiro" by a pure  
60 culture biomass of *Rhizopus oryzae*, molds selected for this purpose.

61

## 62 **2. MATERIALS AND METHODS**

### 63 **2.1. Collection of samples**

64 Fifty seven (57) samples (fermented cassava Fufu) were collected from the eighteen (18) sites in North  
65 Kivu used for the aflatoxin assay and thirty six (36) samples obtained from the experimental fermentation  
66 with the *Rhizopus oryzae* strain.

### 67 **2.2. Assessment of the presence of aflatoxins in the Fufu of North Kivu markets**

68 Approximately two hundred and fifty grams of dry ferment cassava chips were aseptically collected in  
69 markets in twenty one locations in North Kivu due to three samples per site for aflatoxin and isolation of  
70 non-toxinogenic and potent *Rhizopus oryzae* in the directed fermentation of cassava [17].

71 During the sampling, a macroscopic analysis of the organoleptic properties of the dry fermenting cassava  
72 Fufu was carried out on site before being sent to the laboratory for the isolation of the seeds and the  
73 aflatoxin determination. The analysis concerned the surface color of cassava chips and their aroma [18,  
74 19].

### 75 **2.3. Determination of aflatoxins in fermented cassava samples**

76 Aflatoxins were evaluated by biological detection and spectrophotometric assay. For biological detection,  
77 aflatoxins were detected by inhibiting the growth of sensitive *E. coli* C600 on nutrient agar following their  
78 diffusion.

79 The aflatoxin assay was done by the rapid multitoxin assay method using the Acquity Spectrophotometer  
80 HPTLC and the Quattro Preparer XE mass spectrophotometric at South Africa's Perishable Products  
81 Export Control Board (PPECB) (RSA).

## 82 2.4. Directed fermentation of cassava by non-toxigenic microferment

83 A culture of *Rhizopus oryzae* has been selected to serve as a microferment to replace the scrapings of  
84 the traditional fermentation called "Musiyiro" in Nande dialect. After massive seeding on a first batch of  
85 cassava under aseptic conditions, scrapings consisting of *R. oryzae* Mycelium were subcultured  
86 successively six times on six batches of cassava in the presence of two controls, the first taking up the  
87 traditional fermentation and the second one seeded with *Rhizopus oryzae* strain and incubated under  
88 aseptic conditions.

89 The incubation was done under the traditional fermentation conditions until the softening of cassava in  
90 order to put in the peasant conditions.

91 The fermented, dried cassava was removed from the site for analysis and sent to the laboratory for  
92 aflatoxin determination [20].

## 93 2.5. Statistical data analysis

94 Data were analyzed by descriptive statistics and analysis of variance (ANOVA) using Statistix Ver.8  
95 software. Difference in the levels of aflatoxin contamination was determined by the comparison of mean  
96 using least significant difference (LSD) at 5% level of significance. The mean contents of aflatoxins were  
97 transformed into  $\log(x+1)$  to normalize data prior to analysis.

## 98 3. RESULTS AND DISCUSSION

99 **Table 1. Overall aflatoxins contamination incidence in cassava samples**

Classes of aflatoxins	Cassava Samples			
	Sample size	Positive sample (%)	Mean	Range
Aflatoxin B1	63	29	0.96 <sup>a</sup>	0.34-1.95
Aflatoxin B2	63	29	0.36 <sup>b</sup>	0.12-0.74
Aflatoxin G1	63	nd	nd	nd
Aflatoxin G2	63	nd	nd	nd

100 \*The mean aflatoxin levels with the different superscript letters in the same column are significantly different ( $p < 0.05$ );  
101 mean aflatoxin levels were transformed into  $\log(x+1)$  prior to analysis

102 nd= the levels of the aflatoxin analyzed were lower than the limit of detection

103  
104 Cassava fufu samples analyzed did contain two types of aflatoxins. Aflatoxins B1 and B2 were present in  
105 fermented cassava and in 4 of our 21 fermented cassava samples from North Kivu province while  
106 aflatoxins G1 and G2 were absent in all fermented cassava samples analyzed. Aflatoxins B1 are  
107 significantly higher ( $p < 0.05$ ) than B2. A plausible reason could be that aflatoxins levels in samples of  
108 cassava analyzed were below the limit of detection (LOD): 0.2 ppb and 0.5 ppb for aflatoxin G1 and G2;  
109 respectively. Also, previous studies suggest that cassava is unlikely to be a source of aflatoxin [21, 22].  
110 Another study investigated the fungal and aflatoxins contamination of cassava products and found that  
111 aflatoxins B1, B2, G1 and G2 were lower than the limit of detection (2ppb) of analytical method (VICAM  
112 Afla Test immunoaffinity fluorometric method) that they used [23].

113 However, other studies have shown the presence of aflatoxins in cassava products [5, 24, 25]. The study  
114 conducted by Jonathan *et al* [25] showed that major spoilage (biodeteriorating) fungi of Attiéké from  
115 Ejigbo, Iwo and Adjame in West Africa were mostly molds with *Aspergillus niger* and *A. flavus* having

116 highest occurrence and *Candida albicans* and their percentage occurrence has direct effect on its food  
 117 values.

118

119 **Table 2. Aflatoxin B1 contamination (ppb) in cassava samples**

Source of samples	Cassava Samples		
	Sample size	Positive sample (%)	Mean
Alungupa	3	40	0.89 <sup>a</sup>
Beni ville	3	20	1.2 <sup>a</sup>
Butembo	3	nd	nd
Goma	3	nd	nd
Itendi	3	nd	nd
Kabasha	3	nd	nd
Kalunguta	3	nd	nd
Kanyabayonga	3	nd	nd
Kayina	3	20	1.05 <sup>a</sup>
Kiantshaba	3	nd	nd
Kiwandja	3	nd	nd
Lubero	3	nd	nd
Mabaya	3	nd	nd
Mbau	3	nd	nd
Mukuliya	3	nd	nd
Musienene	3	nd	nd
Oicha	3	20	0.71 <sup>a</sup>
Rutshuru	3	nd	nd

120 \*The mean aflatoxin B1 contents with the same superscript letters in the same column are not significantly different  
 121 ( $p>0.05$ ); mean aflatoxin levels were transformed into  $\log(x+1)$  prior to analysis  
 122 nd= the levels of the aflatoxin analyzed were lower than the limit of detection (0.15 ppb)

123 Table 2 shows that of the 21 markets sampled in North Kivu, only four markets whose cassava products  
 124 were contaminated by aflatoxin; it's about Oicha, Alungupa, Beni city and Kayina. The highest aflatoxin  
 125 contamination was noted in samples that came from Beni city market, with the mean of aflatoxins was 1.2  
 126 ppb for aflatoxin B1 followed by Kayina (1.05 ppb) and Alupunga market (0.89ppb). The least  
 127 contamination was observed to Oicha market. The same observation was made to aflatoxin B2 contents  
 128 as reported in Table 3. This presence is noticed in the samples from the 4 sites including Kayina, Oicha,  
 129 Alungupa and Beni city. The almost permanent presence of aflatoxins in the samples coming from the  
 130 same axes can be justified by the fact of permanent use of the previous fermentation scrapings by site  
 131 and by axis.

132 **Table 3. Level (ppb) of aflatoxin B2 in cassava samples**

Source of samples	Cassava Samples		
	Sample size	Positive sample (%)	Mean
Alungupa	3	40	0.34 <sup>a</sup>
Beni ville	3	20	0.49 <sup>a</sup>
Butembo	3	nd	nd
Goma	3	nd	nd
Itendi	3	nd	nd
Kabasha	3	nd	nd
Kalunguta	3	nd	nd
Kanyabayonga	3	nd	nd

Kayina	3	20	0.21 <sup>a</sup>
Kiantshaba	3	nd	nd
Kiwandja	3	nd	nd
Lubero	3	nd	nd
Mabaya	3	nd	nd
Mbau	3	nd	nd
Mukuliya	3	nd	nd
Musienene	3	nd	nd
Oicha	3	20	0.41 <sup>a</sup>
Rutshuru	3	nd	nd

133 \*The mean aflatoxin B2 contents with the same superscript letters in the same column are not significantly different  
 134 ( $p>0.05$ ); mean aflatoxin levels were transformed into  $\log(x+1)$  prior to analysis  
 135 nd= the levels of the aflatoxin analyzed were lower than the limit of detection (0.2 ppb)

136 **Table 4. Determination of aflatoxins in cassava by directed fermentation with *R.oryzae***

Fermentation number	Directed fermentation with <i>R.oryzae</i>	Control *
	Aflatoxin ( $\mu\text{g}/\text{kg}$ )	Aflatoxin ( $\mu\text{g}/\text{kg}$ )
F1	0.0	0.5
F2	0.0	1.5
F3	0.0	1
F4	0.0	1.8
F5	0.0	2
F6	0.0	2.5

137 \*with traditional fermentation, from Beni city

138  
 139 The survey established implies that cassava products are vital source of food and income for majority  
 140 people in North Kivu. Generally, in Beni city, cassava products which contained fungal growth retained  
 141 those till the time of preparation for milling when are scraped. This condition may contribute to high  
 142 contamination of aflatoxins when compared to others districts.

143 The determination of aflatoxins in cassava samples of directed fermentation with *R.oryzae* showed that  
 144 there was no aflatoxin; while the traditional fermentation cassava control incubated at the same site  
 145 contained aflatoxins B1 and B2. After six subcultures of the scrapings constituting the microferment in the  
 146 traditional fermentation, table 4 shows disparities in the production of aflatoxins in control batches.

147 Their concentration is different with a tendency to increase which probably reflects the dominance and  
 148 stability of aflatoxinogenic strains in the site after several subcultures. These results are consistent with  
 149 the studies of Yandju, 1995. The results of aflatoxin determination in 6 batches of cassava inoculated with  
 150 pure *R. oryzae* microferment showed an absence of aflatoxins in all batches, although incubation was  
 151 done under the same conditions in a traditional setting.

152 Indeed, the inoculation of the selected germs during a natural fermentation, eventually impose a pure and  
 153 quantitatively large initial biomass. This would gradually colonize cassava processing sites and destroy  
 154 the effects of toxigenic microorganisms that play a negative role in cassava fermentation under peasant  
 155 conditions [15, 17]. Most recently, R. Bandyopadhyay at al (2016) developed a technique using a non-  
 156 aflatoxinogenic *Aspergillus* strain to provide biological control of maize in storage. To date, it is the only  
 157 product called "Aflasale" consisting of a suspension of aflatoxinogenic mold spores to protect agricultural  
 158 products against the production of aflatoxins by aflatoxinogenic molds.

159 The results obtained in this study suggest that *Rhizopus oryzae* strains used in controlled fermentation  
 160 have been a potent inhibitor of aflatoxins during dry fermentation.

161

#### 162 4. CONCLUSION

163

164 The results of this study indicates the presence of aflatoxins B1 and B2 in the cassava fufu samples from  
165 the Alungupa, Beni city, Kayina and Oicha sites while all the samples obtained from fermentations  
166 directed with *R.oryzae* did not have traces of aflatoxin.

167 Furthermore, the determination of aflatoxins in cassava samples fermented by scrapings from the  
168 biomass of *Rhizopus oryzae* non-aflatoxinogenes showed a complete absence of these even when the  
169 *Rhizopus* strains were sown without asepsis under peasant conditions.

170 It is therefore very advantageous to use non-aflatoxigenic strains for the dry fermentation of cassava. It  
171 will produce large quantities and popularize. This would undoubtedly reduce the risks associated with  
172 aflatoxins.

#### 173 REFERENCES

- 174 1. FAO (2011) 'Global information and early warning system on food and agriculture', FAO Food  
175 Outlook: Global Market Analysis, November.
- 176 2. Wareing PW, Westby A, Gibbs JA, Allotey LT, Halm M. Consumer preferences and fungal and  
177 mycotoxin contamination of dried cassava products from Ghana. *Int. J. Food Sci. Tech.*2001;  
178 36(1):1–10.
- 179 3. Bankole SA, Adebajo A. Mycotoxins in food in West Africa: current situation and possibilities of  
180 controlling it. *Afr. J. Biotech.*2003; 2(9): 254–263.
- 181 4. Westby A. Cassava utilisation, storage and small-scale processing. In: Hillocks R, Tresh J, Bellotti A,  
182 editors. *Cassava: Biology, Production and Utilization*. Ed.CABI Publishing, Wallingford, UK; 2002.
- 183 5. Adjovi CSY, Gnonlonfin BJJ, Bailly S, Bailly JD, Tadriss S et al. Occurrence of mycotoxins in cassava  
184 (*Manihot esculenta* Crantz) and its products. *International Journal of Food Safety, Nutrition and Public*  
185 *Health*. 2015; 5: 217-247. <10.1504/IJFSNPH.2015.070157>. <hal- 01594243>
- 186 6. Boutrif E. Prevention of Aflatoxin in Pistachios. *Food, Nutrition and Agriculture*.1998; 2:1.
- 187 7. Akande KE, Abubakar MM, Adegbola TA, Bogoro SE. Nutritional and health implications of  
188 mycotoxins in animal feeds: A review. *Pak J Nutr*. 2006; 5:398–403.
- 189 8. Bryden WL. Mycotoxin contamination of the feed supply chain: Implications for animal productivity  
190 and feed security. *Animal Feed Science and Technology*. 2012; 173: 134-58.
- 191 9. Ogiehor IS, Ikenebomeh MJ, Ekundayo AO. The bioload and aflatoxin content of market garri from  
192 some selected states in southern Nigeria: Public health significance. *Afr Health Sci*. 2007; 7:223–227.
- 193 10. Wu F, Khlangwiset P. Health and economic impacts and cost-effectiveness of aflatoxin-reduction  
194 strategies in Africa: Case studies in biocontrol and postharvest interventions. *Food Additives &*  
195 *Contaminants*. 2010; 27: 496–509.
- 196 11. Shephard GS. Aflatoxin and food safety: Recent African perspectives. *J Toxicol*. 2003; 22:267–286.
- 197 12. Strosnider H, Azziz-Baumgartner E, Banziger M, Bhat RV, Breiman R, Brune MN et al. Workgroup  
198 report: Public health strategies for reducing aflatoxin exposure in developing countries. *Environmental*  
199 *Health Perspectives*. 2006; 114: 1989–1903.
- 200 13. Jonathan SG, Abdul-Lateef MB, Ayansina ADV. Fungal and aflatoxin detection in fresh and stored  
201 'garri ijebu' (locally processed food) *Manihot esculenta*. *Report and Opinion*. 2013; 5: 13-19.
- 202 14. Ashraf MW. Determination of aflatoxin levels in some dairy food products and dry nuts consumed in  
203 Saudi Arabia. *Food Public Health*. 2012; 2:39–42.
- 204 15. Yandju DL, Matondo KL, Munyanganizi B. Les moisissures toxigènes impliquées dans le  
205 ramollissement des racines tubéreuses du manioc en fermentation sèche. *Transformation Alimentaire*  
206 *du Manioc*. ed. ORSTOM, 1995.French.
- 207 16. Manjula K, Hell K, Fandohan P, Abass A, Bandyopadhyay R. Aflatoxin and fumonisin contamination  
208 of cassava products and maize grain from markets in Tanzania and Republic of the Congo. *Toxin*  
209 *Reviews*. 2009; 2:63 - 69.
- 210 17. Djoulde RD. Mise au point d'un ferment mixte destine a la bioconversion des tubercules de manioc  
211 cyanogene. *Sciences du Vivant [q-bio]*. Migration - université en cours d'affectation, 2004. French.
- 212 18. Aguruo CU, Onda MA, Omoni V, Tand Ogonna IO. Characterization of moulds associated with  
213 processed garri stored for 40 days at ambient temperature in Makurdi, Nigeria. *African Journal of*

- 214 Biotechnology. 2014;13(5):673-677. DOI: 10.5897/AJB2013.12187 ISSN 1684-5315 ©2014  
215 Academic Journals <http://www.academicjournals.org/AJB>
- 216 19. Brudzynski A, Van Pee W, Kornaszewski W. The occurrence of aflatoxin B1 in peanuts, corn and  
217 dried cassava sold at the local market in Kinshasa, Zaire; its coincidence with high hepatoma  
218 morbidity among the population. Zesz Probl Postepow Nauk Roln. 1977; 189:189–117.
- 219 20. Peng W, FengYZ, Maohua Q. Study on the Novel Strain *Rhizopus Oryzae* ATCC2809's Growth and  
220 Fermentation. International Journal of Scientific Engineering and Research. 2013; 1(3): 2347-3878
- 221 21. Gnonlonfin GJB, Adjovi CSY, Katerere DR, Shephard GS, Sanni A, Brimer L. Microflora and absence  
222 of aflatoxin contamination of commercialized cassava chips in Benin, West Africa. Food Control.  
223 2012; 25(2): 333-337
- 224 22. Matsiko F, Kanyange C, Ingabire G, Dusingizimana T, Vasanthakaalam H, Kimonyo A. Detection and  
225 quantification of aflatoxin in cassava and maize flour sold in Kigali open markets, Rwanda.  
226 International Food Research Journal. 2017; 24(1): 459-464.
- 227 23. Chiona M, Ntawuruhunga P, Benesi IRM, Matumba L, Moyo CC Aflatoxins contamination in processed  
228 cassava in Malawi and Zambia. African journal of food and Agricultural nutrition and  
229 Development. 2014; 14(3): 8809-8820.
- 230 24. Gacheru PK, Abong GO, Okoth MW, Lamuka PO, Shibairo SA, Katama CKM. Microbiological safety  
231 and quality of dried cassava chips and flour sold in the nairobi and coastal regions of Kenya. African  
232 Crop Science Journal. 2016; 24 (1):137–143.
- 233 25. Jonathan SG, Bello TS, Asomoloye MD. Food Values, Spoilage Moulds and Aflatoxin Detection in  
234 'Attiéké' (A Cassava Fermented Product). Journal of Microbial & Biochemical Technology. 2017; 9(5):  
235 244-248. DOI: [10.4172/1948-5948.1000372](https://doi.org/10.4172/1948-5948.1000372).

236

237

238

239

240

241