

DETERMINANTS AND PREVALENCE OF METABOLIC SYNDROME IN SUBJECTS ATTENDING THE GENERAL OUTPATIENT DEPARTMENT OF THE FEDERAL MEDICAL CENTRE KATSINA STATE, NORTH WEST NIGERIA

ABSTRACT

Metabolic syndrome (MS) is a term that describe the presence of conditions that increase an individual risk for heart disease and other disorders such as diabetes and stroke, and its occurrence is in the rise in residents of North-Western Nigeria, The aim of this study was to investigate the prevalence and determinants of metabolic syndrome in respondents from Katsina senatorial zone attending the Out-patient Department of the Federal Medical Center Katsina, Katsina State, Nigeria. A total of 211 (male 109; female 102) respondents aged 10 to 80 years were recruited for the study. Anthropometric parameters and blood pressure of respondents were determined using standard methods; serum lipid profile was determined using enzymatic methods. From the results, 11.8% of the male respondents were under weight, 51.6% were within the normal range, 25.4% of the study male population was found to be overweight and 12.8% were obese. 13.5% of the female respondents were under weight, 51.9% were within the normal range, 22.8% of the study female population was found to be overweight and 13.1% were obese. All serum lipids measurements with the exception of serum LDL-C concentration correlated positively with age. There were no statistically significant differences between the frequencies of total cholesterol, HDL, LDL and TG between the male and female respondents. The most common form of Dyslipidemia in the male and female respondents is low HDL-C. BMI, SBP, LDL-C, HDLC and TRIG were associated metabolic syndrome with the association being significant for SBP and HDL-C (0.05; 0.03) in the male respondent, in the female respondents BMI, SBP, DBP, LDL-C and TRIG were associated metabolic syndrome with the association being significant for SBP, DBP, and TRIG (0.04; 0.04; 0.04) respectively. The prevalence of metabolic syndrome was highest in male (31.75%) than in the female respondents (28.33).The prevalence of metabolic syndrome is common in the population under study with the male respondents having the highest prevalence. A robust and well designed intervention program by concerned authorities is desirable to address complications of the risk factors for metabolic syndrome in the population.

Keywords: Obesity, Metabolic, hypertension, Lipid, Katsina

1. INTRODUCTION

Worldwide there is an increase in reported cases of metabolic syndrome (MS) prevalence in populations (1) Metabolic syndrome (MS) symbolizes the co-occurrence of risk factors that increase one's risk for heart disease and other disorders such as diabetes and stroke. MS is a clustering of metabolic disturbances such as abdominal obesity, high blood pressure (BP), increased blood glucose level, and dyslipidemia, all of which increases the risk of cardiovascular disease (CVD) and type 2 diabetes mellitus (2). The worldwide increase in

the prevalence of MS cannot be fully explained by lifestyle factors such as sedentary behavior and caloric intake alone. Exposures to environmental toxicants, such as heavy metals, have been implicated (3). It has been reported that metabolic syndrome is common in residents of North-Western Nigeria, commoner in the females than males (4).

The study was aimed at determining the prevalence of metabolic syndrome and dyslipidemia and their components among selected male and female subjects from Katsina Senatorial zone attending the general out-patient Department of the Federal Medical Center, Katsina, Katsina State, Nigeria, a referral Hospital. Results from the study will provide novel approaches to preventing and managing non-communicable diseases.

2. MATERIAL AND METHODS

2.1 STUDY AREA

The study was conducted during 2016-2017 in Katsina State, Nigeria located between latitude 12^o15'N and longitude of 7^o30'E in the North West Zone of Nigeria, with an area of 24,192km² (9,341 sq meters) (5). The State has a rainy season that begins in April and ends in October, while the dry season starts in November and last till March. The average annual rainfall, temperature, and relative humidity of Katsina State are 1,312 mm, 27.3°C and 50.2%, respectively.

2.2 SAMPLING TECHNIQUE

In this cross sectional study the Federal Medical Center Katsina a tertiary referral hospital was chosen. The method of Yusuf et al., (6) was adopted in selection of subjects that participate in the study, where with the use of stratified sampling technique; proportionate allocation was given to each age group to make up the required sample size depending on the population of the subjects in each age group.

All out patients within the study age groups were taken into. However, in each age group, systematic sampling method was employed in which the subjects in each age group were given numbers serially according to their age grouping, thereby giving each subject an equal chance. The random number table was used to select the first number and thereafter subjects were picked at regular interval (sample interval) so as to meet the sample size requirement in each group. The sample interval was determined by dividing the total number of subjects in the age groups by the sample size:

$$2500/250 = 10.$$

A total of 250 out patients were selected for the study, out of which 211 have complete data.

Only those with complete data taken were included in the study analysis. The residential township of each subject was accessed from the database of the Health Evaluation Centre of the Federal Medical Centre. Of the 211 individuals who were enrolled in the final analysis, 109 are male and 102 are female. All data were anonymised, and the ethical approval of the study protocol was given by the Ethical Committee Katsina State Ministry of Health.

2.3 ANTHROPOMETRIC AND BLOOD PRESSURE MEASUREMENTS

The weight of each subject was measured, being bare footed and in light clothing, using WEYLUX weighing scale, model 424J; Sliding Beam Column Scale, (Short Pillar with height

of 560mm). Measurements were recorded to the nearest one (1) kilogram (Kg). Height of the subjects was measured using ACCUSTAT Ross Stadiometer, 44817 (Genentech Incorporated). The measurements were recorded to the nearest 1cm. Waist measurements were taken with the use of a non-stretch metallic tape with a narrow blade and a blank lead-in. Waist circumference was measured on bare skin in the narrowest part of the abdomen between the ribs and iliac crest. The BMI (for age groups 10-80 years) was then computed using the standard formula [BMI=weight (kg)/height (m²)] (6). Blood pressure of the subjects was measured using the method of Wesseling (7). Aneroid Sphygmomanometer model OGO2, Kenzmedico Company Limited, Saitama Japan was used, with each of the participant seated in a quiet place on a chair with a back support.

2.4 BIOCHEMICAL MEASUREMENTS

Blood samples were taken from each subject by veni puncture after which the samples were allowed to clot at room temperature before being centrifuged in order to separate the serum and then frozen before the analysis.

2.4.2 LIPID PROFILE

The Serum TC concentration was measured by the end point colorimetric method of Allain et al., (8) with the use of T60 spectrophotometer and test kits obtainable from Spectrum diagnostics. In this method, the cholesterol was determined after enzymatic hydrolysis and oxidation. The indicator quinoneimine was formed from the reaction between hydrogen peroxide and 4-aminoantipyrine in the presence of phenol and peroxidase.

The serum HDL-Cholesterol of the male and female subjects was measured with a T60 spectrophotometer with test kits obtained from Spectrum diagnostics. In this method, low density lipoprotein and chylomicron fractions are precipitated quantitatively by addition of phosphotungstic acid in the presence of Mg₂⁺ ions. After centrifugation, the cholesterol concentration in the HDL fraction which remained in the supernatant was then determined (9).

The serum LDL-Cholesterol concentration of the male and female subjects was measured with a T60 spectrophotometer with test kits obtainable from Spectrum diagnostics. In this method, low density lipoproteins are precipitated by heparin at their iso-electric point (pH=5.04). After centrifugation, the HDL cholesterol and VLDL remained in the supernatant. The cholesterol concentration in the LDL fraction was determined by enzymatic method (10).

The serum Triacyl glycerol concentration of the male and female subjects was measured with a T60 spectrophotometer with test kits obtainable from Spectrum diagnostics. In this method, the triglycerides are determined after enzymatic hydrolysis with lipases. The indicator is a quinoneimine formed from hydrogen peroxide, 4-amino phenazone and 4-chlorophenol under the catalytic influence of peroxidase (11).

The serum TC, LDL-C and TG were considered elevated if they were ≥5.2 mmol/L, 3.4 mmol/L and 1.7 mmol/L, respectively, and HDL-C was taken as low if it was <1.0 mmol/L (12). Dyslipidemia was considered when at least one lipid disorder (hypertriglyceridaemia, hypercholesterolemia and or low HDL-C) was present.

2.5 STATISTICAL ANALYSIS

The results were presented as mean ± standard deviation. Relationship and association between parameters were explored using ANOVA and student's t-test. Multivariate logistic

analysis stratified by gender was performed to explore the association between measured parameters and metabolic syndrome.

3. RESULTS AND DISCUSSION

3.1 RESULTS

3.1.1 ANTHROPOMETRIC PARAMETERS, BLOOD PRESSURE AND SERUM LIPID PROFILE IN MALE AND FEMALE SUBJECTS

Being the same for both sexes and for all ages of adults, the BMI is the most useful population-level measure of overweight and obesity (13) More than 1.9 billion adults, 18 years and older, were overweight in the year 2016. Out of which over 650 million were obese (13). The BMI of the male subjects are presented in tables 1. From the results, 11.8% of the male subjects were under weight, 51.6% were within the normal range, 25.4% of the study male population was found to be overweight and 12.8% were obese. The BMI of the female subjects are presented in tables 2. From the results, 13.5% of the female subjects from the three senatorial zones were under weight, 51.9% were within the normal range, 22.8% of the study female population was found to be overweight and 13.1% were obese. Tables 1 and 2 also represent the mean serum lipid levels in male and female subjects. Mean TC level is highest in female subjects (3.73 mmol/l) compared to the male subjects (3.72 mmol/l), LDL-cholesterol is higher in male subjects (2.05 mmol/l) compared to the females (2.03 mmol/l), HDL-cholesterol is highest in female subjects (1.09 mmol/l) compared to the male subjects (0.91 mmol/l) and Triglycerides is highest in the male subjects (1.24 mmol/l) compared to the female subjects (1.16 mmol/l). But all the differences failed to achieve significance. The pattern of mean serum lipids concentration is similar for the male and female subjects. Mean TC levels are significantly higher ($p < 0.05$) in the age groups 71-80years, 61-70years and 51-60years than in the age groups 10-20 years, 21-30 years and 41-50 years. The male subjects in the age groups 61-70, 31-40, 51-60 and 71-80 have a significantly higher ($P < 0.05$) LDL-C values than the age groups 41-50, 21-30 and 10-20. The HDL-C values are significantly higher ($p < 0.05$) in the age groups 41-50, 71-80, 61-70 and 31-40 than the age groups 10-20, 21-30 and 51-60. The age groups 31-40, 71-80, 61-70 and 51-60 have a significantly higher ($p < 0.05$) mean TRIG values compared to the age groups 10-20, 41-50 and 21-30. There were no statistically significant differences between the frequencies of total cholesterol, HDL, LDL and TG between the male and female subjects in the study.

Table 1 Mean BMI, Waist Circumference (cm), Systolic and Diastolic Blood Pressure (mmHg), and Serum Lipid Profile (mmol/L) in Katsina Senatorial Zone Male Subjects

Age (years)	BMI	WC	Systolic Blood Pressure	Diastolic Blood Pressure	TC	LDL-C	HDL-C	TRIG
10-20 n=19	18.5955 ±2.1570	73.8400± 8.4050	114.6700 ± 9.9620	75.0000 ±10.6000	3.0370 ^b ± 0.2630	1.2100 ^d ± 0.1650	1.0250 ±0.0380	1.1170 ±0.1560
21-30 n=16	22.1026 ±1.1320	75.2000± 6.2460	127.2100 ±11.6700	80.6000 ± 2.5010	3.4580 ^b ± 0.6200	1.4300 ^d ± 0.2750	0.9520 ±0.0670	1.2230 ±0.0800
31-40 n=15	24.1989 ±1.1350	84.3800± 10.5700	134.6700 ± 9.7920	82.0000 ± 2.8500	3.4500± 0.8980	2.2950 ^c ± 0.6040	1.0430 ±0.1200	1.3140 ±0.1620
41-50 n=16	25.6410 ±0.8326	88.5200± 5.2800	132.3700 ±10.6050	81.3530 ± 3.6220	3.6840 ^b ± 1.1530	1.8730 ^d ± 0.6760	1.1130 ±0.1810	1.2200 ±0.1040
51-60 n=15	26.3740 ±0.4568	88.7600± 9.6300	136.7900 ± 7.9130	84.2500 ± 4.7120	3.9080 ^a ± 0.7990	2.2580 ^c ± 0.5970	1.0180 ±0.0810	1.2700 ±0.0630
61-70	25.7310	85.9300±	141.0500	82.1330	4.1840 ^a ±	2.6280 ^c ±	1.0680	1.2700

n=17	±0.1637	10.6300	± 7.3370	± 3.3780	1.3140	0.9260	±0.1820	±0.0750
71-80	24.9673	83.6100±	142.8600	86.2500	4.3420 ^a ±	2.0530 ^c ±	1.0700	1.2880
n=11	±0.4218	3.5500	± 5.0210	± 5.7230	1.3000	0.2660	±0.2120	±0.0190

Values are given as Mean ± Standard Deviation. KEY: BMI= Body Mass Index, WC= Waist Circumference, TC =Total Cholesterol, LDL-C= Low Density Lipoprotein Cholesterol, HDL-C=High Density Lipoprotein Cholesterol, TRIG= Triglyceride.

Table 2 Mean BMI, Waist Circumference (cm), Systolic and Diastolic Blood Pressure (mmHg), and Serum Lipid Profile (mmol/L) in Katsina Senatorial Zone Female Subjects

Age (years)	BMI	WC	Systolic Blood Pressure	Diastolic Blood Pressure	TC	LDL-C	HDL-C	TRIG
10-20 n=21	17.7970± 2.5560	69.6400± 3.5600	107.6400± 8.3350	74.9410 ± 7.2930	3.2130± 0.1830	1.3410± 0.5120	0.9920± 0.0580	1.1210± 0.1540
21-30 n=14	22.2790± 0.6270	71.4200± 7.2500	117.8000 ± 4.4750	78.7860 ± 5.2650	3.5180± 0.7260	1.7620± 0.3210	1.1180± 0.2610	1.1020± 0.0410
31-40 n=16	25.1960± 1.7830	73.7800± 5.3300	120.3300 ± 3.4160	79.5000 ± 3.8990	3.7740± 0.9530	1.7430± 0.4360	1.2250± 0.2550	1.1280± 0.1560
41-50 n=14	26.3280± 0.6640	85.2600± 4.2800	134.5000± 10.0660	80.3640 ± 2.0140	3.5120± 0.8210	1.7480± 0.5860	1.0460± 0.1750	1.1160± 0.1130
51-60 n=15	26.9350± 0.2850	83.6600± 6.2100	141.8200± 10.3820	82.3890 ± 3.1270	3.8320± 0.7420	2.2250± 0.5370	1.1140± 0.8350	1.2530± 0.0480
61-70 n=12	26.5930± 0.4210	83.1700± 5.9600	143.1800 ± 9.3420	82.5560 ± 3.1660	4.2160± 0.9150	2.7750± 0.3140	1.1130± 0.6350	1.2200± 0.9230
71-80 n=10	26.1290± 0.7290	79.2600± 4.5500	141.9400± 18.4700	88.0000 ± 4.6900	4.0140± 0.8430	2.6400± 0.9520	1.0430± 0.0390	1.1620± 0.0720

Values are given as Mean ± Standard Deviation. KEY: BMI= Body Mass Index, WC= Waist Circumference, TC =Total Cholesterol, LDL-C= Low Density Lipoprotein Cholesterol, HDL-C=High Density Lipoprotein Cholesterol, TRIG= Triglyceride.

3.1.2 DETERMINANTS OF METABOLIC SYNDROME

The determinants of metabolic syndrome in the male and female subjects are shown in table 3 with decreased HDL-C being the most common component of metabolic syndrome in both the male and female subjects, while elevated TRIG were the least. The sequence for the determinants of metabolic syndrome occurs in the order: ↓HDL > Hypertension > ↑WC > ↑TRIG. From the table as was observed for the subjects. The highest hypertensive subjects (40.37%) were observed in the male subjects and the least (37.25%) was observed for the female subjects. For the subjects with an elevated waist circumference the male subjects have the highest (31.19%) while the female has the least (28.43%). The male subjects

recorded the highest number of subjects with elevated triglycerides (16.51%) and the female subjects have the least (15.69%). The highest decreased HDL-C was in the male subjects (41.28%) while the least was in the female subjects (37.26%). The prevalence of metabolic syndrome was highest in male (31.75%) than in the female subjects (28.33%).

Table 3 Determinants of Metabolic Syndrome (%) in Male and Female Subjects

Determinant	Number (%)	
	Male (n=109)	Female (n=102)
Hypertension	44(40.37)	38(37.25)
↑WC	34(31.19)	29(28.43)
↑TRIG	18(16.51)	16(15.69)
↓HDL-C	45(41.28)	38(37.26)

3.1.3 DYSLIPIDEMIA

The most common form of Dyslipidemia in the male and female subjects is low HDL-C (Table 4), with the male subjects exhibiting highest percentage decrease (41.28%) and the female subjects the least (37.26%). The male subjects from Katsina have the highest percentage elevated triglyceride (16.51%), while the female subjects have the least (15.69%). The percentage elevated LDL-C is highest in female subjects (10.78%) and lowest in male subjects (7.34%). Elevated TC was highest in female subjects (18.63%) and lowest in the male subjects (12.84%).

Table 4 Dyslipidemia (%) in Male and Female Subjects

Dyslipidemia	Number (%)	
	Male (n=109)	Female (n=102)
↑TC	14(12.84)	19(18.63)
↑LDL-C	8(7.34)	11(10.78)
↑TRIG	18(16.51)	16(15.69)
↓HDL-C	45(41.28)	38(37.26)

Table 5 represents the results of multivariate logistic regression analysis for metabolic syndrome risk Factors in the subjects stratified by gender. From the table BMI, SBP, LDL-C, HDL-C and TRIG were associated metabolic syndrome with the association being significant for SBP and HDL-C (0.05; 0.03) in the male subjects. From the table BMI, SBP, DBP, LDL-C and TRIG were associated metabolic syndrome with the association being significant for SBP, DBP and TRIG (0.04; 0.04; 0.04) respectively in the female subjects

Table 5 Multivariate Logistic Regression Analysis for Metabolic Syndrome Risk Factors in Subjects Stratified By Gender

Risk Factor	Male			Female		
	OR	95% CI	Pvalue	OR	95% CI	Pvalue
BMI	1.05	23.50- 24.40	0.54	1.05	23.80- 25.10	0.56
WC	0.65	86.60- 90.50	0.27	0.82	77.40- 78.70	0.77
SBP	1.21	132.00- 133.00	0.05	1.13	129.00- 130.00	0.04
DBP	0.83	81.20- 82.20	0.48	1.24	80.30- 81.60	0.04
LDL-C	1.11	1.58- 2.52	0.15	1.11	1.28-2.94	0.86

HDL-C	1.13	0.44- 1.39	0.03	0.85	0.47-1.70	0.14
TRIG	1.38	0.77- 1.72	0.56	1.19	0.54-1.80	0.04

Key: OR=Odd Ratio, BMI= Body Mass Index, WC= Waist Circumference, SBP= Systolic Blood Pressure, DBP= Diastolic Blood Pressure, LDL-C= Low Density Lipoprotein Cholesterol, HDL-C= High Density Lipoprotein Cholesterol, TRIG= Triglyceride

3.2 DISCUSSION

The results of the prevalence of anthropometry (Underweight, normal weight, overweight and obesity) in the male and female subjects differs from the prevalence reported for traders from Bodija Ibadan, Nigeria (14) with the subjects in the present study having a comparatively lower overweight and obese but a higher normal weight values. The prevalence of obesity in the present study is similar to the world prevalence values for adults as reported by Planchart et al., (3). The values are also lower than the overweight and obesity prevalence for men (70.8%) and women (77.7%) African-Americans (15), the prevalence in Sokoto reported by Sabir et al., (4), obesity prevalence (81% men; 93% women) in Ghanaian and Nigerian born African-Americans (16) and obesity prevalence (17.2%) in rural workers from Idemili South Local Government Area, Southeast Nigeria (17).

The waist circumference mean values reported in the present study (Men range= 73.84-88.52 cm; Women range= 69.64-85.26 cm) is higher than the mean value reported (Men= 74.78 cm; Women= 71.72 cm) in a study conducted in Vietnam (18) and lower than the mean values (108.3, 102.1, 115.7) reported for post menopausal women with diabetes in Sydney Australia (19), it is also falls within the range of the mean value (86.2) reported for men and women subjects in a study conducted in Sokoto city, Nigeria (4).

The prevalence of high blood pressure in the male (40.37%) and female (37.25%) subjects in this study is lower when compared to the prevalence reported (46.1%) for subjects from Sokoto metropolis, Sokoto state, Nigeria (4), but the prevalence is higher than the prevalence of elevated blood pressure (19.3%) reported by Okaka and Eiya (20) in a rural community in Southern Nigeria, the prevalence (16%) reported for Ghanaian and Nigerian born African-Americans (16) and the prevalence (23.1%) reported by Oguoma et al., (12) in a Nigerian population with impaired fasting blood glucose level and diabetes mellitus.

The mean serum lipid levels in the male and female subject's falls within the values considered acceptable for African Americans (21). The levels of serum lipid values were low when compared to the general US population (21). The values are also lower when compared to serum lipid values reported by Okaka and Eiya (20) in a rural population from southern Nigeria, the results of Akintunde and Salawu (22) on serum lipids concentration of Staff of Ladoke Akintola University Ogbomosho, Nigeria, Tehran lipid and glucose study (23), results of lipid profile of Ghanaian and Nigerian-born African-Americans (16) and the lipid profile reported by Bently and Rotimi (15) for African-Americans. The trend of relative low serum lipid profile observed in this study is similar to what was reported in lipids profile studies conducted in Northern Nigeria (24; 25; 26; 27; 28) and the results of a study conducted a rural population from south-western Uganda (20). All serum lipids measurements with the exception of serum LDL-C concentration correlated positively with age. This finding is in line with the Bogalusa heart study (30) and the Framingham Offspring Study (31) that shows positive correlation between lipid profile and age and the study of Sairam et al., (32) conducted in India, but differs from the Honolulu Heart Program study (33), and the findings of Adrabo et al., (34), whose study brought into light the hypothesis

that there is an increase in the levels of cholesterol with age until the sixties but the level start to decrease beyond that age.

The prevalence of low HDL-C exhibited by the male female subjects is similar to what was reported in staff of Ladoke Akintola University of Technology, Ogbomosho (22) and from Sokoto, Nigeria (4) and in a Nigerian population with impaired fasting blood glucose level and diabetes mellitus (12). But the prevalence is higher than the prevalence of low HDL-C (16.6%) reported for African-Americans (15), but lower than the prevalence (71.3%) reported in a rural population in South-western Uganda (29).

Prevalence of elevated TRIG in the male (16.51%) and female (15.69%) subjects in the present study is lower compared to the prevalence of 38.0% reported for rural workers from Idemili South Local Government Area, Southeast Nigeria (17).

The Low high-density lipoprotein-cholesterol level as the most prevalent dyslipidemia in the present study is similar to what was reported among Northern Mexican adolescents (35), but the result is in contrast to the reported abdominal obesity, raised diastolic blood pressure and hyper triglyceridemia as determinants of metabolic syndrome in rural workers from Idemili South Local Government Area, Southeast Nigeria (17).

The male and female subjects may be predisposed to atrial fibrillation (AF), as the low cholesterol levels and high cholesterol variability shown by the subjects were earlier associated with a higher risk of AF development (36).

4. CONCLUSION

The prevalence of metabolic syndrome is common in the population under study with the male subjects having the highest prevalence. A robust and well designed intervention program by concerned authorities is desirable to address complications of the risk factors for metabolic syndrome.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee (Katsina HREC assigned number: MOH/ADM/SUB/1174/1/190) and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

- 1 Lee JK, Ryoo JH, Choi JH, Park SK,. Serum Uric Acid Level and the Incidence of Metabolic Syndrome in Middle-aged Korean Men: A 5-Year Follow-up Study. *J Prev Med Public Health*, 2014; 47(6): 317–326.
- 2 Osgood K, Krakoff J, Thearle M. Serum uric acid predicts both current and future components of the metabolic syndrome. *Metab Syndr Relat Disord.*, 2013; 11(3):157–162

3 Planchart A, Green A, Hoyo C, Mattingly CJ,. Heavy Metal Exposure and Metabolic Syndrome: Evidence from Human and Model System Studies. *Current Environmental Health Reports*, 2018; <https://doi.org/10.1007/s40572-018-0182-3>

4 Sabir AA, Jimoh A, Iwuala SO, Isezuo SA, Bilbis LS, Aminu KU, Abubakar SA, Saidu Y Metabolic syndrome in urban city of North-Western Nigeria: prevalence and determinants *Pan Afr Med J*. 2016; 23: 19

5 Katsina State investor's Katsina State Government handbook, Yaliam Press Ltd 2016: 12-15

6 Yusuf SM, Mijinyawa MS, Musa BM, Gezawa ID, Uloko AE,. Overweight and obesity among adolescents in Kano, *Nigeria Journal Metabolic syndrome*, 2013; 2(1) DOI: 104172/2167-0943.1000126.

7 Wesseling KH, A century of non-invasive arterial pressure measurement: From Marey to Panez and Finapres. *Homeostasis*, 1995; 36: 50-66.

8 Allain CC, Poon LS, Chan CSG, Richmond W. Enzymatic determination of total cholesterol in serum. *Clinical Chemistry*, 1974; 20:470-475.

9 Friedewald WT, Levy RI, Fredrickson DS,. Estimation of the concentration of LDL cholesterol in plasma without use of preparative ultracentrifuge. *Clinical Chemistry* 1972; 499-502.

10 Wicland H, Seidel D, A simple method for the precipitation of low density lipoproteins. *Journal of Lipid Research*, 1983; 24:904-909

11 Tiez NW. Clinical guide to laboratory tests, 2nd Edition 1990, W. B. Saunders Company. Philadelphia, U. S. A.; 554-556

12 Oguoma VM, Nwose EU, Ulasi II, Akintunde AA, Chukwukelu EE, Bwititi PT, *et al*. Cardiovascular disease risk factors in a Nigerian population with impaired fasting blood glucose level and diabetes mellitus *BMC Public Health BMC series* 2017; 17:36

13World Health Organization (2018) Preventing noncommunicable diseases (NCDs) by reducing environmental risk factors <http://apps.who.int/iris/bitstream/handle/10665/258796/WHO-FWC-EPE-17.01-eng.pdf;jsessionid=7368E2B81AC7C9CE8B40C38C7EED1F9E>

14 Charles-Davies MA, Arinola OG, Fasanmade AA, Olaniyi JA, Oyewole OE, Owolabi MO, *et al*. Indices of Metabolic Syndrome in 534 Apparently Healthy Nigerian Traders. *Journal of US-China Medical Science*, 2012; Volume 9, No. 2 (Serial No. 87), pp. 91–100

15 Bentley AR, Rotimi, CN, Interethnic Differences in Serum Lipids: Implications for Cardiometabolic Disease Risk in African Ancestry Populations. *Global Heart*, 2017; <https://doi.org/10.1016/j.gheart.2017.01.011>

16 Commodore-Mensah Y, Hill M, Allen J, Cooper LA, Blumenthal R, Agyemang C, *et al*. Sex Differences in Cardiovascular Disease Risk of Ghanaian- and Nigerian-Born West African Immigrants in the United States: The Afro-Cardiac Study. *Journal of the American Heart Association*, 2016; 5: e002385

17 Ayogu RNB, Nwajuaku C, Udentia EA,. Components and Risk Factors of Metabolic Syndrome among Rural Nigerian Workers. *Niger Med. J.* 2019; 60(2): 53–61. doi: 10.4103/nmj.NMJ_53_19

18 Tran NTT, Blizzard CL, Luong KN, Truong NLV, Tran BQ, Otahal P, *et al.* The importance of waist circumference and body mass index in cross-sectional relationships with risk of cardiovascular disease in Vietnam. *PLoS ONE*, 2018 13(5): e0198202. <https://doi.org/10.1371/journal.pone.0198202>

19 Seimon RV, Wild-Taylor AL, Gibson AA, Harper C, McClintock S, Hamish A. *et al.* Less Waste on Waist Measurements: Determination of Optimal Waist Circumference Measurement Site to Predict Visceral Adipose Tissue in Postmenopausal Women with Obesity. *Nutrients*, 2018; 10, 239; doi: 10.3390/nu10020239

20 Okaka EI, Eiya BO,. Prevalence and pattern of dyslipidemia in a rural community in Southern Nigeria. *Afr J Med Health Sci.*, 2013; 12:82-6

21 Rifai NI, Warmick GR, Dominiczak MH. Handbook of lipoprotein testing 1996; Washington DC Association for Clinical Chemistry Press.

22 Akintunde AA, Salawu AA, Opadijo OG,. Prevalence of traditional cardiovascular risk factors among staff of Ladoke Akintola University of Technology Ogbomoso, Nigeria. *Niger J Clin Pract.*, 2014; 17: 750-5

23 Kalantari S. Childhood cardiovascular risk factors, a predictor of late adolescent overweight. *Adv Biomed Res.*, 2016; 5:56

24 Glew RH, Margaret W, Carole AC, Samuel MC, Michael C, Selina NO, *et al.* Cardiovascular disease risk factors and diet of Fulani pastoralist of Northern Nigeria. *American Journal of Clinical Nutrition*, 2001; 7(6):730-736.

25 Atiku MK, Yusuf AB. Marital status and occupation versus serum total cholesterol and HDL-cholesterol levels in healthy adults from Kano metropolis, Nigeria. *Bayero Journal of Pure and Applied Sciences*, 2011; 4(1):110-111

26 Hafiz A, Ibrahim AY, Atiku MK,. Nutritional and health status of adolescents from selected secondary schools in Katsina State Nigeria *Nigerian Journal of Nutritional Sciences*, 2012; 2(33):11-14

27 Shuaibu L, Ibrahim AY, Suleiman S. Body mass index and lipid profile: A case study of female school going adolescents in Katsina metropolis, Nigeria. *British Journal of Applied Science and Technology*, 2014; 4(25):3665-3677.

28 Yaradua AI, Shuaibu L, Madaha M,. A Comparative Analysis of Lipid Profile among Rural and Urban School Going Male Adolescents in Katsina State Nigeria. *International Journal of Biochemistry Research and Review*, 2015; 5(4): 250-258

29 Asiki G, Murphy GAV, Baisley K, Nsubuga RN, Karabarinde A, Newton R, *et al.* Prevalence of Dyslipidaemia and Associated Risk Factors in a Rural Population in South-Western Uganda: A Community Based Survey. *PLoS ONE*, 2015; 10(5): e0126166. <https://doi.org/10.1371/journal.pone.0126166>

30 Freedman DS, Dutz WA, Srinivasan SR, Berenson GS (1999) The relationship of overweight to cardiovascular risk factors among children and adolescents. The Bogalusa Heart Study, *Pediatrics*; 163 (6):175-181

31 Schaefer EJ, Lamon-Fava S, Cohn SD, Schaefer MM, Ordovas JM, Castelli WP, Wilson PW. Effects of age, gender, and menopausal status on plasma low density lipoprotein cholesterol and apolipoprotein B levels in the Framingham Offspring Study. *J Lipid Res.* 1994 May;35(5):779-92.

32 Sairam S, Domalapalli S, Muthu S, Swaminathan J, Ramesh VA, Sekhar I, et al. Hematological and Biochemical Parameters in Apparently Healthy Indian Population: Defining Reference Intervals. *Indian J Clin Biochem.*, 2014; 29(3): 290–297

33 Burchfiel CM, Abbott RD, Sharp DS, Curb JD, Rodriguez BL, Yano K. Distribution and correlates of lipids and lipoproteins in elderly Japanese-American men. The Honolulu Heart Program. *Arterioscler Thromb Vasc Biol.*, 1996; 16(11):1356-64.

34 Marhoum TA, Abdrabo AA, Lutfi MF. Effects of age and gender on serum lipid profile in over 55 years-old apparently healthy Sudanese individuals. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 2013; 3(19) 2013, 10-14.

35 Bibiloni MD, Salas R, De la Garza YE, Villarreal JZ, Sureda A, Tur JA,. Serum Lipid Profile, Prevalence of Dyslipidaemia, and Associated Risk Factors Among Northern Mexican Adolescents. *J Pediatr Gastroenterol Nutr.*, 2016; 63(5): 544–549.

36 Lee HJ, Lee SR, Choi EK, Han KD, Oh S, Low Lipid Levels and High Variability are Associated with the Risk of New-Onset Atrial Fibrillation. *J. Am. Heart Assoc*, 2019; 8:e012771. DOI: 10.1161/JAHA.119.012771

ABBREVIATIONS

MS = Metabolic Syndrome

HREC = Health Research Ethics Committee

ANOVA = Analysis of variance

WC = Waist circumference

SBP = Systolic blood pressure

DBP = Diastolic blood pressure

BMI = Body mass index

TC = Total Cholesterol

LDL-C = Low density lipoprotein Cholesterol

HDL-C = High density lipoprotein Cholesterol

TRIG = Triglyceride

UNDER PEER REVIEW