

Original Research Article

Evaluation of the Phytochemicals, Nutritional and Anti-nutritional Compositions of Fresh, Sprouted and Toasted *Citrullus lanatus* (Watermelon) Seeds

ABSTRACT

Aims: To evaluate the Phytochemicals Nutritional and Anti-nutritional Compositions of Fresh, Sprouted and Toasted *Citrullus lanatus* (Watermelon) Seeds

Study Design: Randomized experimental design.

Place and Duration of Study: Department of Biochemistry and Molecular Biology Laboratory of Nasarawa State University, Keffi, Nasarawa State, Nigeria between January and August, 2018.

Methodology: The phytochemicals, proximate, amino acids, microminerals and antinutrients compositions were determined following standard laboratory procedures. Data were analyzed using one-way ANOVA followed by LSD and Duncan test for levels of significance.

Results: Phytochemical analysis revealed the presence of Tannins, Saponins, Flavonoids, Alkaloids, Cardiac glycosides, Terpenoids in the three samples but anthroquinones was present only in the fresh sample but absent in sprouted and toasted watermelon seeds. Saponins (2.15 ± 0.07 mg/100g), Tannins (40.63 ± 0.13 mg/100g) Cyanogenic glycosides (3.55 ± 0.35 mg/100g). There was no significant ($P > 0.05$) difference in the concentrations of the phytochemicals in all the samples except for cyanogenic glycosides which was significantly ($P < 0.05$) higher in the fresh sample. Percentage protein (24.87 ± 0.48) was significantly ($P < 0.05$) higher in the sprouted sample, ash (%) (2.63 ± 0.11) was significantly ($P < 0.05$) lower in the sprouted, fibre (20.64 ± 0.23) and moisture (%) (2.88 ± 0.19) was significantly ($P < 0.05$) lower in the toasted and carbohydrates (%) (9.81 ± 1.27) was significantly ($P < 0.05$) higher in the sprouted. Results obtained for Amino acids and mineral analyses showed no significant ($P > 0.05$) difference in the three samples and Phytates ($0.31 \pm 0.01\%$) were significantly ($P < 0.05$) higher in the fresh sample while oxalates ($0.85 \pm 0.01\%$) was significantly ($P < 0.05$) higher in the sprouted sample.

Conclusion: Fresh, sprouted and toasted watermelon seeds may possess nutritional and health benefit by the virtue of their richness in phytochemicals, microminerals and amino acids.

Keywords: Amino acids, Bioactive compounds, proximate analysis, microminerals, antinutrients,

INTRODUCTION

Plant seeds are generally good source of food for animals as well as humans, since they contain nutrients necessary for growth, including healthy fats such as omega fats [1]. Seeds of many plants in the *Cucurbitaceae* are rich in oil and proteins, and although none of these oils have been used on an industrial scale, many of the oils are used as cooking oil in some countries in Africa and the Middle East [2]. Watermelon seeds have shown the potential for use in the food industry as they remain intact after removing the pulp and peel [3]. The seeds have also the ability to store well, with both oil and fatty acid content being found to be stable after six months in storage [4]. According to [5], the seeds can be utilized successfully as a source of edible oils for human consumption, and their oil might be an acceptable substitute for highly unsaturated oils [5]. The seeds also contain lipids of nutritional interest, with high concentrations of unsaturated fatty acids, including a high concentration of phytosterols, particularly stigmaterol and β -sitosterol [3]. The use of watermelon seeds as a food source appears to be justified by their reported nutritional value. The dry seeds of *Citrullus lanatus* have been reported on an average to contain 22g of proteins, 30g of fat and 11g of carbohydrate per 100g sample [6]. The seeds are highly nutritive and contain large amounts of proteins and many beneficial minerals such as magnesium, calcium, potassium, iron, phosphorus and zinc [7].

The plant, *Citrullus lanatus* commonly known as watermelon belongs to the *Cucurbitaceae* family and is widely distributed in Africa and Asia, but the exact origin of the crop is subject to strong debate. It is an annual plant which could be cultivated, semi-domesticated and in wild forms [8]. According to [9], the crop originated from southern Africa and occurs naturally in South Africa, Namibia, Botswana, Zimbabwe, Mozambique, Zambia and Malawi. It is also cultivated and thrives effectively in warmer parts of the world [10], and is also widely distributed in tropical and subtropical areas [8], where the crop is well adapted to the prevailing arid and unpredictable climatic conditions of the regions [10];[8]. In Africa, the *Cucumis sativas* accounts for 5.4% of the harvested area devoted to vegetable cultivation and 6.8% worldwide [11]. The fruit is a source of multiple minerals, vitamins, and proteins that are present in the epicarp, pulp and seeds [12]. The plant, like many other cucurbits such as, gourds that include cucumbers (*Cucumis sativas*), squashes (*Cucubita moschata*), luffas (*Luffa sp*) and melons (*Cucumis melo*) are among the economically most important vegetable crops in the world [13].

Therefore, this research seeks to determine and compare the phytochemicals, nutrients and anti – nutrients of the fresh, sprouted and toasted seeds of *Citrullus lanatus*.

Materials and Methods

Materials

Plant Material

The plant material used for the study was *Citrullus lanatus* (Watermelon) seeds. The fruits were purchased from Keffi Main Market, Nasarawa State, Nigeria. The samples were identified and authenticated at the Department of Plant Science and Biotechnology, Nasarawa State University, Keffi, Nasarawa State, Nigeria. The fruits were further broken open to obtain the seeds.

Methods

Sample preparation

The *Citrullus lanatus* fruits were washed with clean tap water, rinsed with distilled water and allowed to dry. The seeds of the fruits were removed by cutting open the fruit with a clean knife and washed with distilled water. The seeds were then air dried for one week. The seeds were divided into three portions. The first portion was toasted in the oven at 100°C until it became golden brown before extraction. The second portion were soaked in water for seventy- two hours, then covered for forty- eight hours for sprouting. After sprouting, they were air dried for one week and thereafter grounded to powder for extraction. The fresh samples were pulverized after drying and extracted.

Extraction of the *Citrullus lanatus* seeds

The processed seeds of the *Citrullus lanatus* were pounded and soaked in 70% ethanol for 48 hours, after which it was sieved and filtered, the filtrates was concentrated using water bath to get the seed extracts which were then stored in a refrigerator until commencement of the experiment.

Qualitative Tests for Phytochemicals

The Phytochemicals screening of the extracts were carried out using standard methods as described by [14] (Sofowora, 1993; [15] Trease and Evans, 2002), to determine the presence of Alkaloids, Saponins, Flavonoids, Tannins, Anthraquinones and Cardiac Glycosides.

Quantitative Determination of Phytochemicals

The quantitative determination of all the phytochemicals were carried out according to the standard methods outlined by [16], for the concentrations of Alkaloids, Saponins, Flavonoids, Tannins, Anthraquinones and Cardiac Glycosides.

Determination of Proximate Composition

The proximate analysis of fresh, toasted and sprouted watermelon seeds were carried out using the standard method outlined by [17], to determine the moisture content, crude protein, total ash and crude fiber contents.

Determination of Mineral Composition

Potassium and sodium were estimated using a modified method of [18]. The samples were digested with perchloric acid and nitric acid and analyzed with digital flame Photometer. Phosphorus was determined by Vanado-Molybdate colorimetric method. Calcium, magnesium and iron contents were analyzed using Atomic Absorption Spectrometry.

Determination of Anti-Nutrients Components

The Saponin content of the samples was determined by double extraction gravimetric method described by [16]. Phytates was determined through phytic acid determination using the procedures described by [19]. Tannins was determined using methods described by [20]. This method was however slightly modified. The total oxalic acid acid of the powdered samples was determined by the modified method of [21].

Amino Acids Analysis

The samples were defatted, for at least 8 hours with chloroform/methanol (2:1 mixture) using soxhlet extractor apparatus, hydrolyzed with 6N HCl and evaporated in a rotary evaporator for 22 hours at 104 - 110°C. The HCl was removed after hydrolysis in vacuum and the residue then injected into and analysed in the Amino Acid Analyzer (Technicon Instrument Co. Ltd., United Kingdom). Amino acid score was calculated for each essential amino acid in a given test protein using the [22] reference pattern.

Statistical analysis

The data obtained were analyzed using one-way analysis of variance (ANOVA) with the help of IBM SPSS package, version 20.0 and the results were expressed as mean \pm standard deviation. The acceptance value of significance was $p < 0.05$ for all the results.

RESULTS AND DISCUSSION

Results

Qualitative Phytochemical composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

The results showed the presence of tannins, Saponins, flavonoids, alkaloids anthraquinones and cardiac glycosides in the fresh, sprouted and toasted watermelon seeds except for anthraquinones that was not detected in sprouted and toasted watermelon seeds (table 1).

Table 1. Qualitative Phytochemical composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Sample	Phytochemicals composition					
	Tannins	Saponins	Flavonoids	Alkaloids	Antraquinones	Cardiac glycosides
Fresh	+	+	+	+	+	+
Sprouted	+	+	+	+	-	+
Toasted	+	+	+	+	-	+

Quantitative Phytochemical composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

As shown in table 2 below, there was no significant ($p > 0.05$) difference in the tannins concentrations of the fresh ($1.20 \pm 0.1 \text{ mg/100g}$), sprouted ($0.89 \pm 0.11 \text{ mg/100g}$), and toasted ($0.62 \pm 0.02 \text{ mg/100g}$) watermelon seeds, similar results was obtained for alkaloids with mean values; $1.36 \pm 0.02 \text{ mg/100g}$, $1.09 \pm 0.06 \text{ mg/100g}$ and $1.09 \pm 0.04 \text{ mg/100g}$ for fresh, sprouted, and toasted watermelon seeds respectively. For cardiac glycosides, the mean value was significantly ($p < 0.05 \text{ mg/100g}$) higher in the fresh ($3.63 \pm 0.01 \text{ mg/100g}$) followed by sprouted ($2.1 \pm 0.09 \text{ mg/100g}$) and lastly the toasted ($0.94 \pm 0.05 \text{ mg/100g}$). There was also no significant ($p > 0.05$) difference in the flavonoids concentrations of the fresh, sprouted and toasted watermelon seeds. Saponin was significantly higher in fresh samples ($2.17 \pm 0.07 \text{ mg/100g}$).

Table 2. Quantitative Phytochemical composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds
Concentrations(mg/100g)

Phytoch.	Fresh	Sprouted	Toasted
Tannins	1.20 ± 0.1^a	0.89 ± 0.11^a	0.62 ± 0.02^a
Alkaloids	1.36 ± 0.02^b	1.09 ± 0.06^b	1.09 ± 0.04^b
Card. Gly.	3.63 ± 0.01^c	2.1 ± 0.09^d	0.94 ± 0.05^e
Flavonoids	2.89 ± 0.07^f	2.4 ± 0.07^f	1.77 ± 0.01^f
Saponin	2.15 ± 0.07^d	0.48 ± 0.03^e	0.11 ± 0.06^e

Results are expressed in Mean \pm SD (n = 3)

Mean values with different letters as superscripts across the horizontal comparisons are considered significant at $p < 0.05$

Proximate composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Table 3 shows percentage protein content of $24.87 \pm 0.48\%$ in the sprouted watermelon seeds, which is significantly ($p < 0.05$) higher compared to that of the fresh ($17.47 \pm 0.22\%$) and toasted ($15.53 \pm 0.31\%$). No significantly ($p > 0.05$) difference was observed in the percentage fat in fresh ($26.67 \pm 0.49\%$), sprouted ($29.14 \pm 0.16\%$) and toasted ($28.89 \pm 0.04\%$) watermelon seeds. For percentage ash, the value was significantly ($p < 0.05$) lower in the sprouted ($2.63 \pm 0.11\%$) compared to the fresh ($4.25 \pm 0.08\%$) and toasted ($3.56 \pm 0.10\%$). Similarly, fibre (20.64 ± 0.23) and moisture (2.88 ± 0.19) were found to be significantly ($p < 0.05$) lower in the toasted compared to the fresh and sprouted. The results obtained for carbohydrates showed percentage composition of 9.81 ± 1.27 in sprouted which was significantly ($p < 0.05$) higher compared to the fresh (5.60 ± 1.27) and toasted ($4.53 \pm 1.27\%$).

Table 3. Proximate composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Samples	Proximate compositions					
	%Protein	%Fat	%Ash	%Fibre	%Moisture	%CHO
Fresh	17.47 ± 0.22^a	26.67 ± 0.49^d	4.25 ± 0.08^e	25.87 ± 0.63^g	5.21 ± 0.16^i	5.60 ± 1.27^k
Sprouted	24.87 ± 0.48^b	29.14 ± 0.16^d	2.63 ± 0.11^f	27.83 ± 0.31^g	5.73 ± 0.21^i	9.81 ± 1.27^l
Toasted	15.53 ± 0.31^a	28.89 ± 0.04^d	3.56 ± 0.10^e	20.64 ± 0.23^h	2.88 ± 0.19^j	4.53 ± 1.27^k

Results are expressed in Mean \pm SD (n = 3)

Mean values with different letters as superscripts across the vertical comparisons are considered significant at $p < 0.05$

Amino Acids Composition of Fresh, Sprouted and Toasted *Citrullus lanatus* Seeds

The amino acid compositions of the fresh, sprouted and toasted *Citrullus lanatus* seeds showed no remarkable mean differences in all the samples analyzed as shown in table 4 below. The twenty essential amino acids were analysed and glutamic acid had the highest concentrations in all the test samples with 13.63 ± 0.23 mg/100g in the fresh, 13.02 ± 0.98 mg/100g in the sprouted and 15.30 ± 0.32 g/100g in the toasted samples followed by arginine with 9.63 ± 0.87 mg/100g in the fresh, 10.84 ± 0.098 g/100g in the sprouted and 10.60 ± 0.45 /100g in the toasted samples while tryptophan had the least concentrations of 0.76 ± 0.21 mg/100g in the fresh, 0.81 ± 0.01 mg/100g in the sprouted and 0.73 ± 0.02 mg/100g in the toasted samples.

Table 4. Amino Acids Composition of fresh, sprouted and toasted *Citrullus lanatus* seeds

Amino Acid	Concentration (mg/100g)		
	Fresh	Sprouted	Toasted
Leucine	7.00 ± 0.01^a	7.30 ± 2.03^a	6.80 ± 2.04^a
Lysine	4.35 ± 0.02^a	4.72 ± 0.06^a	2.86 ± 0.06^b
Isoleucine	3.53 ± 0.23^a	3.50 ± 0.04^a	3.44 ± 0.54^a
Phenylalanine	5.41 ± 0.32^a	5.70 ± 0.92^a	5.32 ± 0.56^a
Tryptophan	0.76 ± 0.21^a	0.81 ± 0.01^a	0.73 ± 0.02^a
Valine	3.30 ± 0.02^a	4.00 ± 1.01^b	3.51 ± 0.31^a
Methionine	2.24 ± 0.22^a	2.62 ± 0.04^a	2.46 ± 0.54^a
Proline	5.40 ± 0.11^a	4.80 ± 0.07^a	3.45 ± 0.11^b

Arginine	9.63±0.87 ^a	10.84±0.00 ^a	10.60±0.45 ^a
Tyrosine	2.41±0.34 ^a	2.06±0.05 ^a	2.41±0.05 ^a
Histidine	1.60±0.90 ^a	1.76±0.03 ^a	1.53±0.01 ^a
Cystine	1.03±0.11 ^a	1.21±0.06 ^a	0.85±0.04 ^a
Alanine	3.56±0.57 ^a	4.02±0.05 ^b	3.94±0.39 ^b
Glutamic acid	13.63±0.23 ^a	13.02±0.98 ^a	15.30±0.32 ^b
Glycine	4.63±0.45 ^a	3.90±0.99 ^b	4.46±0.60 ^a
Threonine	2.25±0.35 ^a	3.00±1.00 ^b	2.00±0.51 ^a
Serine	4.25±0.01 ^a	3.62±0.89 ^b	4.00±1.04 ^a
Aspartic acid	5.70±0.34 ^a	6.26±0.38 ^b	5.64±0.98 ^a

Results are expressed in Mean ±SD (n = 3), mean values with different letters as superscripts across the periods (i.e compared horizontally) are considered statistically significant at p<0.05

Mineral composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Results obtained for the mineral compositions of the fresh, sprouted and toasted *Citrullus lanatus* seeds showed that there was no significant difference among the samples for all the minerals analyzed as shown in table 5 below. Iron (Fe) has the highest mean concentration of 0.006±0.01(mg/100g) in the fresh, 0.006±0.001(mg/100g) in the sprouted and 0.007±0.001(mg/100g) in the toasted samples (table 5).

Table 5. Mineral composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Mineral	Mineral compositions		
	Fresh(mg/100g)	Sprouted(mg/100g)	Toasted(mg/100g)
Zinc (Zn)	0.003±0.00 ^b	0.003±0.001 ^b	0.003±0.00 ^b
Iron (Fe)	0.006±0.01 ^c	0.006±0.001 ^c	0.007±0.001 ^c

Sodium (Na)	0.002±0.001 ^d	0.003±0.003 ^d	0.002±0.001 ^d
Copper (Cu)	0.001±0.002 ^e	0.001±0.00 ^e	0.001±0.00 ^e
Manganese (Mn)	0.002±0.00 ^f	0.002±0.00 ^f	0.001±0.00 ^f

Results are expressed in Mean ± SD (n = 3)

Mean values with different letters as superscripts across the horizontal comparisons are considered significant at p < 0.05

Anti-nutrients composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

As shown in table 6 below, oxalate was found to be significantly (p < 0.05) higher in the sprouted watermelon seeds (0.85±0.01%) when compared to the fresh (0.13±0.01%) and toasted (0.07±0.01%). The concentrations of saponins, tannins, cyanogenic glycosides and phytates were significantly (p < 0.05) higher in the fresh watermelon seeds compared to the sprouted and toasted.

Table 6. Anti-nutrients composition of Fresh, Sprouted and Toasted *Citrullus lanatus* seeds

Samples	Anti-nutrients composition				
	Oxalate(%)	Saponins (%)	Tannins (mg/100g)	Cyanogenic (mg/100g)	
Phytate (%)					
Fresh	0.13±0.01 ^a	2.15±0.07 ^d	40.63± 0.13 ^f	3.55± 0.35 ^h	0.31± 0.01 ^j
Sprouted	0.85±0.01 ^b	0.48±0.03 ^e	28.84±1.09 ^g	2.96±0.11 ⁱ	0.21±0.01 ^k
Toasted	0.07±0.01 ^c	0.11±0.06 ^e	25.05±0.15 ^g	2.95±0.07 ^j	0.18±0.01 ^k

Results are expressed in Mean ± SD (n = 3), Mean values with different letters as superscripts across the vertical comparisons are considered significant at p < 0.05

DISCUSSION

In this research, we evaluated the phytochemical composition, proximate composition, mineral composition, amino acid composition and anti-nutrient composition of the fresh, sprouted and toasted seeds of *Citrullus lanatus* (watermelon). The proximate composition of sprouted *Citrullus lanatus* seeds is shown in table 3 The moisture content of the toasted seeds (2.88±0.19) was significantly lower (p<0.05) than fresh and sprouted seeds (5.21±0.61 and 5.73±0.21) respectively. Moisture content entails the amount of water molecules present in the sample. When the moisture of a sample is high, it makes the sample prone to fungal attack and vice versa. Therefore, toasting the seeds removes water and this in a way serves as a means of preserving the seed as this increases the shelf life. The moisture content for the fresh and sprouted seeds was lower than 7.10% reported by [24]. For percentage ash, the value was significantly (p < 0.05) lower in the sprouted (2.63±0.11%) compared to the fresh (4.25±0.08%) and toasted (3.56±0.10%). The low percentage of ash in the sprouted seeds could be attributed to the seeds using the minerals in the sprouting process. Ash content indicates the percentage of inorganic mineral elements present in the samples. The ash content is comparable to the ash content of *Citrullus Vulgaris* (4.83±0.4%) seed according to [25]. High mineral elements in foods enhance growth and development, and also catalyses metabolic processes in human body (Jacob *et al.*, 2015). The seeds of fresh, sprouted

and toasted water melon seeds shows a high percentage of fibre (25.87 ± 0.63 , 27.83 ± 0.31 and 20.64 ± 0.23 respectively). The percentage of fibre in the watermelon seeds is way higher than 6.40% reported for four varieties of melon [26]. Fibre is believed to reduce the level of cholesterol in the blood, reduces constipation and pile as well as decreases the chance of developing colon cancer. There was no significant ($p > 0.05$) difference observed in the percentage fat in fresh ($26.67 \pm 0.49\%$), sprouted ($29.14 \pm 0.16\%$) and toasted ($28.89 \pm 0.04\%$) watermelon seeds. These values were lower compared to the value obtained for four varieties of water melon (40.26 - 45.21%). These low values do not qualify the seeds as oil seeds. Fat is very vital since it provides the body with tremendous amount of energy. The percentage protein content of the sprouted watermelon seeds $24.87 \pm 0.48\%$, is significantly ($p < 0.05$) higher compared to fresh ($17.47 \pm 0.22\%$) and toasted ($15.53 \pm 0.31\%$). The high protein content observed in the sprouted watermelon seeds may be attributed to the mobilization of proteins required for sprouting to occur. This could also be as a result of mobilization of stored nitrogen of the seeds to aid sprouting, synthesis of enzymes or a compositional changes following the degradation of other constituents. The protein content is low compared to a range of 28.89-31.34%. The results obtained for carbohydrates showed percentage composition of 9.81 ± 1.27 in sprouted which was significantly ($p < 0.05$) higher compared to the fresh (5.60 ± 1.27) and toasted (4.53 ± 1.27). These results compared favourably with 4.52% obtained by [24]. The Carbohydrate content of water melon seeds is low and hence cannot be considered as a source of Carbohydrate.

The anti-nutritional composition of sprouted, toasted and fresh *Citrullus lanatus* seed is presented in table 6. The fresh seeds contain a significantly ($p < 0.05$) higher values of all the anti-nutrients while toasted seeds have the least values. These results indicate that processing especially toasting which involves heat significantly reduces the level of anti-nutrients in the seeds and make it nutritionally acceptable. Tannins precipitate and inhibit amino acids and protein absorption while phytates bind and slow or completely inhibit the absorption of certain minerals (such as zinc, iron, Manganese and to a lesser extent calcium) by cells.

Table 4 shows the result of the amino acid profile of fresh, sprouted and toasted water melon seeds. The results obtained showed that the samples contained all essential amino acids at moderate concentrations with no significant difference ($p > 0.05$) in the fresh, sprouted and toasted samples. The amino acid composition (mg/100g) was highest for glutamic acid (13.63 ± 0.23 , 13.02 ± 0.98 and 15.30 ± 0.32 in fresh, sprouted and toasted seeds respectively). Glutamate is a semi essential amino acid that helps nerve cells to send and receive information from other cells. This was followed by Arginine (9.63 ± 0.87 , 10.84 ± 0.01 , 10.60 ± 0.45) in fresh, sprouted and toasted seeds respectively. Arginine is an essential amino acid that is converted to Nitric oxide which is a powerful neurotransmitter that helps to relax blood vessels and improves circulation. The amino acid profile obtained in this study agrees with the one obtained by [28]

The result of the mineral composition as presented in table 5 shows that Iron was the most abundant with concentration of 0.006 ± 0.001 , 0.006 ± 0.001 and 0.007 ± 0.001 mg/100g in all the three samples (fresh, sprouted and toasted respectively). There was no significant difference among the values. Iron is an essential element for production and it is vital for the proper functioning of hemoglobin, a transport protein responsible for the transport of oxygen in the body. Shortage of Iron leads to anemia. This results agrees with the result of [25]. Zinc is the second abundant minerals in all the samples analyzed. From the results, processing did not affect the mineral composition as there was no statistical difference ($p > 0.05$) among the values. It is needed for the body's defensive (immune system). Zinc also plays a role in cell division, growth and wound healing. Sodium and manganese were the next abundant in fresh, sprouted and toasted watermelon seeds. There was no significant difference among the values. Sodium regulates fluid balance in the body as well as maintain the functions of muscles and nerves while Manganese play roles in enzyme activation and helps in amino acids and protein digestion and utilization. Copper is the least abundant minerals in fresh, sprouted and toasted samples (0.001 ± 0.002 , 0.001 ± 0.00 and 0.001 ± 0.00 mg/100g respectively). There was no significant difference in the values obtained for the three samples. Copper and Iron enables the body to form red blood cells. Manganese also maintains healthy bones, blood vessels and nerves. It prevents cardiovascular disease and osteoporosis. Therefore, from the results obtained, sprouted *Citrullus lanatus* seeds may not be regarded as a good source of minerals. The high value of some of these minerals may satisfy the nutritional needs of the consumer.

CONCLUSION

The results obtained from this study has shown that *Citrullus lanatus* seeds (fresh, sprouted and toasted) may be a good source of protein, minerals and phytochemicals. Toasting increases the shelf life (decreasing the moisture content), decreases the antinutrients considerably and does not seem to affect the concentration of minerals, and amino acids. However, all the three samples contains sodium which is implicated in cardiovascular and renal disorder so it may not be good for people suffering from hypertension.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. **Mathew TJ, Ndamitso MM, Ofori AA, Shaba, EY, Innobeme A, Adamu A** Proximate and mineral compositions of seeds of some conventional and non-conventional fruits in Niger State, Niger, *Academic Research International Journal*, 2014: **5**: 2223 - 9944.
2. **AL-Khalifa AS** Physicochemical characteristics, fatty acid composition, and lipoxygenase activity of crude pumpkin and melon seed oils. *Journal of Agricultural and Food Chemistry*, 1996: **44**: 946-966.
3. **De Conto LC, Gragnani MA, Maus D, Ambiel HCI, Chiu MC, Grimaldi R, Goncalves LAG** Characterization of crude watermelon seed oil by two different extraction methods. *Journal of American Oil Chemists' Society*, 2011: **88**: 1709 - 1714.
4. **Jarret RL, Levy IJ** Oil and fatty acid content in seed of *Citrullus lanatus* Schard. *Journal of Agricultural and Food Chemistry*, 2012:**60**: 5199-5204.
5. **Baboli ZM, Kordi AAS** Characterization and consumption of watermelon seed oil and solvent extraction parameters effects. *Journal of the American Oil Chemists' Society* 2010:**87**: 667 - 671.
6. **Wickens GE, Gooding JR, Field DV** Plants for Arid Lands. George Allen and Unwin, London.1984: 223-321.
7. **Yadav S, Tomar AK, Jithesh O, Khan MA, Yadav RN, Srinivasan A, Singh TP** Purification and partial characterisation of molecular weight vicin-like glycoprotein from seeds of *Citrullus lanatus*. *Protein Journal*, 2011:**30**: 575 - 580.
8. **Acar R, Ozean MM, Kambur G, Durssun N** Some phytochemical properties of edible and forage watermelon seeds. *Iranian Journal of Chemistry and Chemical Engineering*, 2012:**13**: 41-47.
9. **Vaughan JG and Geissler CA** The new Oxford Book of food plants. Oxford University, Press, UK, 2009.
10. **Maggs-Kölling GL, Madsen S and Christiansen JL** A phenetic analysis of morphological variation in *Citrullus lanatus* in Namibia Genetic Resources and Crop Evolution, 2000: **47**: 385-393.
11. **Gunner N and Wehner T** The genes watermelon. Horticultural Sciences 2004:**39**: 1175-1182.
12. **Wani AA, Sogi DS, Wani TA, Shivhare US** Characterization and functional properties of watermelon (*Citrullus lanatus*) seed proteins. *Journal of the Science of Food and Agriculture*, 2011:**91**: 113 - 121.
13. **Loukou AL, Gnakri D, Dje Y, Kippre AV, Malice M, Baudoin JP, Zoro Bi IA** Macronutrient Composition of Three Cucurbit Species Cultivated for Seed Consumption in Cote D'Ivoire. *African Journal of Biotechnology*, 2007:**6** (5): 529-533.
14. **Sofowora A** Medicinal Plants and Traditional Medicine in Africa 2nd Edition. Spectrum Books Ltd, Ibadan, Nigeria, 1993: 289-290.

15. **Trease GE and Evans WC** A text book of pharmacognosy, 15th edition. Academic press, London 2002.
16. **Harbone JB** Phytochemical Methods. Chapman and Hall, New York. 1973.
17. **Ayoade G, Amoo I, Jabar J, Ojo A. and Maduawuchi C** Proximate, Minerals and Amino Acid Profile of (*Canarium Schweinfurthii*) Seed Pulp. *International Journal of Science and Technology*, 2017:6 (1): 657-661
18. **Bonire J, Jalil N and Lori J** Sodium and potassium content of two cultivars of white yam (*Dioscorea rotundata*) and their source soils. *Journal Science Food Agriculture*. 1990:53(2). 271-274.
19. **Lucas, GM. and Markaka P** Phytic acid and other phosphorus compound of bean(*Phaseolusvugaris*) *Journal of Agricultural Education and Chemistry*, 1975:23:13-15.
20. **AOAC (Association of Official Analytical Chemists)**. Official Method of Analysis of the AOAC (W. Horwitz Editor) Thirteenth Edition. Washington D.C, AOAC. 1980
21. **Abeza RH, Blake JT and Fisher EJ** Oxalate determination: Analytical Problems encountered with Certain Plant Species. *Journal of the Association of Official Agricultural Chemists*, 1968:51: 963-965.
22. **Food and Agricultural Organization/World Health Organization**. Protein quality evaluation. Report of Joint FAO/WHO Expert Consultant, FAO Food Nutrition paper; 51, FAO/WHO, Rome, 1991:19 – 21.
23. **Elinge CM, Muhammad A, Atiku FA, Itodo AU, Peni IJ, Sanni OM and Mbongo AN** Proximate, Mineral and Anti-nutritional Composition of Pumpkin (*Cucurbita pepo* L.) Seeds Extracts. *International Journal of Plant Research*. 2012:2(5):146- 150.
24. **Jacob AG, Etong DI, Tijjani A** Proximate, Mineral and Anti-nutritional composition of melon(*Citrullus lanatus*) seeds. *British Journal of Research*, 2015:2 [5] 142-157.
25. **Penuel BL, Khan ME, Maitera NO** Properties of proximate composition and elemental analysis of *citrullus vulgaris*(guna) seed. (*Bulletin of Environment, Pharmacology and loge Science India*). 2013:2(2):39-46.
26. **Abiodun OA and Adeleke RO** Comparative studies of nutritional composition of four melon seeds varieties. *Pakistan Jour. Of Nutrition*. 2010: 9(9): 905-908.
28. **Egbonu AC** Comparative investigation of the proximate and functional properties of watermelon (*Citrullus lanatus*) rind and seed. *Res. J. Environ. Toxicol*. 2015.

UNDER PEER REVIEW