

Original Research Article

The Environmental Friendliness Level Of Boat *Bagan* Fishing Gear In Belitung Regency

ABSTRACT

This research aimed to analyze the environmental friendliness level of boat *bagan* towards the amount of fish caught by the fishermen in Belitung Regency Waters and determined fish composition based on the type and the length of fish. The research was conducted in August 2019. The primary data collection technique used purposive sampling technique while the secondary data was obtained from Tanjungpandan Archipelago Fisheries Port (*Pelabuhan Perikanan Nusantara Tanjungpandan/PPN*). The research analyzed types of fish composition, proportion of main and bycatches, proportion of biologically worth caught fish, proportion of utilized catches, and eco-friendly analysis. Based on the results of research on the environmental friendliness level of boat *bagan* fishing gear in Belitung Regency Waters, it can be concluded that the boat *bagan* catch during the study obtained 10 species including goldstripe sardinella (*Sardinella fimbriata*), spotted sardinella (*Sardinella sirm*), anchovy (*Stoleporus devisi*), squid (*Loligo sp*), short mackerel (*Rastrelliger brachysoma*), common ponyfish (*Leiognathus equulus*), largehead hairtail (*Trichiurus sp*), and yellowstrip scad (*Selaroides leptolepis*), large pelagic fish such as barracuda (*Sphyaena sp*), and demersal fish is also caught such as pufferfish (*Tetraodo sp*). Boat *bagan* fishing gear is classified as less environmentally friendly because the fish caught that are not mature is 48%, the fish that are not the main target fish (bycatch) is 41%, and it catch more than 3 target fish species. However, the utilization of catch is quite good at 99, 12%.

Keywords: Belitung Regency Waters, Boat Bagan, Bycatch, Environmental Friendliness, Main Catch

1. INTRODUCTION

Belitung Regency is one of the area that has fishery potentials and is located in Fisheries Management Area (*Wilayah Pengelolaan Perikanan / WPP*) 711. There are many kinds of important economic marine fish in Belitung Regency. Belitung Regency has fishing activity center namely Tanjungpandan Archipelago Fisheries Port (*Pelabuhan Perikanan Nusantara Tanjungpandan/PPN*). The port location is very strategic because it is close to fishing ground area and marketing center, both domestic and abroad. Capture fisheries is still the mainstay of the marine fishery sector in Belitung regency after the end of tin mining. It consistently becomes the economic driver for Belitung. There are many important economic marine fish in Belitung Regency, such as spotted sardinella, goldstripe sardinella, anchovy, squid, and big fish like skipjack and mackerel. The abundant resources are due to its open location that directly faces South China Sea.

Lift net is a fishing gear that is operated by putting it down and lifting it up vertically. This net is usually made of nylon net that resembles a mosquito net. The net is then tied to square wooden frame. The mesh size is usually very small approximately 0,5 cm. In the usage, lift net usually uses lamps or baits to invite fishes. It is usually operated on a boat, raft, or permanent building. From the shape and usage, lift net can include boat *bagan*, embedded *bagan*, and lift net scoop [1].

Bagan is a fish fishing gear that uses artificial light to attract fish. Belitung Regency fishermen use boat *bagan* as a fishing gear for fishing. In the process of fishing using *bagan* boat, the artificial light used is aimed to collect fish that has positive phototaxis characteristic. According to Yuda (2012), fish that has positive phototaxis characteristic will gather in the artificial light area so that it

is easier for the fishermen to catch them [2].

According to Minister of Maritime Affairs and Fisheries Regulations No.71/2016 concerning Fishing Lane and the Placement of Fishing Equipment in the Fisheries Management Area of the Republic of Indonesia, boat *bagan* uses mesh size ≥ 1 mm, $P < 20$ m, and $L < 20$ m, uses ABPI in the form of lamps with total power $< 2,000$ watt, uses one or two motor boats with total size > 5 until 10 GT.

The catch are divided into two, namely Main Target Catch (*Hasil Tangkapan Sasaran Utama / HTSU*) that means the species are the target of the fishing operation, and the other one is Bycatch (*Hasil Tangkapan Sampingan / HTS*) that means the fish is just an addition and not the target of the fishing operation.

Standardization for fishing gear on fish target is needed because the caught fish size varies. To improve environmental friendliness and *bagan* fishing gear standardization, it is important to note the initial fork length of the length at first maturity (LM).

2. MATHERIALS AND METHODS

The method used in this research is survey method. Survey method is a direct data collection method in the field, and it conducts data collection by concentrating the research on a case intensively so that overall picture can be obtained as a result of data collection and data analysis in a certain period of time and limited to certain areas [3]. The primary data uses purposive sampling, which is a technique to determine research sample with certain considerations so that the obtained data can be more representative [4].

This research was conducted in August 2019. The research location is Belitung Regency Waters. The materials and

equipment used include boat *bagan* fishing gear, rollmeter (1 mm scale), digital scale (1 gr scale), smartphone camera, stationary, and fish catch. In this research, data collection is conducted in some stages, namely:

1. Directly followed fishing operation using boat *bagan* in August 2019 as much as 10 times using boat with a size of 10 GT.
2. The primary data that is collected directly during the research namely catch composition, fork length, individual weight of the catch, the utilization of the catch, and the proportion of the Main Target Catch and Bycatch.
3. Interview with 21 fishermen of *bagan* boat by using prepared questionnaire

2.1 Catch Type Composition

Before being analyzed, the catch is identified to determine the general and the latin name. The identification is in accordance with Fishbase. After being analyzed, the data is grouped according to the species and then the weight and the number are counted. The fish type is then tabulated to see the catch composition.

2.2 Main Target Catch and Bycatch Proportion

Each data of the number and weight of Main Target Catch and Bycatch from the fishing operation are counted in the form of percentage. Then, it is compared to see which of the catch has bigger proportion. The data of fork length of the catch is processed by counting the frequency distribution and is presented in the form of histogram.

2.3 The Proportion of Fish Eligible Catching Biologically

The fish size that is eligible for catching is determined by the length at first maturity. The length measurement is done with fork length (FL) measurement, which is the measurement from mouth to tail fork. The obtained data is presented in descriptive analysis using graphic and the average length value of the most caught fish to find out gonad fish maturity level. After analyzing all the obtained data, if the length of fish that is caught most (L_c) less than the length of fish that has mature gonad first (L_m), it can be concluded that the fishing gear is not selective and vice versa, but this remains based on each species.

2.4 The Proportion of Utilized Catch

All the data of utilized catch is compared with the unutilized catch (discarded) in form of proportion. The measurement of the catch is presented in kg. The observation result of the catch handling process is analyzed to describe the catch that are utilized and unutilized or discarded.

2.5 The Analysis of Environmental Friendliness Level

A fishing gear can be said as an environmentally friendly fishing gear if the main target catch is bigger than the bycatch. The friendliness factor that is used as research to see friendliness level is Mallawa method (2006) of the calculated data and scores on the assessment of the level of fishing gear friendliness (Table 1).

Table 1. The Assessment of Environmental Friendliness Level

Observation	Assessment	Criteria	Score
Main Target Catch (%)	81-100	Very environmentally friendly	4
	61-80	Environmentally friendly	3
	41-60	Less environmentally friendly	2
	1-40	Not environmentally friendly	1
Length at First Maturity (%)	81-100	Very environmentally friendly	4
	61-80	Environmentally friendly	3
	41-60	Less environmentally friendly	2
	1-40	Not environmentally friendly	1
Utilized Catch (%)	81-100	Very environmentally friendly	4
	61-80	Environmentally friendly	3
	41-60	Less environmentally friendly	2
	1-40	Not environmentally friendly	1

Conclusion :

- Total score between 3 and 5 : not environmentally friendly
- Total score between 6 and 8 : less environmentally friendly
- Total score between 9 and 11 : environmentally friendly
- Total score 12 : very environmentally friendly

3. RESULT AND DISCUSSION

Based on the research activities, the following results are obtained:

3.1 Boat *Bagan* Catch

3.1.1 The Fish Type Diversity of Boat *Bagan* Catch

From the identification of the fish type diversity of boat *bagan* catch during the fishing operation of this research, from three units of boats with a total of 10 trips in Belitung Regency Waters, ten marine biota are obtained that consist of eight type of pelagic fish, one demersal fish, and one squid (Table 2).

Table 2. The Fish Type Diversity of the Catch

Name of Fish	Latin Name (<i>species</i>)
Barracuda	<i>Sphyraena</i> sp
Pufferfish	<i>Tetraodo</i> sp
Squid	<i>Loligo</i> sp
Short mackerel	<i>Rastrelliger brachysoma</i>
Common ponyfish	<i>Leiognathus equulus</i>
Largehead hairtail	<i>Trichiurus</i> sp
Spotted sardinella	<i>Sardinella sirm</i>
Yellowstrip scad	<i>Selaroides leptolepis</i>
Goldstripe sardinella	<i>Sardinella fimbriata</i>
Anchovy	<i>Stolephorus devisi</i>

The diversity shows that the *bagan* boat is a multispecies fishing gear; it catch

more than one type of fish, which mean it has low selectivity.

Based on the results of interviews with *bagan* fishermen and the PPN Tanjungpandan Syahbandar Chief, the main target of boat *bagan* fishing gear in the Belitung Regency Waters is goldstripe sardinella (*Sardinella fimbriata*), spotted sardinella (*Sardinella sirm*), and anchovy (*Stolephorus devisi*). These three commodities have high economic value that can increase people's income and are usually caught in large quantities.

3.1.2 The Proportion of Main Target Catch and Bycatch

The total catch of ten trips is 504,4 kg. The number of main target catch is

71,530 fish (89%) with 298,3 kg of weight (59%) (Table 3). The main catch is anchovy (*Stolephorus devisi*) that is 65,127 (81.1%), with 121.4 kg (24.1%) of weight, followed by spotted sardinella (*Sardinella sirm*) that is 3,200 (4%), with 100,9 kg (20%) of weight, then goldstripe sardinella (*Sardinella fimbriata*) that is 3,203 (4%), with 75.9 kg of weight (15,1%) (Table 4).

The number of Bycatch is 8,809 fish (11%) with 206.14 kg (41%) of weight from the total catch (Table 3). The main catch is squid (*Loligo* sp) that is 3,535 (4,4%), with 141.4 kg (28%) of weight (Table 4).

Table 3. The Weight and Number of the Catch

Category	Catch			
	Weight		Amount of fish	
	(kg)	%	(fish)	%
Main Catch	298,3	59%	71530	89%
Bycatch	206,1	41%	8809	11%
Total	504,4	100%	80339	100%

Table 4. The Proportion of the Catch

Name of Fish	Amount of Fish	%	Weight (kg)	%
Barracuda	91	0,1	11,5	2,3
Pufferfish	1	0,0	0,2	0,0
Squid	3535	4,4	141,4	28,0
Short mackerel	3737	4,7	32,7	6,5
Common ponyfish	629	0,8	5,5	1,1
Largehead hairtail	129	0,2	5,4	1,1
Spotted sardinella	3200	4,0	100,9	20,0
Yellowstrip scad	687	0,9	9,5	1,9
Goldstripe sardinella	3203	4,0	75,9	15,1
Anchovy	65127	81,1	121,4	24,1
Total	80339	100	504,4	100

The *bagan* catch is very diverse ranging from fish that has positive phototaxis characteristic to fish that migrate. The composition of this type of catch is

obtained during the eastern monsoon (From July – August). Even though the *bagan* is operated in different season, goldstripe sardinella, spotted sardinella, and anchovy fish remain to become the

main target of the fishing operation of boat *bagan* in PPN Tanjungpandan Belitung Regency.

During this research, the *bagan* boat uses *rumpon* and lamp as to attract fish. According to [5], it is stated that the events of fish attracted to light can be divided into two, namely:

- a. Direct event, the fish is attracted to light and then gather. This is related to phototaxis.
- b. Indirect event, as there is light, plankton and other tiny fish gather, then the target fish gather to feed. Some fish that include in this category are mackerel tuna and barracuda.

The net used on the boat *bagan* is made of waring without a knot, in the form of a rectangular bag, measuring 16 x 6 x 13.5 meters and a mesh size of 0.3 cm. Another factor that causes the variety of catch from a fishing gear is caused by a group of fish between target fish and non-target fish, so that migratory fish can be caught.

To improve the level of environmental friendliness of the fishing gear, the mesh size must be modified so that it can catch the main target fish properly. Prevention can be done by efforts to improve the selectivity of fishing gear

according to [6]. namely by modifying the size of the net mesh from a small size into a larger size so as not to catch fish that are not yet feasible to catch.

3.1.3 Frequency Distribution of Main Catch Fish Fork Length

Boat *bagan* is designed to catch small pelagic fish such as goldstripe sardinella, spotted sardinella and anchovies. The types of fish caught in boat *bagan* usually form a schooling, such as spotted sardinella, goldstripe sardinella, and anchovies. When this research was conducted, small-sized fishes were also caught, allegedly because the construction of the boat chart has a small mesh size, which ranges from 0.3 to 0.5 cm with nylon material so that it allows small fish to be entangled in the net. even the size that is still juvenile.

Goldstripe sardinella caught have different fork lengths ranging from 100 to 155 mm. The sizes range from 121 to 127 mm are the most caught at the time of this research, which is as many as 39 fish, and the smallest amount obtained from the catch is size 149 to 155 mm where there are only 7 fish (Figure 1).




Figure 1. The Distribution of Goldstripe Sardinella Fork Length

Spotted sardinella caught during the research has different lengths of fork ranging from 104 to 166 mm. The sizes range from 125 to 131 mm are the most caught at the time of this research,

which is as many as 58 fish, and the smallest amount obtained from the catch ranges from 104 to 110 mm where there are only two fish (figure 2).




Figure 2. The Distribution of Spotted Sardinella Fork Length

Anchovy caught during research has different fork lengths ranging from 53 to 76 mm. The sizes range from 56 to 58 mm are the most caught at the time of this research, which is 29 fish, and the

smallest amount obtained from the catch ranges from 74 to 76 mm where there are only 10 fish (Figure 3).




Figure 3. The Distribution of Anchovy Fork Length

The size of the fish caught will affect the applicable selling price. Besides that, more importantly, the size of the fish caught will affect the sustainability of the fish resources themselves in particular and the sustainability of the ecosystem in general. The nature of the pelagic fish in groups causes the captured pelagic fish to have several size groups [7]. If from a fishing operation conducted small-sized fish is caught, it can be estimated that other fish caught is small-sized fish as well.

3.1.4 The Proportion of Fish Eligible for Main Catch

The length of the caught fish can be used to determine whether or not the fish is eligible for catching by knowing the length limit of the fish when the gonad (Length at first maturity) is first mature. Catching above the size of the fish when the gonad is first mature can provide an opportunity for the target fish to be able to reproduce and spawn before being caught so that the recruitment phase of small fish into the

adult fish phase can run. Therefore, the size criteria for catching is the most important criteria for determining the environmental friendliness of a fishing operation.

Ningrum et al. (2015) stated that by comparing $L_{c50\%}$ size with LM values, it can be assumed that the first fish caught has spawned or not [8]. If the size of the fish is bigger than LM, the fish have spawned. The length of the gonad's first mature length using an embedded *bagan* in the Sungsang Waters, South Sumatra for anchovies is 62 mm and the estimated age to reach LM is around 3.7 months [9]. Data on the distribution of the length of the anchovy fork is sampled as many as 150 fish. There are 89 fish that are eligible for catching and 61 are not eligible for catching. The proportion of anchovy that is eligible for catching is 59% (Figure 4). Most of the anchovy caught during the study is fish eligible for catching.




Figure 4. The Proportion of Anchovy that is Eligible for Catching

Nugraha (2015) stated that the LM values of male and female spotted sardinella in the Sunda Strait were 137.30 mm and 145.82 mm respectively. In general, spotted sardinella experience gonad maturity, which first occurs in the range of 65-75% of the maximum length (Setyohadi 2010). Data on the distribution of the length of the spotted sardinella fork is

sampled as many as 150 fish. There are 14 fish that are eligible for catching and 136 fish are not eligible for catching. The proportion of spotted sardinella catch that are eligible for catching is 9% (Figure 5). Spotted sardinella caught during the study are mostly fish that are not eligible for catching.




Figure 5. The Proportion of Spotted Sardinella that is Eligible for Catching

Bintoro (2019) stated that the length of the first mature gonad goldstripe

sardinella in the Bali Strait for males and females was 11.95 cm and 10.79 cm

respectively [10]. In the study of Aryuningka (2016) in the Sunda Strait, the first mature female goldstripe sardinella gonad is 157 mm. Data on the distribution of the length of the fork of the goldstripe sardinella is sampled as many as 150 fish. There are 131 fish that are eligible for catching and 19 fish that are not eligible for catching. The

proportion of goldstripe sardinella eligible for catching is 87% (Figure 6). Goldstripe sardinella caught during the study are mostly fish eligible for catching.




Figure 6. The Proportion of Goldstripe Sardinella Eligible for Catching

According to [11], the total number of fish eligible for catching has a proportion of 52% (Figure 7), so it can be said that boat *bagan* is less environmentally

friendly. This is because boat *bagan* has a mesh size that is too small, which is 0.3 cm, so that many small fish that are not mature are caught.




Figure 7. The Total Proportion of Catching Eligibility

3.2 The Composition of Catch Utilization

Utilization of catch is a factor that can improve the level of environmental friendliness of a fishing gear. Some bycatch are utilized and some are not utilized by fishermen. The catch is usually used because of high economic value, safe for consumption and demanded by the community. Catch that are not utilized are usually not of

economic value, dangerous to eat and less demanded by the community.

Catch caught are usually used by fishermen for sale and self-consumption. The catch sold is 388.1 kg (76.35%). The catch consumed is 114.9 kg (22.77%) (Table 5). Catch disposed are 4.4 kg (0.88%). The catch that are disposed of at the time of this research are due to the little weight that is lack of economical value and the danger of the fish like pufferfish that is dangerous if consumed.

Table 5. The Utilization of Main Target Catch and Bycatch

Utilization	Main Catch		Bycatch		Total	
	Weight (kg)	%	Weight (kg)	%	Weight (kg)	%
1	Utilized					
On Sale	223,2	74,82%	161,9	78,55%	388,1	76,35%
Self Consumption	75,1	25,18%	39,8	19,30%	114,9	22,77%
2	Not Utilized					
Not Sold	0	0%	0	0%	0	0%
Discarded	0	0%	4,4	2,15%	4,4	0,88%
Total	298,297	100%	206,1	100%	504,4	100%

Overall, the catch obtained, both the main catch and bycatch, are mostly used by fishermen. Main catch utilized are 298.3 kg (100%), and bycatch used by fishermen are 201.7 kg (97.85%) and the bycatch that is not used is 4.4 kg (2.15%) from the total catch caught including pufferfish (*Tetraodo sp*), common ponyfish (*Leiognathus sp*), largehead hairtail (*Trichiurus sp*), and yellowstrip scad (*Selaroides leptolepis*).

The analysis of environmental friendliness level of boat *bagan* is done to determine the level of friendliness to the environment of these fishing gear in order to implement sustainable fisheries in accordance with the provisions of responsible fishing practices.

The analysis environmental friendliness level can be seen from several parameters, namely the composition of the catch, the proportion of the main catch and bycatch, the proportion of fish eligible for catching, and the scoring value to determine the level of friendliness of the boat *bagan*. A

3.3 The Analysis of Environmental Friendliness Level

summary of the results of an assessment of the level of

environmental friendliness of the boat *bagan* unit is presented in Table 6.

Table 6. Assessment Results of the Environmental Friendliness Level

Observation	Assessment	Criteria	Score
Main Catch	59%	Less environmentally friendly	2
Length at First Maturity	52%	Less environmentally friendly	2
Utilization of Catches	99,12%	Very environmentally friendly	4
Total Score			8

Based on the assessment of the level of environmental friendliness by using a score, the total score obtained from the three factors used is 8. 8 out of 6, 7 and 8, based on the scores obtained from the three factors used in determining the level of environmental friendliness of the boat *bagan*, the boat *bagan* in PPN Tanjungpandan is classified as less environmentally friendly.

The main catch of the boat *bagan* are anchovy, spotted sardinella and goldstripe sardinella as much as 298.3 kg (59%) of the total catch. According to [11], based on this proportion, the boat *bagan* unit is less environmentally friendly when viewed in terms of catch weight. The amount of bycatch obtained is too much. More than 3 species causes the selectivity value of the boat *bagan* is not environmentally friendly.

The level of environmental friendliness based on the length of the fork of the main catch can be seen from the long distribution of existing classes. Catching fish above the size of the first time the gonads mature can provide opportunities for the target fish to be able to reproduce and spawn before being caught, so the process of breeding and the phase of small fish into adult fish can run continuously. Thus, the determination of the eligibility of the fish to be caught is closely related to determining the level of environmental friendliness of the fishing

operation. This is based on [6], that one of the selective fishing processes is that it does not endanger the sustainability of target fish resources. The number of fish eligible for catching during the research is 234 out of 450 fish of the target fish sampled, With a percentage of 52%, based on this information it can be seen that the target fish caught in the boat *bagan* mostly include fish that are eligible for catching, so the number of fish is eligible for catching is 52%, according to [11], the boat *bagan* is less environmentally friendly.

Based on the results of the study of environmental friendliness level of several criteria, it can be said that all criteria show less environmentally friendly, that is, if viewed from the composition of the catch and catch size, this is due to the size of the mesh used in fishing operations that is too small so that it can catch many non-target fish and can catch fish that are not yet eligible to catch. Boat *bagan* units operating in PPN Tanjungpandan have fulfilled Minister of Maritime Affairs and Fisheries Regulations No.71/2016 concerning Fishing Lane and the Placement of Fishing Equipment in the Fisheries Management Area of the Republic of Indonesia, so it is necessary to have law enforcement or legal rules in accordance with the implementation in the field so that the fishing gear is more selective so that it only catch the

target fish, or renovates the boat *bagan* framework to increase its selectivity.

4. CONCLUSIONS

Based on the results of research on the environmental friendliness level of boat *bagan* fishing gear in Belitung Regency Waters, it can be concluded that the boat *bagan* catch during the study obtained 10 species including goldstripe sardinella (*Sardinella fimbriata*), spotted sardinella (*Sardinella sirm*), anchovy (*Stoleporus devisi*), squid (*Loligo sp*), short mackerel (*Rastrelliger brachysoma*), common ponyfish (*Leiognathus equulus*), largehead hairtail (*Trichiurus sp*), and yellowstrip scad (*Selaroides leptolepis*), large pelagic fish such as barracuda (*Sphyraena sp*), and demersal fish is also caught such as pufferfish (*Tetraodo sp*).

Boat *bagan* fishing gear is classified as less environmentally friendly because the fish caught that are not mature is 48%, the fish that are not the main target fish (bycatch) is 41%, and it catch more than 3 target fish species. However, the utilization of catch is quite good at 99, 12%.

5. REFERENCES

- 1 Subani W dan HR Barus. 1989. Marine Fish and Shrimp Fishing Equipment in Indonesia. *Journal of Marine Fisheries Research*. No. 50. Jakarta: Indonesian Fisheries Research and Development Agency. Agriculture department. Page. 1-248 pages, ISBN/ISSN 0216-7727
- 2 Yuda L.K., D. Iriana., A. M. A. Khan. 2012. The Environmental Friendliness Level of *Bagan* Fishing Gear in Pelabuhanratu Waters, Sukabumi Regency. *Marine and Fisheries Journal*. Vol. 3, No. 3, ISSN 2088-3137
- 3 Sugiyono. 2009. *Quantitative, qualitative and R&D Research Methods*. Bandung: Alfabeta.
- 4 Sugiyono. 2014. *Quantitative, qualitative, and R&D Approach Educational Research Methods*. Bandung: Alfabeta
- 5 Ayodhya, A. U. 1981. *Fishing Methods*. Bogor. Dewi Sri Foundation. 91 pages.
- 6 Monintja D. R. 2000. *Utilization of Coastal Resources in the Field of Capture Fisheries*. Training Proceeding on Integrated Coastal Management. Center for Coastal and Ocean Resource Study. Bogor Agricultural Institute. Bogor
- 7 Takwim dan H. Setiawan. 2009. The Main Catch and Bycatch Purse Seine Unit in Pekalongan Regency. *Department of Capture Fishery Resources 1-14*
- 8 Ningrum, V. P., Ghofar, A., Ain, C. 2015. Some Biological Aspects of Rajungan Fisheries (*Portunus pelagicus*) in Betahwalang and surrounding waters. *Journal of Fisheries Science*. 11(1): 62-71
- 9 Fauziyah, Saleh K, Hadi, Supriyadi F. 2012. Response of Difference in Light Intensity of Petromac Lamps towards Embedded *Bagan* Catch in Sungsang Waters, South Sumatra. *Maspari Journal*. 4(2): 215-224
- 10 Bintoro. G., D. Setyohadi, T.D. Lelono., F. Maharani., 2019. Biology and Population Dynamics Analysis of Fringescale Sardine (*Sardinella fimbriata*) in Bali Strait Waters, Indonesia. *IOP Conf. Series: Earth and Environmental Science 391(2019) 012024*

- 11 Mallawa A. 2006. *Study of Estimating the Potential of Fisheries and Marine Resources of Selayar Regency*. Makassar: Hasanuddin University, Faculty of Marine and Fisheries Sciences

UNDER PEER REVIEW