

1 **Epidemiological study of intestinal parasites in school children in Vandeikya**  
2 **LGA, Benue State, Nigeria**

3  
4  
5 **ABSTRACT**

6 A study was conducted to determine the epidemiology of intestinal parasites in  
7 school children in Vandeikya LGA, Benue State, Nigeria was conducted. Two  
8 hundred and ninety three (293) stool samples from school children were examined  
9 in selected schools across the Local Government Area. 32 of 293 children were  
10 infected with parasites which include: *Ascaris lumbricoides*, *Entamoeba*  
11 *histolytica*, Hookworm, *Strongyloides stercularis* and *Taenia spp*. Hookworm had  
12 the highest prevalence rate of 9(3.1%), *Entamoeba histolytica* 7(2.3%), *Taenia spp*  
13 had 3(1.0%), *Strongyloide stercularis* had 2(0.7%) while *Ascaris lumbricoides*  
14 recorded the least prevalence rate of 1(0.3%). There were also of parasitic infection  
15 in the pupil. *Entamoeba histolytica*, and *Ascaris lumbricoides* recorded the highest  
16 prevalence rate of 5(1.7%). *Entamoeba histolytica*, Hook worm and *Strongyloides*  
17 *stercularis* recorded 2(0.7%) while *Entamoeba histolytica*, *Ascaris lumbricoides*  
18 and Hook worm recorded the least prevalence 2(0.7%). There is significant  
19 difference ( $P < 0.05$ ) between the age groups and prevalence of intestinal parasites  
20  $\chi^2$  calculated (10.117) is greater than  $\chi^2$  tabulated (4.891) at  $df = 2$ , while there was  
21 no significant difference ( $P > 0.05$ ) in the infection rate based on sex, because the  
22 value of  $\chi^2$  calculated (3.245) is less than  $\chi^2$  tabulated value (5.991) at  $df = 2$ .  
23 Intestinal parasites are prevalent in Vandeikya LGA, Benue State. Risk factors like  
24 open defecation, use of stream and well water should be minimized in order to  
25 prevent infection.

26 **Keywords: Prevalence, intestinal parasites, mixed infection**

27

## 28 **Background**

29 Intestinal parasites are helminthes (worms) and protozoa that reside in the  
30 intestines of their hosts; disease burden is higher in children and women (WHO,  
31 2017). At least 880 million children in need of treatment for intestinal worms  
32 (WHO, 2017). According to World Health Organization estimates, 500 million  
33 people in the world are infected with *Entamoeba histolytica*, this protozoon causes  
34 symptomatic illness in about 50 million people and is responsible for 100,000  
35 mortalities (Lozano *et al.*, 2012). Intestinal parasites may cause morbidities like:  
36 weakness, inflammation of the intestines, abdominal pains, nausea and dysentery  
37 in those infected (Ashtiani *et al.*, 2011). In addition, infection in children may  
38 result in reduced ability to learn, reduced food absorption leading to malnutrition,  
39 anaemia, stunted growth, and may lead to death (Amuta *et al.*, 2013). Intestinal  
40 parasites are endemic in the tropics where favorable climatic, environmental and  
41 sociocultural factors permit transmission (Alli *et al.*, 2011a). The infection rate for  
42 these parasites has primarily been attributed to poverty, unhygienic environmental  
43 conditions and over-dispersion of parasites (Amuta *et al.*, 2013). Significantly  
44 higher prevalence of intestinal parasites was reported in children from homes  
45 where livestock are reared and in children of farmers (Butera *et al.*, 2019).  
46 According to a study conducted in Ethiopia, hygiene factors like the inconsistent  
47 use of shoes is strongly associated with prevalence of hookworm infection, and  
48 long untidy finger nails and drinking of well water are important risk factor for *E.*  
49 *histolytica* (Hailegebriel, 2017). Wei Liao *et al.* (2017) also reported drinking of  
50 rain water rainwater as a possible risk factor for intestinal parasites. In contrast,  
51 Erismman *et al.* (2016) found no association between household drinking water  
52 source in school aged children in Burkina Faso. Proper understanding of  
53 prevalence, distribution and associated risk factors for intestinal parasites could

54 help in development of strategy aimed at transmission interruption and control of  
55 the diseases they cause. This study was therefore aimed at: 1) investigating the  
56 prevalence and distribution of intestinal parasites in school aged children in  
57 Vandeikya LGA and (2) determining the association between prevalence of  
58 parasite species in children and three risk factors: drinking water source,  
59 occupation of parents and type of toilet used.

## 60 **MATERIALS AND METHODS**

### 61 **Study area**

62 The research was carried out in select public and privates schools across  
63 Vandeikya Local Government Area of Benue State.

### 64 **Ethical clearance**

65 Ethical clearance was obtained from research ethics board. Informed consent and  
66 was obtained from parents, school authorities and informed accent from research  
67 subjects.

### 68 **Samples collection**

69 School children that participated in the study were aged between 3 – 15 years.  
70 With the assistance and cooperation a their teachers, school children were given  
71 20ml universal bottles the previous day and asked to return it the next morning  
72 with their stool sample (morning stool). Questionnaires were used to obtain data on  
73 age, sex, drinking water source, type of toilet system used at home and occupation  
74 of parents. The stool samples were transported to laboratory at General Hospital  
75 Gboko where they were analyzed.

### 76 **Method of identification of intestinal parasites**

77 The methods that was used to identify the presence of intestinal parasites was the  
78 direct microscopy of wet preparation, and the concentration method stool sample  
79 was emulsified in saturated salt solution in the test tube, the supernatant was pour

80 out and the concentrate was centrifuge for 5 minutes. A drop or two of saline  
81 solution was drop on the microscopic slide properly covered with cover slip and  
82 viewed under a microscope with  $\times 10$  and  $\times 40$  objective lens.

### 83 **Statistical analysis**

84 Chi-square test was used to determine the homogeneity of the disease in the  
85 different schools.

### 86 **RESULTS**

87 Table 1 shows age related prevalences of a range of intestinal parasites present in  
88 stool samples of research subjects: The total prevalences of the intestinal parasites  
89 were as follows: *Ascaris lumbricoides* 1 (0.3%), *Entamoeba histolytica* 7 (2.3%),  
90 Hookworms 9 (3.1%), *Strongyloides stercularis* 2 (0.7%), *Taenia* spp. 3 (1.0%).  
91 Among children aged 0-4 years, Hookworm was the most prevalent intestinal  
92 parasite with prevalence of 2 (2.8%), and no intestinal parasite was observed  
93 among children aged 15-19 years. There was no significant difference in  
94 prevalence between the different age groups.

95 Table 2 shows sex related prevalences of intestinal parasites in stool samples of  
96 research subjects. In males, prevalences in *Ascaris lumbricoides* was 1 (0.6%),  
97 *Entamoeba histolytica* 4 (2.4%), Hookworms 1 (0.6%), *Strongyloides stercularis* 6  
98 (0.7%), and *Taenia* spp 0 (0.0%). Prevalences in females were as follows: *Ascaris*  
99 *lumbricoides* 0 (0.0%), *Entamoeba histolytica* 3(2.2%), Hookworms 5 (3.7%),  
100 *Strongyloides stercularis* 1 (0.7%), *Taenia* spp. 19 (14%). Females had higher total  
101 prevalences of intestinal parasites 19(14%) than males 13(8.3%).

102 Table 3 shows the prevalence of mixed intestinal parasite infections in school  
103 children with respect to age. *Entamoeba histolytica*, *Ascaris lumbricoides* and  
104 Hookworm had prevalence of 2 (0.7%); *Entamoeba histolytica* and *Ascaris*  
105 *lumbricoides* had the highest prevalence rate of 5(1.7%); and *Entamoeba*  
106 *histolytica*, Hook worm and *Strongyloides stercolaris* had prevalence of 2(0.7%).

107 Table 4 showed the prevalence of mixed intestinal parasites in school children with  
 108 respect to sex. Prevalence of mixed infections was higher among males 4(2.5%)  
 109 than among females 5(3.7%), giving a 9(3.1%) total prevalence of mixed  
 110 infections.

111 Figure 1. Represents the prevalence of intestinal parasites in relation to the type of  
 112 toilet used by the subjects. Those that used nearby bush were more infected 45% of  
 113 total infections with intestinal parasites, while those that used Pit and water system  
 114 toilets had 40% and 15% total infection respectively.

115 Figure 2 shows the relationship between source of drinking water and the  
 116 prevalence of intestinal parasites. School children whose obtain drinking water  
 117 from streams were most infected accounting for 60% of total infections, those who  
 118 obtain drinking water from wells had the second highest infection rates (30%) and  
 119 those who obtain drinking water from borehole were the least infected accounting  
 120 for (10%) of the total infection.

121 Figure 3: shows the relationship between parental occupation of school children  
 122 and infection with intestinal parasites. Children whose parents were farmers  
 123 accounted for (50%) of those infected, children whose parents were Traders  
 124 accounted for (30%) of those infected and children while parents were Civil-  
 125 servants accounted for 20% of those infected.

126 **Table 1:** Age related prevalence of intestinal parasites

Age	No. Examined	<i>A. lumbricoides</i> (%)	<i>E. histolytica</i> (%)	Hookworm (%)	<i>S. stercoralis</i> (%)	Taenia spp (%)
0 – 4	67	0(0.0)	1(1.49)	2(2.99)	1(1.49)	0(0.0)
5 – 9	146	1(0.68)	3(2.05)	4(2.74)	1(0.68)	3(2.05)
10 – 14	77	0(0.0)	3(3.90)	3(3.90)	0(0.0)	0(0.0)

15 – 19	3	0(0.0)	0(0.0)	0(0.0)	0(0.0)	0(0.0)
Total	293	1(0.34)	7(2.39)	9(3.07)	2(0.68)	3(1.02)

$\chi^2$  Cal = 10.117, df = 12, P>0.05

127

128 **Table 2:** Sex related prevalence of intestinal parasites

129

Sex	No. Examined	<i>A. lumbricoides</i> (%)	<i>E. histolytica</i> (%)	Hookworm (%)	<i>S. stercoralis</i> (%)	Taenia spp (%)	Total (%)
Male	157	1(0.64)	4(2.55)	4(2.55)	1(0.64)	0(0.0)	13(8.28)
Female	136	0(0.0)	3(2.21)	5(3.68)	1(0.74)	3(2.21)	19(13.97)
Total	293	1(0.34)	7(2.39)	9(3.07)	2(0.68)	3(1.02)	32(10.9)

130

$\chi^2$  Cal = 3.245, df = 4, P>0.05

131

132

133

134 **Table 3:** Prevalence of mixed infections with intestinal parasites in different age groups

Age	No. examined	No. infected (prevalence)		
		<i>E. histolytica</i>	<i>E. histolytica</i>	<i>E. histolytica</i>
		<i>A. lumbricoides</i>	<i>A. lumbricoides</i>	Hookworm
		Hookworm		<i>S. stercoralis</i>
		(%)	(%)	(%)
0 – 4	67	0(0.0)	1(1.49)	1(1.49)
5 – 9	146	1(0.68)	2(1.4)	1(0.68)
10 – 14	80	1(1.25)	2(2.50)	0(0.0)
Total	293	2(0.68)	5(1.71)	2(0.68)

135  $\chi^2$  Cal = 2,948, df = 4, P>0.05

136

137


138 **Table 4:** Prevalence of mixed infections with intestinal parasites in different sexes

139

Sex	No. examined	No. infected (prevalence)			
		<i>E. histolytica</i>	<i>E. histolytica</i>	<i>E. histolytica</i>	Total (%)
		<i>A. lumbricoides</i>	<i>A. lumbricoides</i>	Hookworm	
		Hookworm		<i>S. stercoralis</i>	
		(%)	(%)	(%)	
Male	157	1(0.64)	2(1.27)	1(0.64)	4(2.55)
Female	136	1(0.74)	3(2.21)	1(0.74)	5(3.68)
Total	293	2(0.68)	5(1.71)	2(0.68)	9(3.07)

140  $\chi^2$  Cal = 0.052, df = 2, P>0.05


141



142

143 **Figure 1:** Relationship between toilet facility of the pupils and parasitic  
 144 infection.


145



146

147 **Figure 2:** Relationship between source of drinking water of the pupils and parasitic  
 148 infection.





149

150 Figure 3: Relationship between parental occupation of the pupil and parasitic  
 151 infection.

## 152 **DISCUSSION**

153 The study revealed the presence of *A. lumbricoides*, *E. histolytica*, *Hookworms*, *S.*  
 154 *stercoralis*, and *Taenia* spp infections in school aged children. Hookworms and *E.*  
 155 *histolytica* had the highest prevalences. The temperature in the tropics is known to  
 156 favour the prevalence of a range of intestinal parasites and could be the reason for  
 157 their presence in the study area. Seasonal variations in abiotic factors like  
 158 temperature, humidity and rainfall may affect intensity of intestinal parasites (Saki  
 159 *et al.*, 2017), particularly soil transmitted helminthes (Praharaj *et al.*, 2017).  
 160 Hookworms and *E. histolytica* had significantly higher prevalences than the other  
 161 parasites. Although the trophozoite stage of *E. histolytica* which is used to  
 162 diagnose the parasite is fragile and cannot survive outside the host for longer than  
 163 24 hours, the infective stage of the parasite which is the cyst stage can withstand  
 164 desiccation, harsh temperature and can linger in the environment for months, this

165 could be the reason for its high prevalence; in addition, the cysts can be transmitted  
166 through water, food. Hookworms are transmitted primarily by walking barefoot on  
167 soil containing parasite larvae, and by ingestion of larvae in the case of  
168 *Ancylostoma duodenale*. Multiple modes of infection could have contributed to  
169 high prevalence of hookworms in this study. In a study by Kumma *et al.* (2019),  
170 *E. histolytica* also had the highest prevalence among intestinal parasites infecting  
171 food vendors in Wolaita Sodo, Ethiopia, while tapeworms and hookworms had  
172 equal prevalences. The author opined that untrimmed fingernails of researcher  
173 subjects could be contributing to high prevalence of intestinal parasites in the  
174 study. Tegegne *et al.* (2018) however reported that *E. histolytica* had second  
175 highest prevalence and *A. lumbricoides* had the highest prevalences among  
176 tuberculosis suspected patients in Gondar, Northwest Ethiopia. However among  
177 protozoan infections, *E. histolytica* had the highest prevalence. The researchers  
178 opined that difference in water supply, economic status and environmental  
179 sanitation could be contributing to high prevalence of parasitic diseases in the  
180 study area.

181 Intestinal parasite prevalences were highest in children aged 5-9 years, as the  
182 children grew older (15-19 years), zero (0%) prevalences were recorded for all  
183 intestinal parasites. Children aged 5-9 may be more prone to playing with soil than  
184 other age groups and are likely to move around bare-foot thus exposing themselves  
185 to parasite infections. In addition this age group may have lower immunity than  
186 their older counterparts which may be an additional factor responsible for observed  
187 higher prevalences with intestinal parasites. Valiathan *et al.* (2016) opined that  
188 aging has an effect of immunity, and this could be a reason for variations in  
189 prevalences among different age groups with similar levels of exposure to  
190 infection. Gebretsadik *et al.* 2018 reported prevalences of intestinal parasite as high  
191 15.5%, in children under 5 years the author noted that poor hand washing habits of

192 this age group could be responsible for the high prevalence. Suntaravitun and  
193 Dokmakaw, 2018 reported higher prevalence of intestinal parasites in older age  
194 groups than in younger age groups which was attributed to possible increased  
195 exposure to contaminated soil in older age groups.

196 Mixed infection (polyparasitism) was recorded among pupils examined; infection  
197 with as many as three different intestinal parasites was recorded in children. Mixed  
198 infections could be as a result of the fact that factors such as high temperature and  
199 poor sanitation favour a range of intestinal parasites and as such children are  
200 exposed to infection with more than one parasite. Particularly, *E. histolytica* was  
201 present in all three groups of mixed infections; this could be as a result of the  
202 ability of its cysts to survive harsh environmental conditions leading to a buildup  
203 of the parasite in the environment. Children aged 5-9 years had higher prevalence  
204 of mixed infections although the difference was not significant; and females also  
205 had higher prevalences of mixed infections although the difference was also not  
206 significant.

207

208

209

210

211

212

213

214

## 215 **Toilet system**

216 Children who defecate in the bush had highest prevalence with intestinal parasite.  
217 This could be because after defecating in the open, children tend to wipe their anus  
218 with leaves of plants or paper and may not wash their hand afterwards. Children  
219 who use pit toilet had the second highest prevalence with the parasite, this could be  
220 because children tend to defecate on the mouth of the pit toilet which is sometimes  
221 unroofed and in bad conditions, this attracts flies causing spread of intestinal  
222 parasite eggs and cysts. Akor *et al.*, 2019 also reported higher prevalence of  
223 intestinal parasite in people who practice open defecation.

## 224 **Source of drinking water**

225 Infection rate relating to the source of drinking water showed that pupil that obtain  
226 their drinking water from the stream recorded the highest number of parasitic  
227 infection, this could be as a result of run off of infected soil into stream during  
228 rainfall and also run off water from gutters, drainages and sewages.

229 The ova of the parasites are easily transported by water and other means thus  
230 contaminating the source of drinking water. Ani and Itiba (2015) reported high  
231 prevalence among people using stream and well as primary source of water in  
232 Ebonyi, Abakaliki, Nigeria.

## 233 **Parental occupation/socioeconomic factors**

234 Children are active and playful and children of farmers in particular because of the  
235 nature of the occupation of their parents , stay for long hours in the farmer with  
236 their parents, thereby making them highly prone to contact with soil contaminated  
237 with intestinal parasites.

238

239 **Conclusion**

240 Intestinal parasites are prevalent in Vandeikya LGA, Benue State. Risk factors like  
241 open defecation, use of stream and well water should be minimized in order to  
242 prevent infection.

243

244 **REFERENCES**

245

246

247 Akor, J. O., Obisike, V. U., Omudu, E. A. and Imandeh, G, N. (2019). The  
248 prevalence of polyparasitism in Oju Local Government Area of Benue State,  
249 Nigeria. *International Journal of Tropical Disease & Health* 39(2): 1-6

250 Ani, O. C. and Itiba, O. L. (2015). Evaluation of parasitic contamination from  
251 local sources of drinking-water in Abakaliki, Ebonyi State, Nigeria. *Nigerian*  
252 *Journal of Parasitology* 36(2): 153-58

253 Gebretsadik, D., Metaferia, Y., Seid, A., Fenta, G. M. and Gedefie, A. (2018).  
254 Prevalence of intestinal parasitic infection among children under 5 years of age at  
255 Dessie Referral Hospital: cross sectional study. *Biomedical Central Research Notes*  
256 11, 771

257 Kumma, W. P., Meskele, w. and Admasie, A. (2019). Prevalence of intestinal  
258 parasitic infections and associated factors among food handlers in Wolaita Sodo  
259 University Students Caterings, Wolaita Sodo, Southern Ethiopia: A Cross-  
260 Sectional Study. *Front. Public Health* 7:140-147

261 Praharal, I., Sarkar, R., Ajjampur, S. S., Roy, S. and Kang, G. (2017). Temporal  
262 trends of intestinal parasites in patients attending a tertiary care hospital in south  
263 india: A seven-year retrospective analysis. *Indian Journal of Medical Research*  
264 146(1): 111-120

- 265 Saki, J., Khademvatan, S., Foroutan-Rad, M. and Gharibzadeh, M. (2017).  
266 Prevalence of intestinal parasitic infections in Haftkel County, Southwest of Iran.  
267 *International Journal of Infection* 4(4): e15593
- 268 Suntaravitun, P. and Dokmakaw, A. (2018). Prevalence of intestinal parasites and  
269 associated risk factors for infection among rural communities of Chachoengsao  
270 province, Thailand. *Korean Journal of Parasitology* 56(1): 33-39
- 271 Tegegne, Y., Wondmagegn, T., Worku, L. and Zeleke, A. J. (2018). Prevalence of  
272 Intestinal Parasites and Associated Factors among Pulmonary Tuberculosis  
273 Suspected Patients Attending University of Gondar Hospital, Gondar, Northwest  
274 Ethiopia. *Journal of Parasitology Research* Article ID 9372145
- 275 Valiathan, R., Ashman, M. and Asthana, D. (2016). Effects of age on the immune  
276 system: infants to elderly. *Scandinavian Journal of Immunology* 83(4): 255-266