

Stem Cell Therapy in Treatment of Non-Communicable Diseases: Possibilities and Challenges

Abstract

Over recent years stem cells have stood out as a promising tool for regenerative medicine, providing alternative therapeutic solutions for many non-communicable diseases. Many clinical trials using stem cells or induced pluripotent stem cells are focused on the refit and regeneration of various tissues and organs in degenerative diseases, whose current treatment only succeeds in delay down the progression of the disease. This review summarizes several current clinical and nonclinical information on the use of embryonic stem cells (ESCs), induced pluripotent stem cells (IPSCs) and mesenchymal stem cells (MSCs) in various diseases. The aim of this review was to expand on the background and therapeutic potential of ESCs, MSCs, and IPSCs whilst linking this to their use within disease therapy with a specific focus on diabetes, kidney disease, and cardiovascular disease with future possibilities and challenges. Also aimed to explain the benefits of transplantation with side effects shortly after transplant and later and interruptions, possibilities and challenges of transplantation.

Key words: Regenerative medicine, Stem cells, Stem cell therapy, Pluripotent stem cells, Embryonic stem cells, Mesenchymal stem cells.

Introduction

Stem cells are undifferentiated cells of a multicellular organism, that can turn into specific cells, as the body needs them. Stem cells have the potential to revolutionize tissue regeneration and engineering (1). Cells in the body have specific purposes, but stem cells are cells that do not yet have a specific role and can become almost any cell that is required. There are three main types of stem cells depending on source: embryonic stem cells (ESCs), induced pluripotent stem cells (iPSCs) and mesenchymal stem cells (MSCs) (2).

For the first time in 1981, researchers could isolate stem cells from mouse embryos. More accurate studies on the biology of mouse stem cells led to discovery of methods for separation of stem cells from the human embryo in 1998. Another name of stem cell therapy is regenerative medicine, it's promotes the repair response of diseased, dysfunctional or injured tissue using stem cells or their derivatives (3). The origin and function of these stem cells varies but they all hold diagnostic and therapeutic potential. A potential route of treatment is using stem cells that have the capability of differentiating into healthy tissue, replacing any lost through disease manifestation and thus justifying the avoidance of lifelong expensive treatments (4).

Stem cells provide new cells for the body as it grows and replace specialized cells that damaged or lost. They have two unique properties that, they can divide repeatedly to produce new cell, and they can change into the other types of cell that make up the body (5).

Classification based on potency

Stem cells can be classified based on potency, by the extent to which they can differentiate into different cell types. The four main classifications are totipotent, pluripotent, multipotent, or unipotent.

Totipotent: The ability to differentiate into all possible cell types. Examples are the zygote and the first few cells that result from the division of the zygote (5).

Pluripotent: The ability to differentiate into almost all cell types. Examples are embryonic stem cells and cells that are isolated from the mesoderm, endoderm, and ectoderm germ layers that are formed in the beginning stages of embryonic stem cell differentiation (6).

Multipotent: The ability to differentiate into a closely related family of cells. Examples include hematopoietic (adult) stem cells that can become red and white blood cells or platelets.

Unipotent: The ability to only produce cells of their own type. Examples include (adult) muscle stem cells (7).

Classification based on their sources

Embryonic stem cells: Embryonic stem cells are self-replicating cells that are potentially immortal (7). They are derived from embryos at a developmental stage before the time of implantation occur in the uterus. They can change into any cell in the body, so these stem cells are said to be pluripotent (8).

Adult stem cells: Adult stem cells are undifferentiated totipotent or multipotent cells that are found in the body after embryonic development and multiply by dividing cells. The main role of adult stem cells in vivo is to preserve and improve the tissue in which they are found. They can replace blood stem cells (or “blood formation”) only with blood cells and various types of skin cells (or “epithelium”) (8).

Induced pluripotent stem cells: Recently, a third type of stem cell has emerged that has properties like embryonic stem cells. Scientists have created these induced pluripotent stem cells (iPS cells) by controlling the expression of specific genes and reprogramming somatic cells into pluripotency (9). They are also pluripotent and can develop into cells of any type (10).

Applications of stem cell therapy

The goal of stem cell therapy is to treat damaged tissue that can't heal itself. Recent stem cell research often encourages patients who have not been treated for diseases in order to alleviate the symptoms of chronic diseases. Stem cell therapy involves more than just implanting cells into the body and growing new healthy tissue. It may also be possible to gratify stem cells already in the body to work overtime and produce new tissue (11).

Diseases and conditions where stem cell treatment is being investigated include: Non-union/Delayed Union Fracture, Osteonecrosis or Avascular Necrosis (AVN), Knee Cartilage Defect, Rheumatoid Arthritis (RA), Spinal Cord injury, Spinal Fusion Treatment, Cerebral Palsy, Autism, Motor Neuron Disease, Multiple Sclerosis, Parkinson's Disease, Alzheimer/Dementia

disease, Cerebellar Atrophy, Cerebellar Ataxia, Spinal Muscular Atrophy, Down Syndrome, Optic Nerve Damage, Retinitis Pigmentosa, Macular Degeneration, Dystrophy Glaucoma Disease, Diabetes (Type 1 & 2) (12), Acute/Chronic Liver Disease, Muscular Dystrophy, Acute/Chronic Kidney Disease (13), Peripheral Arterial Disease, Myocardial Infarction, Lung Disease, Erectile Dysfunction, Anti- Aging Treatment, Scleroderma Disease, Skin Replacement, Male Infertility, Female Infertility, Breast cancer, Scar, Colon cancer, etc. (14).

Possibilities in some most occurring non communicable diseases

Stem cell therapy for kidney disease

Kidney diseases are caused by damage to nephrons, which can be sudden and short lived called acute kidney disease or slow and progressive called chronic kidney disease. Chronic kidney disease can lead to renal failure and is fatal if not treated. Scientists are studying how the kidneys can be restored and the types of kidney cells involved in this process (15).

After variety of insults the kidney has the capacity for regeneration. The factors that caused kidney damage over the past few decades have been carefully studied. There is currently no FDA approved stem cell therapy for kidney disease (16). However, clinical trials are being conducted to determine whether stem cell-based kidney therapy is safe and effective in humans. Scientists have already successfully created an artificial rat kidney that produces urine once transplanted into the animal, making artificial organ transplantation a highly possible reality for human kidney (13, 17).

Types of stem cells investigated to regenerate damaged tissue

Significant advances have been made in promoting stem cell products as a possible treatment of kidney disease, restoring various types of stem cells from renal function in preclinical models of acute and chronic renal failure (18). Both preclinical reports and clinical trials of stem cells used for treatment kidney disease are increasing rapidly. Different types of stem cells, ranging from mesenchymal stem cells (MSCs), adipose stem cells (AdSCs), hematopoietic stem cells (HSCs), amniotic fluid stem cells (AFSCs), renal progenitors (RPs) and so forth, can stimulate renal repair in vivo in models of acute and chronic kidney failure (16).

The promise of stem cell therapies in pre-clinical models of kidney diseases is yet to be translated into more persuasive proof of clinical efficacy. Several clinical trials have confirmed the safety and tolerability of stem cells, and of MSC-based therapies, in patients with renal diseases and kidney transplants. However, long-term monitoring is recommended to rule out the potential risk of cancer and of developing anti-HLA antibodies (19).

Stem cell technology for the treatment of diabetes

Diabetes is a disease with high blood sugar. Most of the glucose comes from the food we eat. The hormone used to transport glucose to the cell as energy is insulin. When our body cannot produce insulin, it is called type 1 diabetes. In type 2, glucose is not available due to insulin resistance (20).

In the past, numerous diabetes treatment technologies have been used, including increased insulin delivery and glucose monitoring systems, new methods for the complete transplantation of the pancreas and graft, and the formation of B cells from the pancreatic ducts or stem cells. Now-a-days these two conditions of diabetes are treated by stem cell technology (21).

Types of stem cells investigated to regenerate damaged tissue

Stem-cell therapy means the replacement of diseased cell or missing cells from progeny of pluripotent or multipotent cells. Many groups of cells are used to turn into beta cells of the pancreas, which produce insulin during differentiation. Both embryonic stem cells (derived from the inner cell mass of a blastocyst) and adult stem cells (found in the postnatal organism) have been used to produce β -cells or otherwise restore the functioning of β -cell (22). Embryonic stem cells (ESC): Stem cells follow appropriate developmental pathway in order become insulin producing cells by using embryonic stem cells transfusing with insulin promoter, resulting insulin producing cells in mouse ESC. Which permitted them to make insulin producing cells (21).

Induced pluripotent cells (IPSC): IPS has high reproducibility and pluripotency. These cells can be divided into cells that produce insulin. So, we can also make the pancreatic beta cells form these cells that can be used for the treatment of diabetes (22).

Mesenchymal stem cell (MSC) therapy: Stem Cell Therapy is an alternative to small islet cell transplantation in patients with type 2 diabetes. Mesenchymal stem cell (MSC) can be obtained from patients with autologous transplantation. However, autologous MSCs from diabetic patients are still remarkably different from ESCs, because of prolonged exposure to hyperglycemia (23). By using these different kinds of stem cell technologies, we can make the insulin producing cells that will be helpful in the cure of diabetes that is worldwide disease. Type 1 diabetes can successfully be treated by using β cell from stem cells where type 2 diabetes is treated by using β cell in combination with drug therapy (24).

Stem cells and cardiac repair

Cardiovascular diseases (CVD), hypertension, coronary heart disease (CHD), stroke, and cardiac arrest (CHF) were the leading causes of death. Finally, stem cells can satisfy a large unmet clinical need and improve the quality of life of millions of people with cardiovascular diseases (25). Reliable evidence suggests that stem cells promise to be a treatment for damaged myocardial regeneration. Ischemic heart failure occurs when there is a lack of oxygen in the tissues of the heart (26).

Types of stem cells investigated to regenerate damaged myocardial tissue

Adult and embryonic stem cells have been investigated to regenerate damaged myocardial tissue in animal models and in a limited number of clinical studies. Specific considerations for the application of various cell types will be discussed in the following sections (27). Embryonic stem (ES) cells: Pluripotent ES cells can produce various types of cells that repair damaged myocardial tissue, including myocardial cells, endothelial cells and smooth muscle cells. For this purpose, it has been shown that mouse and human ES cells spontaneously differentiate in vivo to form endothelial muscle and smooth muscle cells in the body. Human ES cells differentiate into myocytes with the structural and functional properties of myocardial cells. Moreover, ES cells that were transplanted into ischemically injured myocardium in rats differentiated into normal myocardial cells that remained viable for up to four months suggesting that these cells may be candidates for regenerative therapy in humans (28).

Other diseases

Stroke

Stroke triggers the loss of neurons and glial cells in large numbers. In the treatment of this disease, cell therapy opens fresh horizons by facilitating the process of neuronal regeneration. Animal studies and several preclinical studies verify the effectiveness of cell therapy in post-stroke functional enhancement (3).

Spinal cord injury

Spinal cord injury is one of the serious neurological damage caused by neuronal tissue loss and consequently sensory and motor function loss. There is no therapy for this damage to be regenerated. This harm can be remedied by replacing stem or progenitor cells (29).

Inflammatory bowel disease

Crohn and ulcerative colitis were referred to as inflammatory diseases of the intestine. The precise cause of these illnesses is still unknown, but one of their causes is immune system dysfunction. Because stem cells are immunoregulatory cells, and also because of their capacity to transdifferentiate and fuse cells, they appear to have a beneficial impact on improving these illnesses (30).

Liver diseases

Stem cell transplantation has now been proposed in the therapy of cirrhosis as a novel technique. For this purpose, various types of stem cells such as embryonic stem cells, mesenchymal stem cells, annex stem cells and endothelial progenitor cells were used in laboratory studies. Also, laboratory studies have shown that primary hepatocytes can be replaced in liver, spleen, peritoneal cavity and other sites outside the liver (31).

Benefits of stem cell therapy

With so many treatment options out there, you may be surprised what benefits stem cell therapy provides. Some of the benefits include minimal risk, minimal recovery time and minimal worry. Here are 5 more specific benefits to be aware of- no need of surgery and avoid its risks and complications because it is an invasive, non-surgical procedure, required less post-procedural recovery time, does not require the use of general anesthesia, there is no risk of rejection because this therapy uses the biologics extracts from the patient, no risk of communicable disease transmission (32).

Side effects of stem cell transplant

After the transplantation many of the side effects are happens to the patient shortly after the transplant and later. These side effects are briefly described below-

Shortly after the transplants

Many of the problems can happens shortly after the transplants this includes- mouth and throat pain, nausea and vomiting, infection, bleeding and transfusions, interstitial pneumonitis, other lung problems, graft failure and graft-versus-host disease. When the body does not accept the new stem cells the graft fails. Graft failure is more common when the patient and donor are not well matched and when patients get stem cells that have had the T-cells removed. Graft failure can lead to serious bleeding and/or infection and graft-versus-host disease (GVHD) can happen in allogeneic transplants when the immune cells from the donor see the recipient's body as foreign. Doctors think of GVHD as acute or chronic. Acute GVHD starts immediately after transplant and lasts a short time. Chronic GVHD starts later and lasts a long time (33).

Acute GVHD

The average time is about 25 days, but acute GVHD may occur between 10 and 90 days after transplantation. Skin rash, redness of the skin, redness of the palm and initial signs. It can spread throughout your body. Other symptoms include nausea, vomiting, abdominal cramps, diarrhea (moisture, sometimes bleeding), loss of appetite, yellowing of the skin and eyes (jaundice), abdominal pain (abdominal pain) and weight loss (34).

Chronic GVHD

90 to 600 days after the stem cell transplant, chronic GVHD can start anywhere. Rash on the palms of the hands or the soles of the feet are the first signs. The rash can spread and is usually itchy and dry. In severe cases, the skin may blister and peel, like a bad sunburn. Other symptoms of chronic GVHD can include: enlarged liver, bloated abdomen (belly), pain in the upper right part of the abdomen (belly), increased levels of liver enzymes in the blood (seen on blood tests), the skin feels tight, dry burning eyes, dryness or painful sores in the mouth, burning sensations when eating acidic foods, bacterial infections, and blockages in the smaller airways of the lungs, etc. (35).

Problems that may show up later

Organ damage, Relapse (the cancer comes back), Secondary (new) cancers, Abnormal growth of lymph tissues, infertility (the inability to produce children), hormone changes, such as changes in the thyroid or pituitary gland, cataracts (clouding of the lens of the eye, which causes vision loss) (36).

Dietary concerns during stem cell transplant

Patients with stem cell transplantation are very sensitive to microorganisms that can be transmitted through food and beverages. To provide a more protected environment, transplant patients are maintained on a diet upon admission which is secure for the patient. The stem cell transplant diet restrictions continue until the patient's ANC (ANC: The absolute neutrophil count, the number of white blood cells (WBCs) that are neutrophils) is greater than 500 for three consecutive days and if graft vs. host disease is not present (37). Patients who underwent autologous or allogeneic HSCT (hematopoietic stem cell therapy) had more pronounced changes in diet acceptance and longer hospital stays, as well as a greater delay from transplantation until engraftment because they are at increased nutritional risk due to the underlying disease, high metabolic demand and complications related to the conditioning regimen, which mostly affect the gastrointestinal tract, and are thus able to cause symptoms, such as nausea, vomiting, mucositis, odynophagia, diarrhea, abdominal pain and constipation, that make the ingestion of food and absorption of nutrients worse (38, 39). When the appropriate transplantation occurs, the special diet is terminated. The special diet will be stopped when adequate engraftment occurs; however, it is a good idea to continue to use safe food handling and preparation even after the special diet restrictions have ended. Diet guidelines should be maintained before and after therapy (40).

Interruptions, challenges and prospects

Though stem cells offer breathtaking commitment for coming therapies, significant technical hurdles remain. One of the major interruptions is that mass proliferation of stem cells to generate enough quantities of tissue is required for useful transplant purposes. Secondly differentiated stem cells should be able to integrate into the surrounding tissue after transplantation and their functionality for the total duration of the patient's life is another area of concern along with the

fears of immune rejection. Uses of autologous adult stem cells and lifelong immunosuppressive medicine can reduce the harmful side effects. Another major concern is the development of cancer after SCT due to self-renewal and uncontrolled proliferation capabilities of stem cells, in this way making them able to malignant transformation as CSCs. Another major complexity after allogeneic hematopoietic stem cell transplantation is graft-versus-host disease is due to donor T-cell recognition of recipient alloantigen's. Autologous hematopoietic stem cell transplantation also exposes the development of a syndrome like allogeneic graft-versus-host disease causing serious disease in the GI tract (41).

Though this field is still in its primary stage, scientists are hopeful to use iPSCs in transplantation therapy. Use of retroviral vectors to initiate transcription factors for reprogramming objective can be harmful itself with a raise risk of cancer and viral disease development therefore researchers are currently investigating non-viral allotment techniques. Although the nature of the reprogramming method is still not surely clear and the developmental potential of iPSCs derived from various tissues using various methods is unknown. Another main challenge is to recognize adult or cancer stem cells in a tissue population (42).

Many preclinical studies regarding SCT have moved to clinical trials and translational stage. Such as to promote recovery of the damaged heart after myocardial infarction (MI), stem cells have been delivered by intracoronary infusion. Where, recent reviews of these clinical trials report that less than half of the trials found only small improvements in cardiac function. Significant tasks yet remain to enhance the usefulness of stem cell therapy for CVD including improved identification, recruitment, and in vitro growth of autologous stem cells, besides identification of mobilizing and homing agents, and development of novel culture conditions to improve stem cell survival and engraftment. Other clinical trials are planned to address treatment options for diabetes and kidney diseases (43).

To date (08/04/2019), over 880 MSC-based clinical trials, complete or ongoing, have been registered in the US National Institutes of Health database, including hematological diseases, graft versus host disease, diabetes, organ transplantation, inflammatory diseases, and diseases in the lung, liver, bone as well as cardiovascular, neurological and autoimmune diseases. Most of the above clinical trials are carried out in early phases (phase I–II), suggesting that the efficiency of SC treatment remains to be further investigated in the long-term (44).

Conclusion

There is a lot of research on investigating how stem cells can be used to treat different types of diseases. It is expected that; stem cell therapy might be used to develop for many types of diseases in future for which there are no effective treatments now. The obtained pluripotent stem cells and adult stem cells are important candidates for regenerative therapy because of the nature of self-healing and the wide range of properties of pluripotency. Stem cells have been tested for use in various diseases, such as diabetes, kidney disease, spinal cord injuries, heart disease, stroke and Parkinson's disease, as well as various forms of blood diseases. Where kidney disease has been studied, there is no proven treatment for this disease with stem cells because of its complex structure. On the other hand, type 1 diabetes can successfully treated by using β cell from stem cells where type 2 diabetes is treated by using β cell in combination with drug therapy. And for the patient with different heart disease stem cell therapeutics are very promising. As stem cell therapy has many benefits for treating disease it also has some side effects too. Stem cells have a bright future for the therapeutic world according to different literature. We hope to see new horizon of therapeutics in the form of bone marrow transplant, skin replacement, organ development, and replacement of lost tissue such as hairs, tooth, retina and cochlear cells which will add new dimension to the treatment and healing process of ailing patients.

Ethics declarations

Ethics approval: Not applicable.

References

1. Menendez JA, Joven J. Energy metabolism and metabolic sensors in stem cells: the metabostem crossroads of aging and cancer. *Oxidative Stress and Inflammation in Non-communicable Diseases-Molecular Mechanisms and Perspectives in Therapeutics*: Springer; 2014. p. 117-40.
2. Marks PW, Witten CM, Califf RM. Clarifying stem-cell therapy's benefits and risks. *N Engl J Med*. 2017;376(11):1007-9.
3. Larijani B, NASLI EE, Amini P, Nikbin B, Alimoghaddam K, Amiri S, et al. Stem cell therapy in treatment of different diseases. 2012.

4. Schöler HR. The potential of stem cells: An inventory. *Humanbiotechnology as social challenge*: Routledge; 2016. p. 45-72.
5. Ghajar CM, Peinado H, Mori H, Matei IR, Evason KJ, Brazier H, et al. The perivascular niche regulates breast tumour dormancy. *Nature cell biology*. 2013;15(7):807.
6. Ulloa-Montoya F, Verfaillie CM, Hu W-S. Culture systems for pluripotent stem cells. *Journal of bioscience and bioengineering*. 2005;100(1):12-27.
7. Jeffrey M, Ponzetti M, David S. stem cells An Interactive Qualifying Project Report, Submitted to the Faculty of worcester polytechnic institute In partial fulfillment of the requirements for the Degree of Bachelor of Science, 2006. *AJPCT*. 2014;2(7):919-30.
8. Crispi V, Matsakas A. Duchenne muscular dystrophy: genome editing gives new hope for treatment. *Postgraduate medical journal*. 2018;94(1111):296-304.
9. Bolós V, López Á, Aparicio LA. Cancer Stem Cells as a New Opportunity for Therapeutic Intervention. *Cancer Stem Cells Theories and Practice*: IntechOpen; 2011.
10. Bangladesh Demographic and Health Survey Bangladesh: Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh. 2014.
11. Trounson A, McDonald C. Stem cell therapies in clinical trials: progress and challenges. *Cell stem cell*. 2015;17(1):11-22.
12. Libby P, Nathan DM, Abraham K, Brunzell JD, Fradkin JE, Haffner SM, et al. Report of the national heart, lung, and blood institute-national institute of diabetes and digestive and kidney diseases working group on cardiovascular complications of type 1 diabetes mellitus. *Circulation*. 2005;111(25):3489-93.
13. Ikarashi K, Li B, Suwa M, Kawamura K, Morioka T, Yao J, et al. Bone marrow cells contribute to regeneration of damaged glomerular endothelial cells. *Kidney international*. 2005;67(5):1925-33.
14. Dulak J, Szade K, Szade A, Nowak W, Józkwicz A. Adult stem cells: hopes and hypes of regenerative medicine. *Acta Biochimica Polonica*. 2015;62(3).
15. Togel F, Hu Z, Weiss K, Isaac J, Lange C, Westenfelder C. Administered mesenchymal stem cells protect against ischemic acute renal failure through differentiation-independent mechanisms. *American Journal of Physiology-Renal Physiology*. 2005;289(1):F31-F42.

16. Kale S, Karihaloo A, Clark PR, Kashgarian M, Krause DS, Cantley LG. Bone marrow stem cells contribute to repair of the ischemically injured renal tubule. *The Journal of clinical investigation*. 2003;112(1):42-9.
17. Hickson LJ, Eirin A, Lerman LO. Challenges and opportunities for stem cell therapy in patients with chronic kidney disease. *Kidney international*. 2016;89(4):767-78.
18. Hauser PV, De Fazio R, Bruno S, Sdei S, Grange C, Bussolati B, et al. Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. *The American journal of pathology*. 2010;177(4):2011-21.
19. van Koppen A, Joles JA, van Balkom BW, Lim SK, de Kleijn D, Giles RH, et al. Human embryonic mesenchymal stem cell-derived conditioned medium rescues kidney function in rats with established chronic kidney disease. *PloS one*. 2012;7(6):e38746.
20. Janghorbani M, Van Dam RM, Willett WC, Hu FB. Systematic review of type 1 and type 2 diabetes mellitus and risk of fracture. *American journal of epidemiology*. 2007;166(5):495-505.
21. Kelly OG, Chan MY, Martinson LA, Kadoya K, Ostertag TM, Ross KG, et al. Cell-surface markers for the isolation of pancreatic cell types derived from human embryonic stem cells. *Nature biotechnology*. 2011;29(8):750.
22. Zhang D, Jiang W, Liu M, Sui X, Yin X, Chen S, et al. Highly efficient differentiation of human ES cells and iPS cells into mature pancreatic insulin-producing cells. *Cell research*. 2009;19(4):429.
23. Urban VS, Kiss J, Kovacs J, Gocza E, Vas V, Monostori É, et al. Mesenchymal stem cells cooperate with bone marrow cells in therapy of diabetes. *Stem cells*. 2008;26(1):244-53.
24. Zeeshan N, Naveed M, Irshad DFA, Ahsan A, Abrar M, Ghafoor S. *Journal of Cell Science & Therapy*. 2017.
25. Terzic A, Behfar A. Regenerative heart failure therapy headed for optimization. *European heart journal*. 2014;35(19):1231.
26. Min J-Y, Yang Y, Converso KL, Liu L, Huang Q, Morgan JP, et al. Transplantation of embryonic stem cells improves cardiac function in postinfarcted rats. *Journal of Applied Physiology*. 2002;92(1):288-96.
27. Vittet D, Prandini M, Berthier R, Schweitzer A, Martin-Sisteron H, Uzan G, et al. Embryonic stem cells differentiate in vitro to endothelial cells. 2011.

28. Bartunek J, Behfar A, Dolatabadi D, Vanderheyden M, Ostojic M, Dens J, et al. Cardiopoietic stem cell therapy in heart failure: the C-CURE (Cardiopoietic stem Cell therapy in heart failURE) multicenter randomized trial with lineage-specified biologics. *Journal of the American College of Cardiology*. 2013;61(23):2329-38.
29. Nandoe Tewarie RS, Hurtado A, Bartels RH, Grotenhuis A, Oudega M. Stem cell-based therapies for spinal cord injury. *The journal of spinal cord medicine*. 2009;32(2):105-14.
30. Yan L, Cai C, Li J, Xu S, Chang Q, Li Y, et al. Present status and perspectives of stem cell-based therapies for gastrointestinal diseases. *Stem Cell Reviews and Reports*. 2009;5(3):278-82.
31. Meier RP, Müller YD, Morel P, Gonelle-Gispert C, Bühler LH. Transplantation of mesenchymal stem cells for the treatment of liver diseases, is there enough evidence? *Stem cell research*. 2013;11(3):1348-64.
32. Murnaghan M, Li G, Marsh DR. Nonsteroidal anti-inflammatory drug-induced fracture nonunion: an inhibition of angiogenesis? *JBJS*. 2006;88(suppl_3):140-7.
33. Joggerst SJ, Hatzopoulos AK. Stem cell therapy for cardiac repair: benefits and barriers. *Expert reviews in molecular medicine*. 2009;11.
34. Harrison N, Mitterbauer M, Tobudic S, Kalhs P, Rabitsch W, Greinix H, et al. Incidence and characteristics of invasive fungal diseases in allogeneic hematopoietic stem cell transplant recipients: a retrospective cohort study. *BMC infectious diseases*. 2015;15(1):584.
35. Jagasia M, Arora M, Flowers ME, Chao NJ, McCarthy PL, Cutler CS, et al. Risk factors for acute GVHD and survival after hematopoietic cell transplantation. *Blood*. 2012;119(1):296-307.
36. Couriel DR, Hosing C, Saliba R, Shpall EJ, Anderlini P, Rhodes B, et al. Extracorporeal photochemotherapy for the treatment of steroid-resistant chronic GVHD. *Blood*. 2006;107(8):3074-80.
37. So EJ, Lee JS, Kim JY. Nutritional intake and nutritional status by the type of hematopoietic stem cell transplantation. *Clinical nutrition research*. 2012;1(1):3-12.
38. de Defranchi RLB, Bordalejo A, Cañueto I, Villar A, Navarro E. Evolution of nutritional status in patients with autologous and allogeneic hematopoietic stem cell transplant. *Supportive care in cancer*. 2015;23(5):1341-7.

39. Andersen S, Brown T, Kennedy G, Banks M. Implementation of an evidenced based nutrition support pathway for haematopoietic progenitor cell transplant patients. *Clinical nutrition*. 2015;34(3):536-40.
40. Prockmann S, Freitas AHR, Ferreira MG, Vieira FGK, de Salles RK. Evaluation of diet acceptance by patients with haematological cancer during chemotherapeutic treatment. *Nutricion hospitalaria*. 2015;32(2):779-84.
41. Nygaard GO. Cognition, disease activity and MRI changes in early multiple sclerosis. 2016.
42. Titomanlio L, Kavelaars A, Dalous J, Mani S, El Ghouzzi V, Heijnen C, et al. Stem cell therapy for neonatal brain injury: perspectives and challenges. *Annals of neurology*. 2011;70(5):698-712.
43. Neeri F. *BRITISH BIOMEDICAL BULLETIN*. 2015.
44. Rota C, Morigi M, Imberti B. Stem Cell Therapies in Kidney Diseases: Progress and Challenges. *International journal of molecular sciences*. 2019;20(11):2790.