

## Original Research Article

### Physiological and biochemical changes under salinity and drought stress in ricebean [*Vigna umbellata* (Thunb.) Ohwi and Ohashi] seedlings

---

#### ABSTRACT

**Aims:** To study the effect of iso-osmotic potentials of drought and salinity during seedling growth stage in ricebean.

**Study design:** Completely randomised design.

**Place and Duration of Study:** The lab experiment was conducted during the year of 2017-2018 and 2018-2019 in ricebean variety Bidhan 1 at Department of Plant Physiology, Bidhan Chandra Krishi Viswavidyalaya, Kalyani, Nadia, West Bengal, India.

**Methodology:** For studying the effect of iso-osmotic potential of salinity and drought stress, the solutions of NaCl and PEG 6000 with -0.2, -0.4 and -0.8 MPa osmotic potential were used and the experiment was conducted in sand culture using modified Hoagland solution [1] under laboratory condition of diffused light, at around 80±1% relative humidity (R.H.) and at a temperature of 28±1°C.

**Results:** All the biochemical parameters under study, in general were adversely affected by the both stress with the effects being more drastic as the intensity of stress increased. The highest intensity of salinity stress was found to produce more adverse effects than drought in respect of RLWC, leaf chlorophyll as well as protein content in leaves of ricebean in the present experiment. While the content of soluble sugar, starch and phenol in the leaf were more drastically affected by drought stress.

**Conclusion:** The drought stress was found to register more drastic effects on seedling growth as compared to iso-osmotic potential of salinity stress, especially, at the highest intensity of stress in ricebean cultivar Bidhan 1.

*Key words: Ricebean, Salinity stress, Drought stress, Leaf Protein, Lipid peroxidation, Total Phenol, Leaf starch and Total soluble sugar.*

## 1. INTRODUCTION

Salinity stress can affect plants initially by creating an osmotic stress then it induces ion toxicity that lead to cyto-toxicity, metabolic impairment, nutrient imbalance and finally death of the plant. Initially, the presence of salts in high concentration makes very difficult for plants to withdraw water from soil due to very low osmotic potential. In effect, the plants suffer from a sort of osmotic stress which causes yield reduction. At the later stages of stress, due to the absorption of sodium and chloride ions in high concentration plants suffer from cyto-toxicity which result in reduction of growth, leaf burn and plant death. The presence of high concentration of  $\text{Na}^+$  and  $\text{Cl}^-$  ion also reduces the availability of other ions like  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , thus, causing nutritional disorders.

Drought is a meteorological term that normally occurs under depleting soil moisture and the intensity of drought increases under atmospheric conditions conducive to increased water loss by transpiration and/ or evaporation. Water deficit is one of the major abiotic stresses, which adversely affects crop growth and yield. These changes are mainly related to altered metabolic functions, one of those is either loss of or reduced synthesis of photosynthetic pigments, uptake and translocation of ion, carbohydrate biosynthesis, nutrient metabolism and synthesis of growth promoters. These changes in the metabolic functions and synthesis of photosynthetic pigments are closely associated with biomass production in plant [2]. A common adverse effect of water stress on crop plants is the reduction in fresh and dry biomass [3]. Plant productivity under drought stress is strongly related to the processes of dry matter partitioning and temporal biomass distribution [4].

The present experiment has been designed to study the comparative effects of different levels of salinity and drought stress on some physiological and biochemical parameters of ricebean during seedling growth stage.

## 2. MATERIAL AND METHODS

For studying the effect of salinity and drought stress on seedling growth of ricebean the experiment was conducted in sand culture using modified Hoagland solution [1] under laboratory condition of diffused light, at around  $80\pm 1\%$  relative humidity (R.H.) and at a temperature of  $28\pm 1^\circ\text{C}$ . For this purpose, the seeds of ricebean cultivar Bidhan-1 were surface sterilized with 0.1% (w/v)  $\text{HgCl}_2$  for 3 minutes followed by thorough washing in distilled water. Then the seeds were germinated for 48 hours at  $28\pm 1^\circ\text{C}$  using glass distilled water. The pre-germinated seeds were then transferred to plastic beakers of capacity one litre containing neutral sand. Five pre-germinated seeds were transferred to each beaker. Finally three healthy seedlings were allowed to grow each beaker. The seedlings were grown in presence of full strength Hoagland solution prepared as per modification of [1] for 14 days. The nutrient medium was supplemented at 3 days interval. At the age of 14 day the seedlings were subjected to salinity and drought treatments. For this purpose, the appropriate amounts of NaCl and PEG 6000 calculated as per Sosa et al. [5] to obtain the osmotic potential ( $\Psi$ ) of -0.2, -0.4 and -0.8 MPa were mixed with modified Hoagland nutrient solution. Thus, the drought and salinity stress with iso-osmotic potentials were created in the present experiment. A control set having  $\Psi$ s equivalent to 0.0 Mpa osmotic potential without containing NaCl or PEG was also maintained similarly for comparison of results. Observations on different growth and biochemical parameters were recorded on 9 days after treatment application.

The dry weight of seedling were determined by harvesting the seedlings from the beaker at 9 days after treatment (DAT) and then drying in oven at  $80^\circ\text{C}$  till constant weight. Before that the fresh weight in each case also recorded. The chlorophyll content in the leaf was

measured as per Arnon [6], while lipid peroxidation was estimated as per Heath and Packer [7]. Relative leaf water content (RLWC) was determined as per Perez et al. [8]. The content of total phenol, soluble protein and total soluble sugar in the leaves of the seedlings were estimated following the methods of McDonald et al. [9], Lowry et al. [10] and Yoshida et al. [11] respectively.

The mean data in all the cases were subjected to statistical analysis following completely randomised design using INDOSTAT version 7.1 software.

### 3. RESULTS AND DISCUSSION

The total fresh weight of seedling significantly decreased (Table no-1) under all the treatments of drought and salinity stress as compared to that of control. The level of such decrease was higher as the osmotic potential of the growing medium decreased more. Earlier the adverse effects of salinity stress [12, 13] and drought stress [14, 15] on seedling growth in legumes were reported by different workers. Jeannette et al. [13] reported that total fresh weight of root and shoot of cultivated accessions of cowpea was reduced with increased salt stress. Earlier Bibi et al. [16] in sorghum and Khan et al. [17] in wheat reported that the fresh weight of seedling decreased with an increase in PEG concentration. However, in the present experiment drought stress was found to produce more adverse effects on total fresh weight of seedling of ricebean. Thus, the highest intensity (-0.8Mp) of drought and salinity stress led to a reduction of seedling fresh weight by 34.78 and 37.89%, respectively over that of unstressed control.

The dry weight of seedling also significantly decreased under all the treatments of drought and salinity stress as compared to that of control (Table no-1). Like the fresh weight, the level of such decrease in dry weight was also higher as the osmotic potential of the growing medium decreased more. Earlier reduction in dry weight was reported by Anaytullah [18] in rice and in cultivars of blackgram [15] under stress. Gamze [14] showed increasing PEG concentration inhibited seedling fresh and dry weight in pea seedling. The total dry weight registered more adverse effects of drought stress in comparison with salinity stress at iso-osmotic potentials. Thus, the highest intensity (-0.8Mp) of drought and salinity stress led to a reduction of seedling dry weight by 27% and 32%, respectively over that of unstressed control.

**Table-1. Effect of salinity and drought stress on seedling growth in ricebean cv. Bidhan 1**

Treatments	Total fresh weight (gm)	Total dry weight (gm)
<b>Control</b>	1.286	0.041
<b>50 mM NaCl</b>	1.113	0.038
<b>100 mM NaCl</b>	1.044	0.035
<b>200 mM NaCl</b>	0.839	0.029
<b>10% PEG</b>	0.963	0.035
<b>12% PEG</b>	0.827	0.029
<b>18% PEG</b>	0.799	0.028
<b>C.D. (P=0.05)</b>	0.069	0.008

The total chlorophyll content of leaf significantly decreased under all the treatments of drought and salinity stress as compared to that of control, except for PEG 10 % where the mean value slightly exceeded that of control. In all the other cases, the level of decrease in chlorophyll content was higher as the intensity of stress increased (Table no-2). Earlier the adverse effects of drought stress and salinity stress on chlorophyll content were reported by

Pratap and sharma [15], El-Sayed [19] and Aniat-ul-Haq [20]. They also concluded that such decrease in chlorophyll content also resulted in decrease of biomass production. The total chlorophyll content registered more adverse effects of salinity stress in comparison with drought stress at iso-osmotic potentials. The variety Bidhan 1 recorded 35.37 % and 28.51 % reduction in total chlorophyll content under a treatment of 200 mMNaCl and 18 % PEG producing an osmotic potential of -0.8 MPa, respectively, over that of control.

The starch content in the leaves of ricebean seedlings significantly decreased under all the treatments of drought and salinity stress as compared to that of control (Table no-2). This decrease in leaf starch might be attributed to decrease in photosynthetic pigment under stress. Previously, Mohammadkhani and Heidari [21] observed in maize seedling that higher amount of soluble sugars and a lower amount of starch in the leaves under drought stress and concluded the increase in sugar concentration might be a result from the degradation of starch. The variety Bidhan-1 recorded 66.56 % and 68.89 % reduction in starch content in 200 mMNaCl and 18 % PEG producing an osmotic potential of -0.8 MPa, respectively, over that of control. The level of such decrease was higher as the osmotic potential of the growing medium decreased more. The drought stress was found to produce more adverse effects on total content of leaf starch of ricebean in the present experiment.

In case of total soluble sugar content in leaf the lowest intensity of both salinity and drought stress led to an, increase over unstressed control. This increase in sugar content under stress might contribute for osmotic adjustment under stress-induced osmotic shock which was also proposed by Garg et al. [22] and Munns [23]. However, as the stress intensity increased in the present experiment the sugar content significantly decreased under all drought and salinity treatments (Table no-2). The level of such decrease was higher as the osmotic potential of the growing medium decreased more. Earlier the adverse effects of salinity stress [24] and drought stress [25, 26 and 27] on leaf sugar in legumes were reported by different workers. The drought stress was found to produce more adverse effects on total content of sugar in leaf of ricebean in the present experiment.

**Table 2. Effect of salinity and drought stress on contents of chlorophyll, starch and total soluble sugar in the leaves of ricebean cv. Bidhan 1**

Treatments	Chlorophyll <sup>a</sup>	Starch <sup>b</sup>	Sugar <sup>c</sup>
Control	2.77	662.74	83.23
50 mM NaCl	2.36	577.94	111.46
100 mM NaCl	2.35	323.54	83.89
200 mM NaCl	1.79	221.78	75.56
10% PEG	2.89	515.32	108.75
12% PEG	2.53	302.67	81.60
18% PEG	1.98	206.13	43.55
C.D. (P=0.05)	0.50	21.63	9.38

<sup>a</sup> Data expressed as mg g<sup>-1</sup> fresh weight

<sup>b</sup> Data expressed as mg g<sup>-1</sup> dry weight

<sup>c</sup> Data expressed as mg g<sup>-1</sup> dry weight

The leaf protein content significantly decreased under salinity stress, with the effect being more adverse as the osmotic potentials decreased more (Table no-3). In contrast, the content of soluble protein in leaf at PEG 10 % and 12% treatment increased over control by 21.45 and 6.29 % respectively. While at 18% PEG solution creating -0.8 MPa osmotic potential the content decreased by 48.63% as compared to unstressed control. However, the adverse effects of salinity and drought stress on protein in legumes were reported by Verma

et al. [24] and Bhardwaj et al. [28]. The leaf protein content registered more adverse effects of salinity stress in comparison with drought stress at iso-osmotic potentials.

The RLWC significantly decreased under all the treatments of drought and salinity stress as compared to that of control. The level of such decrease was higher as the osmotic potential of the growing medium decreased more. Such decrease in RLWC also indicated the reduction in leaf water potential under osmotic shock in all cases of stress in the present experiment (Table no-3). Thus, the corroborated well the early findings of Chen et al. [29], Bhardwaj et al. [28] and Petrovic et al. [30]. Thus, here salinity stresses of 50,100,200 mM were 73.98, 70.91 and 62.70 % and in case of drought stress imposed by PEG dose 10%,12%,18% the seedlings recorded RLWC of 75.28, 70.67 and 64.92 %, respectively. The salinity stress was found to produce more adverse effects on RLWC of ricebean in the present experiment.

The extent of leaf membrane damage was measured by determining the level of lipid peroxidation which in turn, was estimated as the content of thiobarbituric acid reactive substances (TBARS). The level of lipid peroxidation was significantly increased under all the treatments of drought and salinity stress (Table no-3). The level of such increase was higher as the osmotic potential of the growing medium decreased more. Earlier the adverse effects of salinity stress [31, 24] and drought stress [15] on lipid peroxidation in legumes were reported by different workers. This indicated greater damage of leaf membrane under stress leading to membrane leakiness. The drought stress exhibited more adverse effects than salinity stress at low and medium stress levels in terms of lipid peroxidation, although at the highest intensity of stress the salinity stress was found to be more damaging.

The content of total phenol in leaf significantly decreased under all the treatments of drought and salinity stress as compared to that of control. The level of such decrease was higher as the osmotic potential of the growing medium decreased more. Earlier the adverse effects of salinity stress [24] and drought stress [28] on phenol content in legumes were reported by different workers. The leaf phenol content registered lower means under drought stress in comparison with salinity stress at iso-osmotic potentials in the present experiment. The mean values under salinity dose of 50,100,200 mM NaCl were 4.80, 4.34, 3.09 mM GAE g<sup>-1</sup> fresh weight, respectively and in case of drought stress the mean values 3.86, 2.84, 2.41 mM GAE g<sup>-1</sup> fresh weight at PEG doses of 10%,12%,18% respectively (Table no-3). The drought was found to produce more adverse effects on phenol content than salinity stress in ricebean in the present experiment.

**Table 3. Effect of salinity and drought stress on soluble Protein, relative leaf water content (RLWC), lipid peroxidation and total phenol content in the leaves of ricebean cv. Bidhan 1**

Treatments	Protein <sup>a</sup>	RLWC(%)	Lipid peroxidation <sup>b</sup>	Phenol <sup>c</sup>
Control	164.36	87.62	94.38	5.73
50 mM NaCl	136.67	73.98	125.17	4.80
100 mM NaCl	99.19	70.91	128.45	4.34
200 mM NaCl	57.66	62.70	147.12	3.09
10% PEG	199.62	75.28	134.00	3.86
12% PEG	174.70	70.67	140.31	2.84
18% PEG	84.42	64.92	145.86	2.41
C.D. (P=0.05)	12.08	7.63	12.87	0.76

<sup>a</sup> Data expressed as mg g<sup>-1</sup> fresh weight

<sup>b</sup> Data expressed as µM of TBARS content g<sup>-1</sup> fresh weight

<sup>c</sup> Data expressed as mM of Gallic acid equivalent g<sup>-1</sup> fresh weight

#### 4. CONCLUSION

From the present study, it might be concluded that in general, the drought stress was found to register more drastic effects on seedling growth as compared to iso-osmotic potential of salinity stress, especially, at the highest intensity of stress in ricebean cultivar Bidhan 1. The drought stress also exhibited more negative effects on content of starch, total soluble sugar and phenol content of leaf as well as on membrane structure. While the photosynthetic pigment content in leaf RLWC and leaf protein was found to be more adversely affected by salinity stress.

#### REFERENCES

1. Epstein E. Mineral nutrition of plants: Principles and Perspectives. John Wiley and Sons, New York.1972.
2. Jaleel CU, Manivannan P, Wahid A, Farooq M, Al-juburi HJ, Somasundaram R, Panneerselvam R. Drought stress in plants: a review on morphological characteristics and pigments composition. *Int. J. Agric. Biol.* 2009;11:100–105.
3. Farooq M, Wahid A, Kobayashi N, Fujita, D. and Basra, SMA. Plant drought stress: effects, mechanisms and management. *Agron. Sustain. Dev.* 2009;29:185–212.
4. Kage H, Kochler M, Stutzel H. Root growth and dry matter partitioning of cauliflower under drought stress conditions: measurement and simulation. *European J. Agron.*2004;20:379–394.
5. Sosa L, Lianes A, Reinoso H, Reginato M, Luna V. Osmotic and Specific Ion Effects on the Germination of *Prosopisstrobilifera*. *Annals of Botany.*2005;96:261–267.
6. Arnon DI. Copper enzyme in isolated chloroplast polyphenol oxidase in *Beta vulgaris*. *Plant Physiol.*1949;24:1-15.
7. Heath RL, Packer L. Photoperoxidation in isolated chloroplast. 1. Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.*1968;12:189-198.
8. Perez NCM, Espinosa RG, Castaneda CL, Gallegos JAA, Simpson J. Water relation, histopathology and growth of common bean (*Phaseolus vulgaris* L.) during pathogenesis of *MacrophominaPhaseolina* under drought stress. *Physiol. Mol. Plant Pathol.*2002;60:185- 195.
9. McDonald S, Prenzler PD, Autolovich M, Robards K. Phenolic content and antioxidant activity of olive extracts. *Food Chem.*2001;73:73- 84.
10. Lowry OH, Rosebrogh NJ, Farr L, Randall RJ. Protein measurement with Folin phenol reagent. *J. Biol. Chem.*1951;193:265- 275.
11. Yoshida S, Forno DA, Cock JH, Gomez KA. Laboratory Manual for Physiological studies of Rice, 2<sup>nd</sup> edn. International Rice Research Institute, Los Banos, Philippines.1972.
12. Grieve CM, Suarez DL. Purslane (*PortulacaOleracea*L.): A Halophytic Crop for Drainage Water Reuse Systems. *Plant and Soil.*1997;192:277–283.
13. Jeannette S, Craig R, Lynch JP. Salinity Tolerance of *Phaseolus*Species During Germination and Early Seedling Growth. *Crop Science.*2002;42:1584-1594.
14. Gamze OKCU. Effects of Salt and Drought Stresses on Germination and Seedling Growth of Pea (*Pisumsativum*L.) *Turk J Agric.*2005;29:237-242.
15. Pratap V, Sharma YK. Impact of osmotic stress on seed germination and seedling growth in black gram (*Phaseolus mungo*).*Journal-of-Environmental Biology.*2010;31:721-726.

16. Bibi H, Sadaqat A, Tahir MHN, Akram HM. Screening of sorghum (*Sorghum bicolor* var *moench*) for drought tolerance at seedling stage in polyethylene glycol. *The Journal Animal Plant Science*.2012;22:671-678.
17. Khan MI, Shabbir G, Akram Z, Shah MKN, Ansar M, Cheema NM, Iqbal MS. Character association studies of seedling traits in different wheat genotypes under moisture stress conditions. *SABRAO Journal of Breeding and Genetics*.2013;45:458-467.
18. Anaytullah, Bose B, Yadav RS. PEG induced moisture stress: screening for drought tolerance in rice. *Indian J Plant Physiol*.2007;12(1):88-90.
19. El Sayed HEA. Influence of salinity (NaCl and Na<sub>2</sub>SO<sub>4</sub>) treatments on growth development of broad bean (*Vicia faba* L.) plant. *American Eurasian Journal of Agricultural and Environmental Sciences*.2011;10:600-610.
20. Aniat-ul-Haq, Vamil R, Agnihotri RK. Effect of Osmotic Stress (PEG) on Germination and Seedling Survival of Lentil (*Lens culinaris* MEDIK.). *Research Journal of Agricultural Sciences*.2010;1(3):201-204.
21. Mohammadkhani N, Heidari R. Drought-induced Accumulation of Soluble Sugars and Proline in Two Maize Varieties. *World Applied Sciences Journal*.2008;3:448-453.
22. Garg AK, Kim JK, Owens TG, Ranwala AP, Choi YD, Kochian LV, Wu RJ. Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses. *Proc Natl Acad Sci*.2002;99:15898-15903.
23. Munns R. Genes and salt tolerance: bringing them together. *New Phytologist*.2005;167:645-663.
24. Verma SK, Chaudhary M, Prakash V. Study of the alleviation of salinity effect due to enzymatic and non-enzymatic antioxidants in *Glycine max*. *Research-Journal-of-Pharmaceutical,-Biological-and-Chemical-Sciences*.2012;3:1177-1185.
25. Macar TK, Turan O, Ekmekci Y. Effects of Water Deficit Induced by PEG and NaCl on NaCl on Chickpea (*Cicer arietinum* L.) Cultivars and Lines at Early Seedling Stages. *G.U. Journal of Science*.2009;22:5-14.
26. Zhang HX, Irving LJ, McGill C, Matthew C, DaoWei Z, Kemp P. The effects of salinity and osmotic stress on barley germination rate sodium as an osmotic regulator. *Annals of Botany*.2010;106:1027-1035.
27. Adele M, Maria S, Umberto A, Carmelo S, Albino M. Effect of PEG-induced drought stress on seed germination of four lentil genotypes. *Journal of Plant Interactions*.2014;9(1):354-363.
28. Bhardwaj J, Yadav SK. Comparative study on biochemical parameters and antioxidant enzymes in a drought tolerant and a sensitive variety of horsegram (*Macrotyloma uniflorum*) under drought stress. *American Journal of Plant Physiology*.2012;7:17-29.
29. Chen CS, Xie ZX, Liu XJ. Dynamic transformation of the substances of osmotic adjustment in winter wheat under iso-osmotic salt and drought stresses. *Bulletin of Botanical Research*.2009;29(6):708-713.
30. Petrović G, Jovičić D, Nikolić Z, Tamindžić G, Ignjatov M, Milošević D, Milošević B. Comparative study of drought and salt stress effects on germination and seedling growth of pea. *Genetika*.2016;48:373 -381.
31. Dar ZM, Hemantaranjan A, Panday SK. Antioxidative response of mungbean (*Vigna radiata* L.) to salt stress. *Legume-Research*.2007;30:57-60.