

**Antifungal activity and phytochemical screening of *Cymbopogon citratus*, *Cajanus cajan*, and *Plectranthus amboinicus* leaves collected in Guyana, South America**

---

**ABSTRACT**

**Aims:** Medicinal plants have been a fundamental part of the human health since existence. Guyana is surrounded high in the green shoulder of northern South America and shares Amazon River and Amazon Forest. South American population use plant extracts obtained from traditional medicinal plants as medicine for many infectious diseases. The aim of the study was to evaluate antifungal property and chemical composition of the three medicinal plants *Cymbopogon citratus* (lemon grass), *Cajanus cajan* (pigeon pea), and *Plectranthus amboinicus* (thick leaf thyme) leaves collected from the coastal areas of Guyana.

**Study design:** Experiment based study.

**Place and Duration of Study:** Plants were gathered along the East Coast of Guyana and identified at the Biodiversity Center, University of Guyana, Georgetown, Guyana between January 2017- May 2017.

**Methodology:** Phytochemical extraction was conducted using soxhlet and rotovap apparatus and an aqueous extraction method. Data analysis of the study was done using R-Studio Program for statistical computing and graphics. A Tukey test was done along with ANOVA and Boxplots.

**Results:** Qualitative analysis of phytochemicals were carried out and the presence of terpenoids, steroids, glycoside, alkaloid, tannins and saponins were positive in some plants. Antifungal activity was tested using the poisoned food and well diffusion techniques.

**Conclusion:** In conclusion, *C. cajan* showed significant zones of inhibition using well diffusion technique whereas hexane extract showed significant inhibition with poisoned food technique.

*Keywords: antifungal, plant extract, inhibition, phytochemical, poison food technique, well diffusion technique*

**1. INTRODUCTION**

Nature always has wide variety of therapeutically valued bioactive compounds to offer mankind. Plant product are considered safe for their pharmacological activities and widely recognized for their vast diversity [1] [2]. Human beings are dependent on medicinal plants ever since existence. Past two decades showed an increasing interest in the investigation of different medicinal plant extracts [3]. World Health Organization (WHO) estimates that about 80% of people are still dependent on traditional herb-based medications globally due to their low cost, easy accessibility and likely negligible side effects in comparison to allopathic medicines [4] [5]. Before the advent of synthetic fungicides, plant derivatives were commonly used for fungal control in developing countries. Plant derivatives as fungicides were relatively cheap compared

to imported synthetic fungicides [6]. Damage to crops by fungal pathogens has pressured farmers to use of antifungal control agents [7]. Several of the synthetic fungicides are reported to cause adverse effects on treated soil ecosystems because of their non-biodegradable nature [8]. Substantial use of chemical pesticides induces health problems and environmental hazards in agricultural system, therefore it is of no doubt that natural products of antimicrobial activity are best bio rational alternatives today [9].

*C. citratus*, commonly called as lemon grass is a tall perennial grass which grows in tropical and subtropical habitats [10]. *C. citratus* is well known to have antidepressant, antioxidant, antiseptic, sedative, nervine, bactericidal, and fungicidal properties [11] [12]. It grows well in tropical and subtropical regions of Asia, South America, and Africa [13]. *C. cajan* is known to grow well in tropical and subtropical regions with wide medicinal use [14]. *C. cajan* is an annual perennial shrub typical to dry climates and few parts of South America [15]. Next plants of genus *Plectranthus* included in this study has over 3000 recognized species, spread along countries in Africa, South America, Asia and Australia [16] [17] [18]. The aim of the present study was to identify phytochemicals and antifungal activity of *C. cajan*, *C. citratus* and *P. amboinicus* leaf extracts.

## **2. MATERIAL AND METHODS**

### **2.1 PLANT COLLECTION**

Plants were gathered along the East Coast of Guyana and identified at the Biodiversity Center, University of Guyana, Georgetown, Guyana. The collected leaves of each plant were washed under running tap water, dried and placed in brown paper bags. The paper bags were weighed and placed in a hot air oven, drying at a constant temperature of 55°C until a constant weight was recorded. The dried leaves were powdered using a mixer grinder at GuySuCo Laboratory and stored for the next step [19].

### **2.2 SOXHLET APPARATUS**

Adhering accordingly to the method followed at the Pesticides and Toxic Chemicals Control Board (PTCCB), 64 grams of dried plant material were weighed and placed in a thimble. The thimble (a thick porous cellulose container) was placed into the extraction chamber. The selected solvent was slowly poured through the condenser opening. The boiling flask was heated by a heating mantle. The boiling flask collected the extracted phytochemicals with each evaporation that passed through the siphon arm and the solvent vapor was rapidly cooled in the condenser's cooler. Each plant used a new thimble and fresh solvent.

### **2.3 ROTARY EVAPORATOR (ROTOVAP)**

The rotary evaporator reduces the solutions down to a solid state. The extract containing either hexane or methanol as a solvent were collected in the boiling flasks from the soxhlet apparatus. Each flask with residue were labelled; depicting solvent used and plant species.

### **2.4 AQUEOUS EXTRACTION**

The aqueous extraction was carried out using the standard method [20]. About 15 grams of grounded leaves from each plant species were extracted by successive soaking for 3 days using 35 ml of distilled water in separate container. The extracts were filtered using Whatman No. 1 filter paper using a vacuum. The filtrates were concentrated by evaporation at low

temperature of 30°C using a water bath. The concentrated samples were used to make a 50% stock solution from which the tested concentrations were created. Phytochemical residue of 5 grams was added to 5 ml of the indicated solvent to make 50% stock solution. From the stock solution, 300µl, 400µl and 500µl were used to check the antifungal property. Each test was done in triplicate per plant and solvent.

## 2.5 QUALITATIVE ANALYSIS OF PHYTOCHEMICALS

Phytochemical analysis was done to identify the presence of the phytochemicals; tannins, alkaloids, glycoside, saponins, flavonoids, terpenoids and steroids [21].

## 2.6 CULTURE TECHNIQUE

Antifungal activity was observed on *Aspergillus niger*, the strain was obtained from pure cultures at the University of Guyana, Berbice Campus. *A. niger* was cultured and maintained on Potato Dextrose Agar (PDA). Antifungal activity was performed by Well Diffusion and Poisoned Food technique. Measurements for Poisoned Food Technique was done in accordance to [13] between five to seven days or once the control was completely covered.

Percentage of mycelial growth inhibition was calculated from the formula

$$\text{Mycelial growth inhibition} = \frac{\text{diameter of control} - \text{diameter of sample}}{\text{diameter of control}} * 100$$

## 2.7 ANALYSIS

Data analysis of the study was done using R-Studio Program for statistical computing and graphics. A Tukey test was tested along with ANOVA and Boxplots were constructed. A Tukey test was used to compare concentration, plant and solvents versus techniques. Box and whiskers plot also called boxplots were used as visual representations of the replicates.

## 3. RESULTS AND DISCUSSION

### 3.1 PHYTOCHEMICAL ANALYSIS

Table 1 shows the result of phytochemical screening in different plant extracts. Hexane, a solvent with a low polarity, extracted the most phytochemicals with each plant, while water, a solvent with high polarity, extracted the least. Alkaloid was only present in the water extraction of *C. cajan*, while Terpenoid was present only in *C. citratus*. Glycosides contents were extracted by hexane leaf extract of *C. cajan* and *P. amboinicus* only and steroid contents were extracted by methanol extracts of all the three plant leaves, saponins was the most common phytochemical identified by the plant extracts.

**Table 1 Results on the qualitative tests indicating presence or absence of phytochemicals in each plant solvent.**

Phytochemical	<i>P. amboinicus</i>			<i>C. citratus</i>			<i>C. cajan</i>		
	Hex	Meth	Water	Hex	Meth	Water	Hex	Meth	Water

Saponins	+	+	+	+	+	+	+	+	-
Tannins	-	-	-	-	+	-	-	-	+
Flavonoid	+	-	-	+	-	-	-	-	+
Alkaloid	-	-	-	-	-	-	-	-	+
Terpenoid	-	-	-	+	-	+	-	-	-
Steroid	+	+	-	+	+	-	-	+	-
Glycoside	+	-	-	-	-	-	+	-	-

\**C. citratus* extracted with methanol displayed a green black color that indicated catecholic tannins.

\**C. cajan* extracted with water displayed a blue color that indicated gallic tannins.

\**C. cajan* extracted with hexane displayed an orange color that indicated flavones

### 3.2 POISONED FOOD TECHNIQUE


Table 2 illustrates mean±SE for every plant extract. Tukey test found a significant statistical difference between inhibition percentage and plant extract (Figure 1). *C. cajan* showed a significant inhibition compared to other two plants. Hexane extract of *P. amboinicus* and *C. cajan* showed statistical significance inhibition at 300µl concentration were as hexane extract of *P. amboinicus* and *C. citratus* showed inhibition at 500µl concentration (Figure 2). The Tukey test also showed a statistical significant difference between inhibition percentage and solvents between hexane and the other two solvents -methanol and water (Figure 3). Table 3 shows the inhibition percentage between the tree leaf extract.

**Table 2 Measurements of fungal growth millimeters (mm) and standard error for the Poisoned Food Technique (mean±SE).**


Conc (µl)	<i>P. amboinicus</i>			<i>C. citratus</i>			<i>C. cajan</i>		
	Hex	Meth	Water	Meth	Meth.	Water	Hex	Meth	Water
300	80±5.8	45±5.0	90±0.0	47±1.5	90±0.0	90±0.0	25±4.4	60±7.3	47±2.7
400	70±1.7	50±2.1	90±0.0	40±2.6	90±0.0	90±0.0	30±2.1	60±5.0	90±0.0
500	52±3.1	55±4.4	90±0.0	20±7.6	90±0.0	90±0.0	30±3.3	50±7.3	36±6.4

**Table 3 Mean measurements of zone of inhibition in millimeters (mm) and the standard errors for the Well Diffusion Technique.**

Conc (µl)	<i>P. amboinicus</i>			<i>C. citratus</i>			<i>C. cajan</i>		
	Hex	Meth	Water	Hex	Meth	Water	Hex	Meth	Water
300	6±1.9	5±0.3	0	10±0.3	1±0.6	1±0.0	3±0.3	11±0.9	20±1.7
400	16±2.2	7±0.3	0	18±0.9	7±0.3	2±0.3	5±0.3	15±1.7	25±1.7
500	23±1.8	9±0.9	0	21±1.8	9±0.3	4±0.3	7±0.9	18±3.1	30±4.4



**Figure 1. Boxplot (mean±SD) for different plants and percent inhibition.**



**Figure 2 Boxplot (mean±SD) for different solvents and percent inhibition.**






Figure 3 Boxplot (mean±SD) between concentrations and percentage inhibition.

### 3.3 WELL DIFFUSION TECHNIQUE


The size of the inhibition zones was measured 48 hours after plating. The mean zone of inhibition of the three replicated tests of the plant extract were expressed in millimeter. According to the Tukey test there is statistically significant difference between *C. cajan* and the other two plants. Water extract of *C. cajan* was effective compared to water extracts of *P. amboinicus* and *C. citratus*. There is a significant difference between *C. cajan* and the other two plants. *C. cajan* showed highest zone of inhibition was thirty-five while its lowest was three (Figure 4). Hexane on average produced significant zones of inhibition with all plants but water worked well with *C. cajan*, producing the highest zones of inhibition (Figure 5). The rising gradient of *C. cajan* illustrates effectiveness inhibition with based on various concentration of each plant (Figure 6).



**Figure 4** Boxplot (mean±SD) for different plants and the zones of inhibition (Zoi).



**Figure 5** Boxplot (mean±SD) for different solvents and the zones of inhibitions



**Figure 6** Boxplot (mean±SD) for different concentrations

## 4 DISCUSSION

Various studies in recent years have demonstrated the importance of phytochemicals due to their outstanding health benefits. Plant metabolites are extracted by using various ways and efficiency of extraction method depend on several factors like the nature of phytochemical constituents, the method of extraction, particle size of the sample, extraction time, temperature, pH, solute to solvent ratio, and the solvent polarity [22]. It is very important to choose appropriate solvent system to recover higher extract yield and bioactive compounds from a sample [23]. This study demonstrated a significant difference in the extract yield obtained with different solvents. Literatures have highlighted methanol as the best solvent for extraction [24, 23, 5]. Methanol extract of lemon grass showed greater antimicrobial properties with increasing concentration [25]. Similarly, methanol extract of *Argemone mexicana* leaves and seeds showed greater antibacterial activity than water extracts [26].

Preparation of an extract with an organic solvent was shown to have greater antibacterial activity [27]. GC-MS method showed 11 major peaks in the quantitative phytochemical analysis of methanol and ethanol of *P. amboinicus*, as well as excellent antimicrobial properties [28]. Study has reported that essential oil of *P. amboinicus* possess antifungal activity against *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus ochraceus*, *Aspergillus oryzae*, *Candida versatilis*, *Fusarium moniliforme*, and *Saccharomyces cerevisiae* [29]. *C. citratus* showed antibacterial properties in two of the three main components of the oils identified i.e. the  $\alpha$ -citral (geranial) and  $\beta$ -citral (neral) components [30]. Study has shown antibacterial effect of *C. citratus* with methanol and aqueous extracts at 20  $\mu$ gm [31]. The qualitative phytochemical tests of *C. citratus* showed only the presence of tannins and steroids, indicating extremely low percentage alkaloids, flavonoids and saponins. Ethyl acetate extract of *C. citratus* showed highest antimicrobial activity with *Aspergillus sp* and *Mucor sp* than with aqueous and methanol extracts [32].

The aqueous extraction method in this study found effective antibacterial property but poor antifungal property with *C. cajan* [33]. However, it was stated that *C. cajan* is more active against fungus and that *A. niger* showed 48 components in the GC-MS spectrum [34]. Various studies in the past have demonstrated a major component in pigeon pea leaves that have potential benefits to human health with flavonoids and stilbene [35 36] including treatment of diabetes and jaundice [37, 38, 39].

## 5. CONCLUSION

Some form of inhibition was shown by all three plants and the solvent hexane of *C. cajan* were most successful. The well diffusion technique showed efficient zones of inhibition with *C. cajan*'s water extract. Hexane showed significant inhibition with the Poisoned Food technique especially with *C. citratus* and *C. cajan*.

## REFERENCES

1. Newman DJ, Cragg GM. Natural products as sources of new drugs from 1981 to 2014. *J Nat Prod*. 2014; 79(3): 629-661
2. Thenmozhi K, Anusuya N, Ali MA, Jamuna S, Karthika K, Venkatachalapathi A et al. Pharmacological credence of the folklore use of *Bauhinia malabarica* in the management of jaundice. *Saudi J Biol Sci*. 2018 (25);22-26
3. Bonjar GHS, Farrokhi PR. Anti-bacillus activity of some plants used in traditional medicine of Iran. *Nig J Nat Prod Med*. 2005; 8:34-39



4. Swamy MK, Sinniah UR. A comprehensive review on the phytochemical constituents and pharmacological activities of *Pogostemon cablin Benth*: An aromatic medicinal plant of industrial importance. *Molecules*. 2015a; 20: 8521–8547
5. Swamy MK, Sinniah UR, Akhtar MS. In vitro pharmacological activities and GC-MS analysis of different solvent extracts of *Lantana camara* leaves collected from tropical region of Malaysia. *Evid Based Complementary Altern Med*. 2015: 1–9
6. Galvano F, Piva F, Riteni A, Galvano G. Dietary strategies to counteract the effects of mycotoxins: A review. *J Food Prot*. 2001; 64: 120-131
7. El-Khateeb AY, Elsherbiny EA, Tadros LK, Ali SM, Hamed HM. Phytochemical analysis and antifungal activity of fruit leaves extracts on the mycelial growth of fungal plant pathogens. *J Plant Pathol Microb*. 2013; 4: 4-9
8. Tegegnea G, Pretoriusa JC, Swart WJ. Antifungal properties of *Agapanthus africanus* L. extracts against plant pathogens. *Crop Prot*. 2008; 27: 1052-1060
9. Tiwari RKS, Singh A, Das K, Sinha A. Efficacy of extracts of medicinal plants against *Rhizoctonia solani*. *Ann Plant Prot Sci*. 2007; 15: 460-539
10. Cheel J, Theoduloz C, Rodríguez J, Schmeda-Hirschmann G. Free radical scavengers and antioxidants from Lemongrass (*Cymbopogon citratus* (DC.) Stapf.) *J Agric Food Chem*. 2005; 53: 2511–2517
11. Friedman M, Henika PR, Mandrell RE. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes* and *Salmonella enterica* *J Food Prot*. 2002; 65: 1545–1560
12. Gardner ZE, McGuffin M. American Herbal Products Association's botanical safety handbook. 2nd ed. Boca Raton: CRC Press, American Herbal Products Association; 2013
13. Akhila A. Essential oil-bearing grasses: the genus *Cymbopogon*. New York: CRC Press; 2010
14. Agus C, Achmadi SS, Mubarik NR. Antibacterial activity of naringenin-rich fraction of pigeon pea leaves toward *Salmonella typhi*. *Asian Pac J Trop Biomed*. 2017; 7: 725-728
15. Odeny DA, Jayashree B, Fergusow M, Hoisington D, Cry LG, Gebhardt C. Development, characterization and utilization of microsatellite markers in Pigeon pea. *Plant Breed*. 2007; 126: 30–36
16. Waldia S, Joshi BC, Pathak U, Joshi MC. The genus *Plectranthus* in India and its chemistry. *Chem Biodivers*. 2011; 8: 244–52
17. Daglia M. Polyphenols as antimicrobial agents. *Curr Opin Biotechnol*. 2012; 23: 174–81
18. Swamy MK, Arumugam G, Kaur R, Ghasemzadeh A, Yusoff MM, Sinniah UR. GC-MS based metabolite profiling, antioxidant and antimicrobial properties of different solvent extracts of Malaysian *Plectranthus amboinicus* leaves. *Evid Based Complement Alternat Med*. 2017; 1517683: 10
19. Ashidi J.S, Houghton PJ, Hylands PJ, Efferth T. Ethnobotanical survey and cytotoxicity testing of plants of South-western Nigeria used to treat cancer, with isolation of cytotoxic constituents from *Cajanus cajan* Millsp. leaves. *J Ethnopharmacol*. 2010; 128: 501-512
20. Asuzu IU. Pharmacological evaluation of folklore of *Sphenostylia stenocarpa*. *J. Ethnopharmacol*. 1986; 16: 236-267
21. Siddiqui AA, Ali M. Practical Pharmaceutical chemistry, 1st edition. CBS publishers and distributors New Delhi. 126-131.
22. Do QD, Angkawijaya AE, Tran-Nguyen PR. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila* aromatic *J. Food Drug Anal*. 2014; 22: 296–302
23. Iloki-Assanga SB, Lewis-Luján LM, Lara-Espinoza CL, Gil-Salindo AA, Fernandez-Angulo A, Rubio-Pino, Haines DD. Solvent effects on phytochemical constituent profiles

- and antioxidant activities, using four different extraction formulations for analysis of *Bucida buceras* L. and *Phoradendron californicum*. BMC Res Notes. 2015; 8: 396
24. Sultana B, Anwar F, Ashraf M. Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. Molecules. 2009; 14: 2167–2180
  25. Jafari B, Ebadi A, Aghdam BM, Hassanzade Z. Antibacterial activities of lemon grass methanol extract and essence on pathogenic bacteria. American- Eurasian J Agric Environ Sci. 2012; 12: 1042-1046
  26. Bhattacharjee I, Chatterjee SK, Chatterjee S, Chandra G. Antibacterial potentiality of *Argemone mexicana* solvent extracts against some pathogenic bacteria. Mem Inst Oswaldo Cruz. 2006; 101: 645-648
  27. Nair R, Kalariya T, Sumitra C. Antibacterial activity of some selected Indian medicinal flora. Turk J Biol. 2005; 29: 41-47
  28. Manjamalai A, Singh RSS, Guruvayoorappan C, Grace VMB. Analysis of phytochemical constituents and antimicrobial activity of some medicinal plants in Tamil Nadu, India. Global J Biotechnol Biochem. 2010; 5: 120-128
  29. Murthy PS, Ramalakshmi K, Srinivas P. Fungitoxic activity of Indian borage (*Plectranthus amboinicus*) volatiles. Food Chem. 2009; 114: 1014-8
  30. Onawunmi GO, Yisak WA, Ogunlana EO. Antibacterial constituents in the essential oil of *Cymbopogon citratus* (DC.) Stapf. J Ethnopharmacol. 1984; 12: 279-286
  31. Ojo OO, Anibijuwon II, Ojo OO. Studies on extracts of three medicinal plants of South Western Nigeria: *Hoslundia opposita*, *Lantana camara* and *Cymbopogon citratus*. Adv Nat Appl Sci. 2010; 4: 93-98
  32. Bhavya T and Padma S. Antifungal activity of *Cymbopogon citratus* leaves extract against herbal drug contaminants. World J Pharm Pharmaceut Sci. 2014; 3(6): 768-775
  33. Ezeifeke GO, Orji MU, Mbata TI, Patrick AO. Antimicrobial activities of *Cajanus cajan*, *Garcinia Kola* and *Xylopiya aethiopica* on pathogenic microorganisms. Biotech. 2004; 3: 41-43.
  34. Abdel KM, Zohor M, Inas O, EL- Hafez M. GC-MS Analysis and biological activity of fixed oil from Sudanese *Cajanus cajan* (leguminosae) seeds. Int J Adv Res. 2016; 4: 1373-1380
  35. Fu YJ, Liu W, Zu YG, Tong MH, Li SM, Yan MM, Efferth T, Luo H. Enzyme assisted extraction of luteolin and apigenin from pigeon pea [*Cajanuscajan* (L.) Millsp.] leaves. Food Chem. 2008; 1111: 508-512
  36. Luo QF, Sun L, Si JY, Chen DH. Hypocholesterolemic effect of stilbenes containing extract-fraction from *Cajanus cajan* L. on diet-induced hypercholesterolemia in mice. Phytomedicine. 2008; 15: 932-939
  37. Nahar L, Nasrin F, Zahan R, Haque A, Haque E, Mosaddik A. Comparative study of antidiabetic activity of *Cajanus cajan* and *Tamarindus indica* in alloxan-induced diabetic mice with a reference to in vitro antioxidant activity. Pharmacog Res. 2014; 6: 180-7
  38. Chander MP, Kartick C, Vijayachari P. Ethnomedicinal knowledge among Karens of Andaman & Nicobar Islands, India. J Ethnopharm. 2015; 162: 127-133
  39. Uchegbu NN, Ishiwu CN. Germinated Pigeon Pea (*Cajanus cajan*): a novel diet for lowering oxidative stress and hyperglycemia. Food Sci Nutri. 2016; 4: 772-777