

**EFFECT OF CRUDE OIL CONCENTRATIONS ON THE VASCULAR BUNDLE SIZE
OF TWO ZEA MAYS VARIETIES**

Abstract

The reliance of humanity on fossil fuels particularly petroleum hydrocarbons have been on the increase. This had led to the increase pollution of agricultural lands. This study was aimed at investigating the effect of crude oil concentrations on the vascular bundles size of two *Zea mays* varieties – Oba Super Zaria and SMZ 37 Kaduna commonly cultivated in the South-eastern Nigeria. The experiment study was conducted in the green house in the Department of Biological Sciences, faculty of Natural Sciences, Kogi State University, Ayingba, Kogi State. The experiment was setup as a 2×4 factorial laid out in a Completely Randomized Design replicated three times with four treatment concentrations (0-control, 15, 30 and 45ml) of petroleum crude used and two maize varieties used. Significant effects ($P < 0.001$) of the concentrations as well as the varieties were observed on the vascular bundles. The result showed a significant ($P < 0.05$) decrease in vascular bundle with increasing concentration was observed with 45ml concentration having the smallest vascular bundles. The vascular bundles of Oba super Zaria decreased significantly with increasing concentrations of crude oil ($p \leq 0.05$), while significant differences was observed 45 ml treatment in SMZ 37 Kaduna as compared to the control. Based on the results obtained from this investigation, OSZM appeared to be more susceptible to soils affected with crude oil.

Keywords: Crude Oil ; *Zea Mays*; Pollution; Soil; Vascular Bundle

Introduction

Maize is ranked as the third most essential cereal crop following wheat and rice in the world production of cereal crops (FAO, 2002). Maize is one of the main staple consumed in Nigeria, as it makes available an inexpensive nutritious food that is of great assistance in sustaining the rapid increasing population. It also serves as raw material in many industrial processes and in the feeding of farm animals (Olaniyan and Lucas, 2004). Maize is grown in most agro ecological

30 areas especially in the Niger Delta region where oil industrial activities are predominant
31 (Agbogidi, *et al.*, 2007). The growth and yield of crops are adversely affected by crude oil
32 pollution, and this has been reported by many workers to include poor and stunted growth, low
33 production and sometimes outright death of crops (Anoliefo, 1991; Anoliefo and Vwioko, 1994;
34 Agbogidi, *et al.*, 2007).

35 The reliance of humanity on fossil fuels particularly petroleum hydrocarbons have been on the
36 increase. This had led to the increase pollution of agricultural lands, as a result of spillage of
37 crude oil during the process of extraction and processing (Ayotamuno and Kogbara, 2007). In
38 Nigeria, reports have it that from 1976 to 1998, a sum of about 5,724 incidences of oil spills
39 which introduced crude oil to both the terrestrial and aquatic ecosystems to the tune of
40 2,571,113.90 barrels (Aroh *et al.*, 2010). Since then, crude oil contamination of farmlands has
41 become common experience in the country, and many of these farmlands have been abandoned
42 in the aftermath of pollution.

43 The presence of crude oil in the soil renders it inadequate for plant growth, because of the
44 reduced level of available plant nutrients as well as the rising toxic levels of certain micro
45 elements (Barua *et al.*, 2011). The susceptibility of plants to crude exposure is high, and plants
46 may end up dying in a matter of few weeks or months. Some plant species are able to grow in
47 hydrocarbon polluted soils and bring about their degradation via the rhizosphere part of their
48 root. This is because the rhizosphere supports the growth of many microorganisms which
49 increases biomass and microbial activity, thereby accelerating degradation process (Quinones-
50 Aquilar *et al.*, 2003; Agbogidi, *et al.*, 2005).

51 The essential elements of vascular systems are the xylem, concerned with the transport of water
52 and dissolved salts, and the phloem, which translocates synthesized but soluble materials around
53 the plant to places of active growth or regions of use or storage (Cutler *et al.*, 2007). Xylem and
54 phloem are normally associated and together form the vascular bundles which are often enclosed
55 in a sheath of fibres. In addition, contains an outer sheath of parenchyma cells (the bundle
56 sheaths) in some instances. Vascular bundles make up the 'plumbing system' of primary tissues,
57 and organs without secondary growth in thickness (Cutler *et al.*, 2007). This study was carried
58 out to ascertain the growth potentials of a commonly grown crop in the country, *Zea mays*
59 (maize), on a crude oil contaminated soil.

60 **MATERIALS AND METHODS**

61 The soil used in this study was obtained from the top soil collected from a farm land. The soil
62 was treated by sieving and autoclaving at 82⁰C for 30 minutes to remove roots that may serve as
63 an alternative host to pathogens that may affect the growth of the plant. Two varieties of maize
64 used for the study were the Oba Super Zaria maize, and SMZ 37 Kaduna. Unrefined crude oil
65 was gotten from the Nigeria National Petroleum Corporation, (NNPC). The crude oil was mixed
66 with water to vary the concentration: 15ml, 30ml, and 45ml of the crude oil were mixed in 500ml
67 of distilled water.

68 Five hundred and fifty gram of treated top soil was weighed and put into 24 pots and each pot
69 was planted with three seed of maize. The planting pots were perforated at the bottom and sides
70 to allow for aeration and drainage of excess water. After germination, the pots (soil) were all
71 polluted with crude oil at different concentration (15, 30 and 45ml), except the control, at two
72 weeks after germination. The experiment was observed for two week with continuous watery.
73 The experiment was setup as a 2 × 4 factorial laid out in a Completely Randomized Design
74 replicated three times. The experiment study was conducted in the green house in the
75 Department of Biological Sciences, faculty of Natural Sciences, Kogi State University, Ayingba,
76 Kogi State.

77 The stems were harvested and sections were made using a Reichert sledge microtome. The
78 sections were stained, using the counter staining procedures of Jane 1962, as adopted by
79 Ajuziogu *et.al.* (2018). The counter staining procedures stained the lignified tissues of the
80 vascular bundles red, and contrasted them from the unlignified tissues of the ground meristem.
81 Fifteen measurement of vascular bundle diameter were made for each of the various treatment
82 samples and the control at × 100 magnifications using a calibrated microscope.

83 Data obtained were subjected to a two-way analysis of variance via a univariate analysis using
84 IBM statistical package for social sciences (SPSS) version 20. Significant means were separated
85 using least significant difference test generated from GenStat Discovery Edition 4 statistical
86 package.

87

88 **RESULT**

89 The analysis of variance on the effect of crude oil concentration and the varietal effect on the
90 vascular bundle size was highly significant ($P < 0.001$) across the effect of the varieties,


91 concentration and the combined effect of maize variety interacting with crude oil concentration
92 (Table 1).

93 **Table 1: Analysis of Variance showing the effect of crude oil concentration on two varieties**
94 **of maize**


Source	Sum of Squares	df	Mean Square	F
Variety	0.09	1	0.09	139.67***
Concentration	0.18	3	0.06	93.06***
Variety × Concentration	0.13	3	0.04	67.69***
Error	0.01	16	0.001	
Total	0.40	23		

95 ***- Significant at $P < 0.001$

96 The mean differences of the vascular bundle across the varieties as present on Fig. 1 shows that
97 Oba Super Zaria Maize (OSZM) has significantly ($P < 0.05$) wider vessels as compared to SMZ
98 37 Kaduna. As recorded on Fig 2, there was a crude oil dose dependent effect on the vascular
99 bundle. As compared to the control which had an average vascular bundle of $0.72 \pm 0.08\text{mm}$, a
100 significant ($P < 0.05$) decrease in vascular bundle with increasing concentration was observed
101 with 45ml concentration having the smallest vessels (Fig 2).



103 **Fig 1: Mean vascular bundleof two maize variety exposed to varied concentration of crude**
 104 **oil**



105
 106 **Fig 2: Mean crude oil concentration effect on the vascular bundleof two maize variety**
 107 At 0 ml of crude oil, Oba Super Zaria maize, (OSZM) had a mean diameter of 0.91 ± 0.00 mm
 108 which was significantly higher ($p < 0.05$) than every other treatment combinations. This was
 109 followed by the vascular bundleof the same variety exposed to 15 ml of crude oil with an
 110 average diameter of 0.61 ± 0.01 mm which was also significantly wider as compared to higher
 111 concentrations (Table 2).
 112 The smallest vascular bundleof 0.49 ± 0.03 mm was observed from SMZ 37 Kaduna variety
 113 exposed to 45ml concentration of crude oil. This was however not significantly different from
 114 the vascular bundleobserved from OSZM variety treated with the same concentration (45ml) and
 115 SMZ 37 Kaduna variety exposed to 30ml (Table 2).

116
 117 **Table 2: Interactive ffect of crude oil concentration on the vascular bundleof two maize**
 118 **varieties**

Concentration	Suprzaria Maize	SMZ 37 Kaduna
0 ml	0.91 ± 0.00^a	0.54 ± 0.00^{cd}
15 ml	0.61 ± 0.01^b	0.54 ± 0.00^{cd}
30 ml	0.55 ± 0.01^c	0.52 ± 0.01^{cde}
45 ml	0.50 ± 0.02^{ce}	0.49 ± 0.03^e

119 *Means with different alphabets on the table are significantly different using Least Significant Difference (LSD) at
 120 $P < 0.05$

121 **DISCUSSION**

122 The study showed that an increase in crude oil concentration in the soil decrease the diameter of
123 the vascular bundle in maize. This might possibly be attributed to the differences in polarity
124 between water and crude oil. Water is a high polar solvent while hydrocarbon oil (crude oil) is a
125 non polar solvent, therefore a mixture of water and hydrocarbon oil will result in formation of
126 two layers, with the hydrocarbon oil layer above and the water below. However, pollution of soil
127 with this mixture of water and hydrocarbon oil would possibly result to the masking or shielding
128 of water molecules and other minerals in the soil by the hydrocarbon oil. As a result of this, there
129 would be no or less uptake of water and other nutrient contained in the soil by plant. These in
130 turn would lead to shrinking of the vascular bundles and retardation of plants growth within
131 polluted environment due to impose draught by hydrocarbon oil. Increasing the concentration of
132 crude oil led to a proportional increase in the rate of shrinking of the vascular bundles and hence,
133 reduction in the size of the plant. This reduction of vascular bundle will also lead to slow growth
134 and low yield of grains. These negative effects of crude oil pollution on crops had been earlier
135 highlighted by Odu (1983), Bello *et al.* (1999), Ekundayo *et al.* (2001) and Dung *et al.* (2008).
136 Therefore, this study supported the findings of the above named authors and confirmed that the
137 results were similar from the point of view of output reduction on crude oil pollution. Similarly
138 Ubochi *et al.* (2019) had reported decrease in nutrients uptake by plants as a result of the oil
139 content in coconut oil effluent.

140 In view of the dose-dependent decrease in vascular bundleof maize, Anoliefo and Edegbai
141 (2001) reported that low level of oil in pollution could be easily be degraded by natural
142 rehabilitation in soils, increase organic matter in soil and improve the fertility, physical and
143 chemical properties of the soil. The differences in the size of vascular bundles of maize varieties
144 in response to crude oil levels corroborates the reports of Baker (1970) and Naegele (1974) that
145 the effect of crude oil on plants is dependent on the variety amongst other factors. The
146 differences in plants reaction to pollution are due to an innate genetic response of the plant
147 system as modified by environmental influences (Baker, 1970; Naegele, 1974). Based on the
148 results obtained from this investigation, OSZM appeared to be susceptible to soils affected with
149 crude oil.

150

151

152 **REFERENCES**

- 153 Agbogidi, O.M., Eruotor, P.G., and Akparobi, S.O. (2005). Effects of location and crude oil
154 levels on the growth of seven maize varieties (*Zea mays* L.). In: Uguru MI, Iroegbu CU,
155 Ejere VC (eds.). *Proceedings of the 30th Annual Conference of the Genetics Society of*
156 *Nigeria*, held at the University of Nigeria, Nsukka, Enugu State, between 5th and 8th of
157 September. pp. 95-101.
- 158 Agbogidi, O. M., Eruotor, P.G. and Akparobi, S.O. (2007). Effects of time of application of
159 crude oil to soil on the growth of maize (*Zea mays* L.). *Research Journal of*
160 *Environmental Toxicology*, 1 (3), 116 – 123.
- 161 Ajuziogu, G.C, Onyeke, C.C. Ayogu, V.O., Asuzu, C.U., Urama, D. C. Odoh, N.V. (2018).
162 Treatability indices of some plant species of Fabaceae in Nigeria. *Wood Research*, 63 (6)
163 971-978.
- 164 Anoliefo, G. O. (1991). *Forcados blend crude oil effects on respiratory mechanism, mineral*
165 *element composition and growth of Citrullus vulgaris* Schead. Unpublished Ph.D Thesis
166 University of Benin, Benin-City. p. 293.
- 167 Anoliefo, G.O. and Edegbai, B.O. (2001). Effect of crude oil as a soil contaminant on the growth
168 of two egg plant species, *Solanum melongena* L. and *S. incanum*. *Journal of Agriculture,*
169 *Forestry and Fisheries*, 1: 1-25.
- 170 Anoliefo, G. O. and Vwioko, D. E. (1994). Effects of spent lubricating oil on the growth of
171 *Capsicum annum* (L) and *Lycopersicon esculentum* (Miller). *Environment and Pollution*,
172 88:361-364.
- 173 Aroh, K.N. Ubong, I.U. Eze, C.L. Harry, I.M. Umo-Otong, J.C. and Gobo, A.E. (2010). Oil spill
174 incidents and pipeline vandalization in Nigeria, impact on public health and negation to
175 attainment of Millennium development goal: the Ishiagu example. *Disaster Prevention*
176 *and Management*, 19 (1): 70-87
- 177 Ayotamuno, J M. and Kogbara, R. B. (2007). Determining the tolerance level of *Zea mays*
178 (maize) to a crude oil polluted agricultural soil. *African Journal of Biotechnology*, 6 (11):
179 1332-1337
- 180 Baker, J.M. (1970) The Effects of Oils on Plants. *Environmental Pollution*, 1: 27-44.
181
- 182 Barua, D., Buragohain, J. and Sarma, S. K. (2011). Certain physico-chemical changes in the soil
183 brought about by contamination of crude oil in two oil fields of Assam, NE India.
184 *European Journal of Experimental Biology*, 1 (3):154-161
- 185 Bello, E.I. Aladesanwa, R.D. Akinlabi, S.A. and Mohammed, T.I. (1999). Effects of gas flaring
186 on the growth and yield of maize (*Zea mays* L.) in South – Eastern Nigeria. *Applied*
187 *Tropical Agriculture*, 4 (1): 42 – 47

- 188 Cutler DF, Botha T, Stevenson D, Plant anatomy: an applied approach. Blackwell Publishing,
189 Malden, MA, USA, 2007, pp 133
- 190 Dung, E. J. Bombom, L. S., and Agusomu, T. D. (2008). The effects of gas flaring on crops in the
191 Niger Delta, *Nigeria. Geo Journal*, 73 (4): 297 – 305.
- 192 Ekundayo, E.O. Emede, T.O. and Osayande, D. J. (2001). Effects of crude oil spillage on growth
193 and yield of maize (*Zea mays* L.) in soil of Midwestern Nigeria. *Plant Food for Human*
194 *Nutrition (formerly Qualitas Plantum)*, 56 (4): 313 – 324.
- 195 Food and Agriculture Organization (FAO) (2002). World Agriculture: towards 2015/2030.
196 Summary report, Rome. Available at <http://www.fao.org/3/y3557e/y3557e00.htm>
197 (Accessed 27th October, 2019)
- 198 Naegele, J. A. (1974). Effect of pollution on plants. In: Sax, I. N. (ed.). *Industrial pollution*. Van
199 Nostrand Reinhold Publishing Co, New York.
- 200 Odu, C.T.I. (1983). The oil industry and the environment. *The Nigerian Accountant*, 16 (1): 23-
201 26, 47.
- 202 Olaniyan, A. B. and Lucas, E. O. (2004). Maize hybrids cultivation in Nigeria - A review.
203 *Journal of Food, Agriculture & Environment*, 2 (3&4): 177-181.
- 204 Quinones-Aquilar, E. E., Ferrera-Cerrato, R. Gavi, R. F., Fernandez, L., Rodriguez, V. R. and
205 Alarcom, A. (2003). Emergence and growth of maize in a crude oil polluted soil.
206 *Agrociencia*, 37: 585-594.
- 207 Ubochi, K. C., Nweze, N. C. and Ojua, E. O. (2019). Effects of coconut oil effluent (ICE)
208 irrigation practice on some soil chemical properties and nutrients composition of *Talinum*
209 *fruticosum* L. *International Journal of Ecology and Environmental Sciences*, 45 (3): 303-
210 310
211