

**Inhibitory Studies of *Tamarindus indica* Seed Extract and Fractions  
on Hematological Activities of *Bitis arietans* Venom.**

**ABSTRACT**

Envenomation is a serious public health and occupational hazard in tropical and subtropical countries. Antivenom serum treatment has been related to extreme hypersensitivity, high cost, and inaccessibility. Consequently, therapeutic plants have been investigated to give an alternative treatment. The aim of this study was to evaluate the ability of the methanolic extract and fractions of *Tamarindus indica* seeds to neutralize the hemolytic, anticoagulant and hemorrhagic activities of *Bitis arietans* venom. The powdered form of *Tamarindus indica* seed was extracted using methanol. The methanol extract was further fractionated using butanol solvent. The aqueous fraction of the methanolic extract inhibited hemolysis caused by crude venom by 95.5%, while methanolic extract significantly reduced the anticoagulant time of crude venom to 10.2%. The *in vivo* hemorrhagic activity of the crude venom of *Bitis arietans* was neutralized by both the crude methanolic extract and the subsequent two fractions of *Tamarindus indica* seed. However, the crude methanolic extract was found to be the most active by reducing the hemorrhage diameter caused by crude venom to 39%. The findings of this present research work demonstrated a very good antivenom properties of the seeds of *Tamarindus indica*, which suggests that it could be a potential source of effective, and safe compounds with inhibitory activities against *Bitis arietans* envenomation.

**KEYWORDS:** Antivenom, *Bitis arietans*, *Tamarindus indica*, Hemolytic, Anticoagulant, Hemorrhagic.

**1. INTRODUCTION**

Snakebite is an important public health and occupational hazard often faced by farmers, farm laborers, hunters and nomads of tropical and subtropical countries present in Asia, Africa and South America [1].

Envenomation by venomous snake is a common acute life-threatening and time-limiting medical emergency. Globally, approximately 5.4 million snakebites occur each year, resulting in 1.8 to 2.7 million cases of envenomings, with mortality ranging from 81 410 to 137 880 deaths and around three times as many amputations and other permanent disabilities each year [2, 3]. The extent of snakebite globally, resulted the World Health Organization (WHO) declaration of snakebite as a neglected tropical disease in 2009. However, it was removed in 2013, and re-declared as category A on the 9th of June 2017 [4].

In Nigeria alone an estimate of 497 per 100,000 populations are bitten annually by venomous snakes. Nigeria is therefore reported to have one fifth of all West African region cases of snakebites [5]. However, an accurate measure of snakebites remains elusive as most victims approach traditional healers for aid and treatment. Snakebite is a major challenge that affects farmers, nomads and rural dwellers of all ages. Inadequate facilities at the primary health center (PHC), ignorance of conventional treatment of snakebite by medical doctors; further delays inappropriate treatment of victims leads to increased morbidity and mortality [6]. Treatment with antivenom serum remains the only specific therapy for snakebite envenomation. However, it is heterologous and therefore liable to cause adverse reactions, such as early anaphylactic, pyrogenic and delayed reactions [7]. Medicinal plants are rich sources of natural inhibitors that tend to reduce the pharmacological activities of various snake venom without resulting to any side effects [8].

*Bitis arietans* commonly known as Puff adder is a venomous snake thought to be the most common and widespread snake in Africa, occurring throughout the continent except in some forested regions and extreme deserts such as the Sahara. *Echis ocellatus* is responsible for most snakebite incidence in Nigeria. *Bitis arietans* is third in terms of medical importance [9]. This is due to a combination of factors, including its wide distribution, common occurrence, large size, potent venom that is produced in large amounts, long fangs, and sitting quietly when approached. The venom has cytotoxic effects and is one of the most toxic of any vipers based on LD<sub>50</sub> [10]. Proteomic analyses has shown that hemorrhagins and hemolysins such as phospholipases A<sub>2</sub>, metalloproteases, serine proteases, disintegrins, L-amino acid oxidase, Kunitz inhibitors, cystatins and C-type lectins are present in *Bitis arietans* venom [11].

*Tamarindus indica* of the Fabaceae, subfamily Caesalpinioideae, is an important food in the tropics. It is a multipurpose tree of which almost every part finds at least some use either nutritional or medicinal [12]. Asian countries such as India and Myanmar have used *Tamarindus indica* for the treatment of snakebite and can be found in their traditional medicine book [13, 14]. However, based on our knowledge, no scientific study has been done to establish this in Nigeria. It has been noted that several anti snake venom activities are known to contain phenolic compounds [15].

Tamarind seeds are reported to contain phenolic antioxidants, procyanidins, epicatechin, taxifolin and eriodictyol. This plant is readily available; therefore this will resolve the problems of accessibility, cost, and storage of antivenom immunotherapy treatment encountered by patients. Hence, the proper utilization of the antivenom properties of *T. indica* seed by researchers, could help provide an alternative treatment that would curb the challenges of envenomation in rural areas.

The aim of this study was to determine the inhibitory effect of methanolic extract and fractions of *Tamarindus indica* seed on hemolytic, hemorrhagic and anti coagulant activities of crude venom of *Bitis arietans*.

## **2. MATERIALS AND METHODS**

### **2.1 Chemicals and Reagents**

All solvents and All other chemicals: sodium dihydrogen phosphate, Hydrogen disodium phosphate, sodium chloride, calcium chloride glycerol, Sodium hydroxide, sodium citrate were all purchased from Sigma Chemical Co. St. Louis, England and were of analytical grade.

### **2.2 Venom Collection**

Two adult *Bitis arietans* were sourced from Fitika village, Giwa Local Government of Kaduna state, and allowed to acclimatize at the Herpetarium, Department of Veterinary Pharmacology and Toxicology, Ahmadu Bello University Zaria. The venom from the *Bitis arietans* was milked as described by Markfarlane [16].

### **2.3 Experimental Animals**

Male and female adult albino mice (20-30g) were used for the research. The mice were procured from animal house of the Department of Pharmacology and Therapeutics Faculty of Pharmaceutical Sciences, Ahmadu Bello University, Zaria.

### **2.4 Preparation of Plant Extract**

*Tamarindus indica* seed was obtained from Sabon-gari, Zaria, Kaduna State and then taken for identification at the Herbarium unit, at Department of Biological Sciences, Ahmadu Bello University Zaria. The dried seeds were crushed into fine powder.

Dried powdered seeds of *Tamarindus indica* (1.2kg) was extracted for 75 hours using n-hexane, ethyl acetate, methanol solvent (2.5L) successively by maceration following the procedure described by Sukito and Tachibana [17].

### **2.5 Fractionation of Extract of *Tamarindus indica* Seed**

The methanol seed extract (138.25g) was dissolved in water (750ml) and placed in a separating funnel and then 1.25L of butanol was added mixed thoroughly and allowed to stand for 24 hours. The butanol fraction was collected carefully subsequently the aqueous fraction. They were both dried using water bath at 40<sup>o</sup>c and then weighed. This procedure was carried out as described by Abubakar [18].

### **2.6 Hemolytic Assay of Crude Venom of *Bitis arietans***

The method described by Gomes and Pallabi [19] was adopted using bovine erythrocytes to determine hemolytic activity. Twenty milliliters (20ml) of bovine blood was collected from Zango abattoir, Zaria, Kaduna State, using sodium citrate as anticoagulant. Blood sample was centrifuged at 646 x g for 10min and the plasma discarded. Five milliliters (5ml) of normal saline was mixed with the packed cell layer and centrifuged at 646 x g for 10min and the supernatant discarded. This procedure was repeated 10 times to obtain plasma-free packed cells. Venom solution (5mg/ml) was mixed with 1ml of 1% cell suspension in saline and the mixture was incubated at 37<sup>o</sup>C for 1hr. the reaction was stopped by adding 3ml of chilled phosphate buffer saline. Then centrifuged at 449 x g for 10mins and absorbance was measured at 540nm. For inhibitory studies 1ml venom solution, with 1 ml of 1% v/v cell suspension in saline was

incubated with 1ml of different concentrations of plant extract fractions at 37°C for 1hr. The reaction was also been stopped by adding 1ml of chilled phosphate buffer then centrifuged at 449 x g 10min and absorbance was measured at 540nm. The control sample consist of 1% cell suspension treated with 3 ml chilled water, was considered as 100 % hemolysis.

% Inhibition of hemolysis =  $100 \times [\text{OD1}-\text{OD2}/\text{OD1}]$

Where: OD1 = Optical density of hypotonic saline solution alone

OD2 = Optical density of test sample.

## **2.7 Anticoagulant activity Assay of Crude Venom of *B. arietans***

Bovine blood sample of 20ml was collected in 3.8% sodium citrate, centrifuged at 701 x g and 4 °C for 2 hours to obtain platelet-poor plasma. Coagulant activity was measured by the method of Condrea *et al* [20] using Bovine citrate plasma (2.0mg/ml). Fifty microliters of CaCl<sub>2</sub> (25mM) was added then, 50uL of venom was added to 0.2 ml of citrated plasma solution and clotting time was recorded at 37°C. For inhibitory studies, 0.2ml of citrated plasma solution was added to 1ml of CaCl<sub>2</sub> (25mM), then the venom alongside different concentrations of plant extract fraction. The mixture was incubated at 37°C for 2 hours and the clotting time was recorded [21].

## **2.8 Haemorrhagic Assay of Crude Venom of *Bitis arietans***

The method described by Omori-Satoh *et al.*, [22] was employed. *Bitis arietans* venom (5mg/ml) was used where 0.2ml was injected sub-cutaneously into the backs of mice. The mice were sacrificed by cervical fracture after 60min, and then skinned and the haemorrhagic foci was measured in mm. For inhibitory studies, the venom was injected alongside with different concentrations of plant extract and fractions into mice. The mice were sacrificed after 2 hours and the haemorrhagic foci was measured in mm as described by [22, 23].

Thirty (30) mice were assembled and divided into 5 groups of 6 mice each.

Group 1: Untreated control was treated with normal saline.

Group 2: Treated control was treated with Crude venom alone.

Group 3: treated with aqueous fraction and Crude Venom

Group 4: treated with n-butanol fraction and Crude Venom

Group 5: treated with methanolic extract and Crude Venom


## 2.9 Statistical Analysis

Experimental results were expressed as means  $\pm$  SD. The data was analyzed by an analysis of variance (at  $p < 0.05$ ) and the means separated by Duncan's multiple range tests. All statistical analysis was performed using coupled Microsoft Excel Software and; Statistical Package for Social Sciences (SPSS) version 20.0

## 3. RESULT


### 3.1 *In Vitro* Studies Activity of Solvent Fractions of *Tamarindus Indica* Seed Extracts on Crude Venom of *Bitis arietans*

The hemolysis caused by distilled water was considered as 100% hemolysis of the RBC, which served as control. The hemolytic activity on RBC by the crude venom of *Bitis arietans* as shown in (Fig 1) revealed a significantly high value of  $65.8\% \pm 0.26$ . At the highest concentration of  $10\text{mgml}^{-1}$  the crude methanolic extract, butanolic fraction and aqueous fraction of *T. indica* seed incubated with  $5\text{mgml}^{-1}$  of crude *B. arietans* venom significantly (at  $p < 0.05$ ) reduced hemolysis of RBC to  $8.1\% \pm 2.51$ ,  $11.7\% \pm 2.84$  and  $4.5\% \pm 2.62$  respectively. At the least concentration of  $2.5\text{mgml}^{-1}$  the methanolic extract, butanolic fraction and aqueous fraction reduced hemolysis to  $55.8\% \pm 1.85$ ,  $73.9\% \pm 4.56$  and  $59.4\% \pm 1.11$  respectively, suggesting that the methanolic extract, aqueous fraction and butanolic fraction *T. indica* seed protected the red blood cells from being lysed by toxins such as  $\text{PLA}_2$  present in *Bitis arietans* venom.



**Figure 1:** Effect of *T. indica* Seed Extracts on the Hemolytic Properties of *B. arietans* Venom. Bars Represent the Mean  $\pm$ SD: with different superscripts are significantly different at  $P < 0.05$  by Duncan's Multiple Range test.

The anticoagulant activity was carried out and the coagulation time was monitored for 2 hours. The crude venom of *B. arietans* did not yield clot formation after 2 hours of incubation, and this was considered 100% anticoagulation time as shown in (Fig 2). Aqueous fraction, methanolic extract and butanolic fraction significantly reduced the anticoagulant time to  $16.70 \pm 1.88\%$ ,  $10.21 \pm 0.62\%$  and  $47.53 \pm 1.61\%$  respectively at  $10 \text{mgml}^{-1}$  concentration. While at the least concentration ( $2.5 \text{mgml}^{-1}$ ), aqueous fraction, the methanolic extract, and butanolic fraction also significantly reduced the anticoagulant time to  $50.21 \pm 3.26\%$ ,  $50.08 \pm 2.24\%$  and  $63.44 \pm 6.04\%$  respectively, but not as compared to the highest concentration.




**Figure 2:** Effect of *T. indica* Seed Extracts on the Anticoagulant Properties of *B. arietans* Venom. Bars Represent the Mean  $\pm$ SD: with different superscripts are significantly different at  $P < 0.05$  by Duncan's Multiple Range test.

### 3.2 *In vivo* Activity of Solvent Fractions of *Tamarindus Indica* Seed Extracts on Crude Venom of *Bitis arietans*

The crude *B. arietans* venom resulted to an MLD of  $32 \pm 3$ mm, which was considered 100% hemorrhage. The methanolic extract, butanolic fraction, and aqueous fraction also significantly (at  $p < 0.05$ ) reduce the diameter of lesion at high concentration to  $39.1\% \pm 0.17$ ,  $40 \pm 0.1$ , and  $42.9 \pm 0.2$  respectively as shown in (Fig 3). This is to suggest that the crude methanolic extract, butanolic fraction and aqueous fraction of *T. indica* seed were able to reduce degradation of the capillary vessel in mice induced by PLA<sub>2</sub> isoform and crude venom of *B arietans*.





**Figure 3:** Effect of *T. indica* Seed Extracts on the Hemorrhagic Properties of *B. arietans* Venom. Bars Represent the Mean  $\pm$ SD; with different superscripts are significantly different at  $P < 0.05$  by Duncan's Multiple Range test.

#### 4. DISCUSSION

The antihemolytic activity implies that fractions of *T. indica*, especially aqueous consist of bioactive compounds that interfere with the mechanism, in which haemolysins such as PLA<sub>2</sub> responsible for hemolysis present in the venom of *Bitis arietans*, interacts with phospholipids on cell membrane that results to hydrolysis of intact membrane releasing free fatty acids and lysopholipids [24, 25, 26].

Medicinal plants have been recorded to deactivate proteins and peptides because of diverse secondary metabolite like polyphenols, terpenoids, and flavonoids which have been reported to neutralize snake venom [27, 28, 29, 30]. Similar results were also shown by the methanolic extract of *Indigofera pulchra* and *Aristolochia abida* plants against hemolysis caused by *Naja nigricollis* venom [18]. This work conforms to

findings of Ushanandini *et al.*, [31], who reported that crude ethanolic extracts of *T.indica* were found to neutralize the hemolytic activity of *Russelli* viper.

Anticoagulation and spontaneous bleeding are some of the hematological effects of viper envenomation in victims. These research study, revealed that treatment with the methanol crude extract and aqueous fraction neutralized the anticoagulant effect of crude venom of *B. arietans* in a concentration dependent manner, where an increase in extract concentration reflected a significant reduction in anticoagulant activity of *B. arietans* venom. These findings suggest that the aqueous fraction constitute of bioactive compounds that inhibits haemolysins and haemorrhagic toxins. PLA<sub>2</sub> is known to prevent the formation of the extrinsic tenase complex (TF, FVIIa and Ca<sup>2+</sup>) and the activation of FX to FXa required for the formation of prothrombinase complex, a vital component of blood coagulation [32, 33]. It is worth noting that the activation of FX and prothrombin are both totally dependent on the presence of phospholipid membrane [34, 35, 36]. Aside from PLA<sub>2</sub> in the venom of *B. arietans*, Zn<sup>2+</sup> metalloproteinase and other nonenzymatic toxins have been shown to degrade A $\alpha$  and B $\beta$  chains of fibrinogen and act upon the coagulation factors and complexes [37, 1, 38]. It is possible that bioactive compound present in aqueous fraction and crude methanolic extract of *T. indica* antagonize the anticoagulant activities of Zn<sup>2+</sup> metalloproteinase and other toxins in an unknown mechanism thereby preventing the hydrolysis of A $\alpha$  and B $\beta$  chains of fibrinogen. This work further authenticates the findings of Ushannidini *et al.*, [31] who reported that the ethanolic extracts of *T. indica* was able to antagonize the defibrinogenating activity of *V. russelli* venom at a concentration of 25ug. Other studies have shown that the aqueous extract of *J. gossypifolia* at high concentration was able to inhibit fibrinogenolytic enzymes from *B. jararaca* by preventing the degradation of A $\alpha$  and B $\beta$  chains of fibrinogen [8].

Severe hemorrhage is a major challenge associated with envenomation of *Bitis arietans*. This study revealed that the crude venom of *B. arietans* induced a significantly high mean diameter of lesion in mice. This can be ascribed to the cumulative action of haemorrhagins such as, PLA<sub>2</sub>, Zn<sup>2+</sup> metalloprotease and nonenzymatic toxins present in *B. arietans* venom. However, irrespective of the cumulative effect of crude *B. arietans* venom, the methanolic extract and fractions of *T. indica* seed was still able to reduce hemorrhage. The reduction of hemorrhage by *T. indica* suggests that, it contains bioactive compounds that directly inhibits the major hemorrhagic toxins which are PLA<sub>2</sub> and Zn<sup>2+</sup> metalloproteinase thus, preventing the degradation of the

extracellular membrane components of the capillary vessels. It is possible that these bioactive compounds interacts with divalent ion such as  $\text{Ca}^{2+}$  and  $\text{Zn}^{2+}$  which are essential cofactors of  $\text{PLA}_2$  and SVMP respectively, thereby leading to the inhibition of their enzymatic activity.

This findings, agrees with the report by Esmeraldino *et al.*, who showed that the butanolic fraction of *C. urucurana* plant was able to inhibit hemorrhage of *B. jararaca* venom by 97%, which was ascribed to catechin, a basic unit of condensed tannins [39]. Similar findings by Pithayankul *et al.*, showed that the butanolic extract of *Eclipta* plant was able to neutralize the lethal and hemorrhagic effect of Malaysia pit viper venom and speculated that it could be due to the hydrogen interaction between the phenolic hydroxyls and the carbonyl group in the peptide bonds of the venom protein [40].

## 5. CONCLUSION

Viper envenomation is a major life-threatening event among healthy individuals in tropical and subtropical countries. In viper venoms, phospholipase  $A_2$  and proteases are extensively studied hydrolytic enzymes due to their relatively high abundance and participation in a wide range of pharmacological and toxic effects.

The methanolic extract, aqueous fraction and butanolic fraction of *T indica* seed was able to neutralize the pharmacological activities of *B. arietans* which included hemolytic, anticoagulant and hemorrhagic effects which are key systemic and local effect in *B. arietans* envenomation. This indicates that methanolic extracts of *T. indica* seed consist of compounds that might interfere with the combined effects of the enzymes and toxins present in *B. arietans* venom. The findings of this present research work demonstrated a very good antivenom properties of the seeds of *Tamarindus indica*, which suggests that it could be a potential source of effective, safe and cost efficient antivenom agent for treatment of *Bitis arietans* envenomation.

## REFERENCE

1. Menaldo, D., L. , Anna L. Jacob-Ferreira, Carolina P. Bernardes, Adélia C. O. Cintra, and Sampaio, S. V. (2015). Purification procedure for the isolation of a P-I metalloprotease and an acidic phospholipase  $A_2$  from *Bothrops atrox* snake venom. *Venomous Animals and Toxins including Tropical Diseases*, 21(28), pp. 1-14.
2. Reis, V. P. D., Boeno, C. N., Evangelista, J. R., Santana, H. M., Serrath, S. N., Lopes, J. A., ... & Dantas, A. D. S. (2020). Increase in the risk of snakebites incidence due to changes in humidity

levels: A time series study in four municipalities of the state of Rondônia. *Revista da Sociedade Brasileira de Medicina Tropical*, 53.

3. WHO. WHO | snakebite envenoming. WHO [Internet]. [Cited 2019 Apr 9]; <https://www.who.int/news-room/fact-sheets/detail/snakebite-envenoming>
4. Chippaux JP. (2017). Snakebite envenomation turns again into a neglected tropical disease. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 23:38, pp.1-2
5. Nasidi, A. (2007). *Snakebite as a serious Public Health problem for Nigeria and Africa*. Paper presented at the WHO Consultative Meeting on Rabies and Envenomings: A Neglected Public Health Issue, Geneva, Switzerland.
6. Bawaskar, H., Bawaskar PH, and PH, B. (2014). Premonitory signs and symptoms of envenoming by common krait (*Bungarus caeruleus*). *Trop Doct*, 44(82), pp 1-5.
7. Gilbert, L. A., Horlbeck, M. A., Adamson, B., Villalta, J. E., Chen, Y., and Whitehead, E. H. (2014). Genome-scale CRISPR-mediated control of gene repression and activation. *Cell*, 159(3), pp 647-661.
8. Fe'lix-Silva, J., Souza, T., Menezes, Y. A. S., Cabral, B. r., ^mara., R. B. G. C., Silva-Junior., A. b. A., and Fernandes-Pedrosa., M. F. (2014). Aqueous Leaf Extract of *Jatropha gossypifolia*L.(Euphorbiaceae) Inhibits Enzymatic and Biological Actions of *Bothrops jararaca*Snake Venom. *Plos one*, 9(8), pp. 1-14.
9. Abubakar, I. S., Abubakar, S. B., Habib, A. G., Nasidi, A., Durfa, N., Yusuf, P. O., ... and Theakston, R. D. G. (2010). Randomised controlled double-blind non-inferiority trial of two antivenoms for saw-scaled or carpet viper (*Echis ocellatus*) envenoming in Nigeria. *PLoS Neglected Tropical Diseases*, 4(7), e767.
10. Mallow, D., Ludwig D, and G., N. ( 2003). True Vipers: Natural History and Toxinology of Old World Vipers. *Krieger Publishing Company, Malabar, Florida.*, pp 359.
11. Megale, Â. A. A., Magnoli, F. C., Kuniyoshi, A. K., Iwai, L. K., Tambourgi, D. V., Portaro, F. C., & da Silva, W. D. (2018). Kn-Ba: a novel serine protease isolated from *Bitis arietans* snake venom with fibrinogenolytic and kinin-releasing activities. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 24(1), 1-11.
12. Kuru, P. (2014). *Tamarindus indica*and its health related effects. *Asian Pacific Journal of Tropical Biomedicine*, 4(9), pp. 676-681.
13. Maung, K. M., & Lynn, Z. (2012). Effects of Tamarind (*Tamarindus indicus* Linn) seed extract on Russell's viper (*Daboia russelli siamensis*) venom. *Tropical biomedicine*, 29(4), 580-587.
14. Al-Ani, I. M., Ismail, S., Maung, K. M., Oothuman, P., & Al-Mahmood, S. M. A. (2017). Histological Study On The Protective Effects Of Tamarind Seed Extract On Cobra Venom In Mice. *Asian J Pharm Clin Res*, 10(10), 301-305.
15. Asad, M. H. H. B., D. E. Sabih, B. A. Chaudhory, I. Ahmad, M. S. Hussain, and Hussain, I. (2014). Anti-hemolytic property of local medicinal plant(s) upon pakistani cobra venom induced hemolysis. *The Journal of Animal & Plant Sciences*, 24(6), pp 1701-1708.

16. Markfarlane, R. G. (1967). Russel's Viper Venoms,. *British Journal of Haematology*, 13, pp 437-451.
17. Sukito, A., and Tachibana, S. (2014). Potent alpha-glucosidaseinhibitors isolated from Ginko biloba leaves. *Park. journal of biological science*, 17(11), pp 1170-1178.
18. Abubakar, M.S., , E. Balogun , E.M. Abdurahman , A.J. Nok , M. Shok , Mohammed, A., and Garba., M. (2006). Ethnomedical Treatment of Poisonous Snakebites: Plant Extract Neutralized Naja nigricollis. *Venom.*, 44(5 ), Pp 343-348.
19. Gomes, A., and Pallabi, D. (1999). Hannahpep: a novel fibrinolytic peptide from the Indian King Cobra (*Ophiophagus hannah*) venom. *Biochemical and biophysical research communications*, 266(2), pp 488-491.
20. Condrea, E., Yang, C. C., & Rosenberg, P. (1983). Anticoagulant activity and plasma phosphatidylserine hydrolysis by snake venom phospholipases A<sub>2</sub>. *Thrombosis and haemostasis*, 49(2), 151.
21. Theakston, R. D. G., and Reid, H. A. (1983). Development of simple standard assay procedures for the characterization of snake venoms. *Bulletin of the world health organization*, 61(6), pp 949.
22. Omori-Satoh, T., Sadahiro, S., Ohsaka, A., and Murata, R. (1972). Purification and characterization of an antihemorrhagic factor in the serum of *Trimeresurus flavoviridis*, a crotalid. *Biochimica et Biophysica Acta (BBA)-Protein Structure*, 285(2), pp 414-426.
23. Moraes, F. V. D., Sousa-e-Silva, M. C. C., Barbaro, K. C., Leitão, M. A., and Furtado, M. F. D. (2003). Biological and immunochemical characterization of *Micrurus altirostris* venom and serum neutralization of its toxic activities. *Toxicon*, 41(1), pp 71-79.
24. Kini, R. M. (2003). Excitement ahead: structure, function and mechanism of snake venom phospholipase A<sub>2</sub> enzymes. *Toxicon*, 42: 827-840.
25. Saikia, D., Bordoloi, N. K., Chattopadhyay, P., Choklingam, S., Ghosh, S. S., and , and Mukherjee, A. K. (2012). Differential mode of attack on membrane phospholipids by an acidic phospholipase A<sub>2</sub> (RVVA-PLA 2-I) from *Daboia russelli* venom. *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1818(12), pp 3149-3157.
26. Gasanov , S. E., Dagda, R. K., and Rael, E. D. (2014). Snake Venom Cytotoxins, Phospholipase A<sub>2</sub>s, and Zn<sup>2+</sup>-dependent Metalloproteinases: Mechanisms of Action and Pharmacological Relevance. *Clinical Toxicology*, 4(1), 1-14.
27. Asad, M. H. H. B, G. Murtaza, S. Siraj, S. A. Khan, S. Azhar, M. S. Hussain, T. Ismail, M. S. Hussain and I. Hussain (2011). Enlisting the scientifically unnoticed medicinal plants of Pakistan as a source of novel therapeutic agents showing anti-venom activity. *African Journal Pharmacy and Pharmacology*, 5(20): 2292-2305.
28. Samy, R. P., Gopalakrishnakone, P., Bow, H., Puspharaj, P. N., and Chow, V. T. K. (2010). Identification and characterization of a phospholipase A<sub>2</sub> from the venom of the Saw-scaled viper: Novel bactericidal and membrane damaging activities. *Biochimie*, 92, pp. 1854- 1866.
29. Wink, M. (2015). Modes of action of herbal medicines and plant secondary metabolites. *Medicines*, 2(3), 251-286.
30. Sarkar, S., Chakraverty, R., Datta, S., and Ghosh, A. (2015). In-Vitro Assays for Neutralization of Snake Venom Using Herbal Drugs: A. *Creative Journal of Pharmaceutical Research* 1(4) 156(169).

31. Ushanandini, S., Nagaraju S, Kumar Harish K, Vedavathi M, Machiach DK, and Kemparaju K. (2006). The anti snake venom properties of *Tamaridus indica* ( Leguminosae) seed extract. *Phytother Res*, 20(10), pp. 851- 858.
32. Chérifi, F., and Laraba-Djebari, F. (2013). Isolated biomolecules of pharmacological interest in hemostasis from *Cerastes cerastes* venom. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 19(1), 11.
33. Faure, G., Gowda, V. T., and Maroun, R. C. (2007). Characterization of a human coagulation factor Xa-binding site on Viperidae snake venom phospholipases A 2 by affinity binding studies and molecular bioinformatics. *BMC structural biology*, 7(1), 82.
34. Xu, Z., Lioi, J., Mu, J., Kamocka, M. M., Liu, X., Chen, D. Z., and Alber, M. (2010). A multiscale model of venous thrombus formation with surface-mediated control of blood coagulation cascade. *Biophysical journal*, 98(9), 1723-1732.
35. Spronk, H. M., ten Cate, H., and van der Meijden, P. E. (2014). Differential roles of tissue factor and phosphatidylserine in activation of coagulation. *Thrombosis research*, 133, S54-S56.
36. Schuijt, T. J., Bakhtiari, K., Daffre, S., DePonte, K., Wiolders, S. J., Marquart, J. A., and Camire, R. M. (2013). Factor Xa activation of factor V is of paramount importance in initiating the coagulation system: lessons from a tick salivary protein. *Circulation*, CIRCULATIONAHA-113.
37. Das, D., Urs, N., Hiremath, V., Vishwanath, B. S., & Doley, R. (2013). Biochemical and biological characterization of *Naja kaouthia* venom from North-East India and its neutralization by polyvalent antivenom. *Journal of venom research*, 4, 31.
38. Cintra, A. C. O., De Toni, L. G. B., Sartim, M. A., Franco, J. J., Caetano, R. C., Murakami, M. T., and Sampaio, S. V. (2012). Batroxase, a new metalloproteinase from *B. atrox* snake venom with strong fibrinolytic activity. *Toxicon*, 60(1), 70-82
39. Esmeraldino, L. E., A.M. Souza, and Sampaio, S. V. (2005). Evaluation of the effect of aqueous extract of *Croton urucurana* Baillon (Euphorbiaceae) on the hemorrhagic activity induced by the venom of *Bothrops jararaca*, using new techniques to quantify hemorrhagic activity in rat skin. *Phytomedicine*, 12, pp.570–576.
40. Pithayanukul, P., Laovachirasuwan., S., Bavovada., R., Pakmanee., N., and Suttisri., R. (2004). Anti-venom potential of butanolic extract of *Eclipta prostrata* against Malayan pit viper venom. *Journal of Ethnopharmacology*, 90, pp.347–352.

## LIST OF ABBREVIATION

ml = Milliliter

mM = Millimolar

mg = Milligram

PLA<sub>2</sub> = Phospholipase A<sub>2</sub>

CP = Citrated plasma

SVMP = Snake Venom Metalloprotease

RBC = Red Blood Cell

Aqu = Aqueous fraction

Butanol = Butanol fraction

Meth = Methanol extract

V = Venom

MLD = Mean Lesion Diameter

WHO = World Health Organization

UNDER PEER REVIEW