

Effect of organic mulching on weed suppression, yield and yield contributing components of sesame (*Sesamum indicum* L.)

Abstract

Sesame is the second in oil crop produced in Ethiopia next to niger seed and it is also 2nd foreign currency generator in the country. Weed is the most significant problem in sesame producing areas. This research was conducted with the aim of reducing weed infestation and boost yield. The experiment was conducted in 2016 cropping season at Humera Agricultural Research Center in RCBD design. Growth performance, yield and yield component and weed data collected. Less abundant and dense *Rhynchosiamalacophylla* weeds (33& (132m⁻²)) were recorded at sudan grass where the most abundant and dense *Rhynchosiamalacophylla* weeds (60.5& (242m⁻²)) recorded at no mulch plot. The highest yield (695kg^{ha}⁻¹) was recorded at sudan grass while the lowest yield (225kg^{ha}⁻¹) was recorded at no mulch. Farmer should apply grass mulch to suppress weed growth and boost yield.

Introduction

Sesame (*Sesamum indicum* L.) is an important oil crop of Ethiopia, which belongs to the Pedaliaceae family. It is short day plant that grows mainly for its vital seed that contains about 50% oil and 25% protein [1]. Sesame is the second in oil crop produced in Ethiopia next to niger seed and it is also 2nd foreign currency generator in the country. Sesame is a warm season oil crop which is chiefly adapted to areas with extended growing period and well aerated soils. Sesame performs well in slightly acid to alkaline soils (pH 5-8) with moderate fertility [2].

Weed is the most significant problem in sesame producing areas of Ethiopia particularly in western zone of Tigray which can cause a yield loss up to 86.3% [3]. This could be due to lack of option for weed control except hand weeding method. In sesame growing areas of western Tigray there is a strong interest in developing weed control options that are organic nature and

30 can keep its international market quality. So application of organic mulching can reduces weed
31 growth and used as a means of weed control option through the world [4]. Organic mulching is
32 important in crop production as they can suppress weed growth by different scholars [4]; [5],[6]
33 [7] and [8] . Moreover, organic mulch can enhance soil fertility after decomposition at the field.
34 Sønsteby et al., [9] also noted wood chips mulch fertile a soil with phosphorus and potassium.
35 Mulch also increases soil moisture by increasing infiltration and reducing evaporation. In this
36 regard [10] reported high soil moisture conservation for sudan grass, sesame and sorghum
37 mulches respectively. This in conformity with [11]. Mulch do not only suppress weed growth,
38 they conserve also moisture in the root zone [12] increase also yield.

39 .


- 40 ➤ To quantify the efficacy of organic mulching on weed infestation
- 41 ➤ To determine effect of organic mulching on yield and yield component

42

43 Material and method

44 2.1. Area description.

45 The field experiment was carried out in 2015 main growing season in Humera Agricultural
46 Research Center Western zone of Tigray Ethiopia. It is positioned between 13°14 to 14°27N and
47 36°27 to 37°32E and about 600 km from Mekelle capital city of Tigray regional state. The
48 dominant soil type of the area is chromic vertisol black in color which is characterized with bulk
49 density of 1.4 g cm⁻³, organic matter content of 1%, pH=8.5 and EC=0.2 mmohs/cm from
50 analyzed result of pre sowing soil sampled. According to Köppen climate classification Humera
51 has a hot semi-arid climate. The annual rainfall ranges between 400 to 600 mm and most of the
52 rain rains in June up to September. The annual mean temperature is 27.6 °C. It is also
53 characterized by hot temperature, erratic rainfall, vast area of plain low lands suitable for large
54 scale and subsistence agriculture including crops and livestock.



55
56 Fig. 1. Map showing study location in Ethiopia

57

58

59 Experimental Design and Treatment

60 The experimental design used was Randomized Complete Block Design (RCBD) with three
61 replications and five treatments. The tested treatments were sesame straw, sorghum straw, rice
62 straw, Sudan grass, and no mulch. The gross plot area was 6m² and the net plot area was

63 3.6m².The distance between the plot and block was 1 and 1.5m, correspondingly. Sesame seed
 64 variety Setit-1 was sown 40 cm and 10 cm between rows and plants correspondingly. The field
 65 was plowed and provided with 100 kg of NPS (19N-38P2O5 + 7S) and 50 kg of urea fertilizer.
 66 The urea was applied in split form 25kg at sowing and 25 at flower initiation. Webworm pest
 67 which cause a potential yield reduction was control with application of diamatet zenawi 2017.
 68 In harmony with the methods adopted by[13] the rate of mulches used was 6 kg per plot which is
 69 equivalent to 10 ton/ha and was applied at row whole immediately after germination of sesame
 70 and the emerged weed was uprooted using a hoe.

71 Measurements and Measurements Method

72 Weed population was determined by counting number of weed plants in 0.5 m*0.5 m quadrant
 73 which is dropped randomly three times in a plot. Data was collected in four weeks after sowing.
 74 Based on weed count, weed abundance, frequency (eq.1) (eq. 2) and weed dominance (eq. 3)
 75 were determined.

76 Weed abundance: it is the population of a species expressed as the number of individuals of
 77 weed per unit area. It was calculated as follows:

78 $A = \frac{\sum w}{N}$ (Equation 1)

79 Where,

80 **W** = number of individuals of a species/sample

81 **N** = number of sample

82
 83 Weed dominance: abundance of individual of species in relation to total abundance. it is
 84 calculated as:

85 $D = \frac{A * 100}{\sum A}$ (Equation 2)

86 $Density = \frac{No.of\ weed\ of\ agiven\ species}{area}$ (Equation 3)

87 Agronomic Traits of Sesame

88 Number of capsules per plant and number of seeds per capsule were counted from five
 89 randomly selected plants per plot. Plant height is one of the growth parameter that was
 90 measured from five plants per plot. To determine the number of seeds per capsule, the seeds

91 of three capsules (lower, medium, and uppermost position on the plant) from each of five
92 plants were counted. Seed yield of each plot was weighed in grams and converted to area
93 basis to determine the yield kg/ha.
94

UNDER PEER REVIEW

95 Weed growth

96 The different grassy and broad leaved weeds observed in association with sesame crop in the
97 study area were sudan grass (*Topas*), *Rahynchosiamalacophylla*(*Teken*), *Ipomoea*
98 *spp*(*Dereya*), *Indigoferaspp*(*Demayto*), *Commelina Foecunda* (*Wuhankur*) and *Xanthium*
99 *abyssinicum*(*Begdzemed*). Among the two categories of weeds, broad leaved weeds showed
100 more variation in species composition.

101

102

UNDER PEER REVIEW

Result and discussion

Table.1.Effect of organic mulching on weed abundance in sesame fields

treatment	RM		Ip spp		XA		CF		INS		Sg	
	Abundance	Density	Abundance	density	Abundance	density	Abundance	Density	Abundance	density	abundance	density
No mulch	60.5	242.0	20.5	82.0	18.0	72.0	26.0	104.0	17.5	70	18.0	72
Rice straw	56.0	224.0	17.5	70.0	11.0	44.0	14.0	56.0	7.5	30	9.0	36
Sesame straw	27.5	110.0	9.0	36.0	3.3	13.2	9.0	36.0	3.5	14	3.0	12
Sorghum straw	33.0	132.0	17.5	70.0	6.5	26.0	10.5	42.0	5.0	20	3.5	14
Sudan grass	33.0	132.0	10.2	40.7	6.8	27.0	11.5	46.0	6.5	26	7.5	30
CV	32.5	32.4	20	20.0	41.8	41.8	37.6	37.0	40.7	40.7	48.6	48.6
LSD	16.3	65.1	3.6	14.24	4.6	18.2	6.4	25.5	3.9	15.6	4.8	19.0

RM=Rahynochosiamalacophylla XA=Xanthiumabyssinicum INS= *Indigofera* CF= *Commelina Foecunda* IP= *Ipomoea spp*
 sg= Sudan grass

Influence of organic mulch on broad leafed weed in abundance and density

all broad leafed weed such as *Rahynchosiamalacophylla*, *ipomea* spp, *Commelina Foecunda*, *Indigofera* spp and *Xanthium abyssinicum* in sesame field showed highly significant difference at ($p < 0.01$) in abundance and density. Abundant and dense weeds of *Rahynchosiamalacophylla*, *ipomea* spp, was noted at no mulch though statistically insignificant with rice straw. In other hand less abundant and dense weeds of *Rahynchosiamalacophylla*, *ipomea* spp, was noted at sorghum, sesame and sudan grass mulch respectively (Table1). *Commelina Foecunda* and *Indigofera* spp weeds were easily influenced by all organic mulches in abundance and density (Table 1). *Xanthium abyssinicum* weed was influenced by all mulch material compared to bare plot though there was insignificant difference among organic mulches except for sesame straw. This influence of organic mulch on weeds could be due to the nature of mulch material suppress weed growth through direct sunlight prevention to growing weed seedling and suffocation. This result in line with Stiegler et al., [14] who reported dry grass reduced weed infestation significantly. In addition a layer of mulching material prevents weed growth by inhabiting light penetration [15].

Organic mulching had high significant difference at ($p < 0.01$) on sudan grass weed in terms of abundance and density. The highest sudan grass weed abundance and density was recorded at bare plot while the lowest at sesame straw mulch (Table 1).

Influence of organic mulching on weed frequency and dominance

The analysis of variance revealed that organic mulch have significant difference ($p < 0.05$) on weed frequency in all weeds except *Rahynchosiamalacophylla*. All weeds were frequently occurred on the bare and rice mulched plots while less frequent weed occurred on organic mulching except in the case *Rahynchosiamalacophylla* which occurred frequently on all treatment (figure 2).

The analysis of variance also shown that organic mulching have insignificant difference ($p > 0.05$) on weed dominance in the case of all weed except on sudan grass (figure 3).




Fig. 2. Effect of organic mulching on weed frequency




Fig. 3. Effect of organic mulching on weed dominance

Plant height

The analysis of variance showed that organic mulching have highly significant difference ($P < 0.01$) on plant height. The tallest plant height (87.1cm) was recorded at Sudan grass while the shortest plant height (56cm) was recorded at bare plots (Table 2). Since the mulch treatment reduced weed infestation as a result the overall performance of the plant superior compared to bare treatment which is weedy. This result in line with Ajibola, A., et al., [16] reported elephant grass produce highest plant compared to no mulch via weed suppression. This highest plant height in mulch plot could be due to less competition to resource for sesame via controlling undesirable weed.

Number of capsule per plant

The analysis of variance revealed that number of capsule per plant showed highly significant difference on number of capsule per plant. The highest numbers of capsule per plant (39.7 & 37.7) were noted at sudan grass and sesame mulch material while the lowest was (27.1) recorded at bare plots (Table 2) respectively. This could be due to the benefit of mulching material in weed suppression. Sudan grass and sesame straw were effective due to their un decomposed nature. Elephant grass mulching produce comparable result with other weed control option such as hand weeding and herbicide [16]. The lowest number of capsule in no mulch plot could be due to sever competition for resource such as water, nutrient and light and this sever competition prevent from conversion of all produced flower to capsule.

Number of seed per capsule

The analysis of variance shown that organic mulching has highly significant difference ($p < 0.01$) on number of seed per capsule compared to bare treatment. All mulching material didn't show significant difference in between except rice straw. The highest number of seed per capsule was noted in Sudan grass, sesame and sorghum mulching material while the lowest was noted bare treatment (Table 2). The less weed competition in mulch covered plots favors better number of

seed per capsule due to adequate availability of resource for grain filling in the mulch treated plots.

Yield (kg/ha)

The analysis of variance revealed that organic mulching have highly significant difference ($p < 0.001$) on yield compared to bare treatment. All mulching materials have significant difference among them except sorghum and sesame insignificant difference in between. The highest yield (695kg/ha) was recorded at sudan grass while the lowest (225kg/ha) was recorded at bare plots (Table 2). Mulch treatment as means of weed control increased yield 67.6% yield compared to bare plots. This might be due to the mulching material had significant effect on important agronomic parameter such as plant height, number of capsule and number of seed per capsule.

Table.2 Influence of mulching on yield and yield component of sesame

Treatment	PH	NSPP	NCP	yield(kg/ha)
no mulch	56.0 ^d	35.0 ^c	27.1 ^c	225.0 ^d
rice	77.9 ^c	39.8 ^b	29.7 ^c	496.0 ^c
sesame	86.0 ^{ab}	43.5 ^a	37.7 ^{ab}	587.9 ^b
sorghum	82.6 ^b	43.0 ^a	31.8 ^{bc}	550.2 ^b
Sudan grass	87.1 ^a	44.3 ^a	39.7 ^a	695.0 ^a
CV (%)	4.5	3.7	15	7.2
LSD	4.2	1.9	6.1	44.7

Conclusion

The result of the study show that organic mulching had great influence on weed abundance, frequency, dominance and density and as a result good plant performance and yield compared to control.

Conflict of interest

The author revealed that there is no conflict of interest on publication of this paper.

Data availability

The data used to support this finding will be available up on request.

UNDER PEER REVIEW

UNDER PEER REVIEW

Reference

1. Trade, M.o., *Oilseed production and marketing situation in Ethiopia*. C:\Documents and Settings\user1\Desktop\Crop Profiles. mht (15/01/08), 2013.
2. Langham, D.R., *Phenology of sesame*. Issues in New Crops and New Uses, Janick & Whipkey, eds., ASHS Press, Alexandria, VA, 2007: p. 144-182.
3. Mizan, A., W. Gebremedhin, and J. Sharma. *Sesame crops versus weeds: when is the critical period of weed control?* in *9th African Crop Science, Conference Proceedings, Cape Town, South Africa, 28 September-2 October 2009*. 2009. African Crop Science Society.
4. Gupta, G., *Effects of mulching and fertilizer application on initial development of some tree species*. Forest Ecology and Management, 1991. **44**(2-4): p. 211-221.
5. Bhadauria, N., et al., *Integrated weed management in sesame*. Indian Journal of Weed Science, 2012. **44**(4): p. 235-237.
6. Döring, T.F., et al., *Effects of straw mulch on soil nitrate dynamics, weeds, and yield and soil erosion in organically grown potatoes*. Field crops research, 2005. **94**(2-3): p. 238-249.
7. Havlin, J., et al., *Crop rotation and tillage effects on soil organic carbon and nitrogen*. Soil Science Society of America Journal, 1990. **54**(2): p. 448-452.
8. Jodaugienė, D., et al., *The impact of different types of organic mulches on weed emergence*. Agronomy Research, 2006. **4**: p. 197-201.
9. Sønsteby, A., A. Nes, and F. Måge, *Effects of bark mulch and NPK fertilizer on yield, leaf nutrient status and soil mineral nitrogen during three years of strawberry production*. Acta Agriculturae Scandinavica, Section B-Soil & Plant Science, 2004. **54**(3): p. 128-134.
10. Teame, G., A. Tsegay, and B. Abrha, *Effect of Organic Mulching on Soil Moisture, Yield, and Yield Contributing Components of Sesame (Sesamum indicum L.)*. International Journal of Agronomy, 2017. **2017**: p. 6.
11. Sinkevičienė, A., et al., *The influence of organic mulches on soil properties and crop yield*. Agronomy Research, 2009. **7**(1): p. 485-491.
12. Sharma, P.K. and C. Acharya, *Carry-over of residual soil moisture with mulching and conservation tillage practices for sowing of rainfed wheat (Triticum aestivum L.) in north-west India*. Soil and Tillage Research, 2000. **57**(1-2): p. 43-52.
13. Ramakrishna, A., et al., *Effect of mulch on soil temperature, moisture, weed infestation and yield of groundnut in northern Vietnam*. Field Crops Research, 2006. **95**(2): p. 115-125.
14. Stiegler, J.C., et al., *Field-based measurement of ammonia volatilization following foliar applications of urea to putting green turf*. Crop science, 2011. **51**(4): p. 1767-1773.
15. Ossoml, E., et al., *Effect of mulch on weed infestation, soil temperature, nutrient concentration, and tuber yield in Ipomoea batatas (L.) Lam. in Papua New Guinea*. Tropical agriculture, 2001.
16. Ajibola, A., T. Modupeola, and A. Adenuga, "Effect of different weed control practices on growth and yield of sesame in southwest Nigeria," Journal of Biological and Chemical Research, vol. 31, no. 2, pp. 1093–1100, 2014.