

---

## Study of $RL$ -Connectedness and $RL$ -Compactness

---

### Abstract

In this paper, we introduce a new kind of locally closed sets called regular locally closed sets (briefly  $RL$ -closed sets) in a topological space which are weaker than the locally closed sets. Regular locally continuous maps and regular locally irresolute maps are also introduced and studied some of their properties. Finally, we introduce the concept of regular locally connectedness and regular locally compactness on a topological space using the  $RL$ -closed sets.

*Keywords:*  $RL$ -closed set,  $RLC$ -continuous map,  $RLC$ -irresolute map,  $RL$ -connectedness,  $RL$ -compactness.

## 1 Introduction

A topological space is called a connected space if that cannot be represented as the union of two or more disjoint union of non-empty open subsets. Connectedness is one of the principal topological properties that are used to distinguish topological spaces. A subset of a topological space is called a connected set if it is a connected space when viewed as a subspace of that topological space.

In the literature, different type of connectedness were defined and studied by different authors: semipreconnected [1] or  $\beta$  connected [2], preconnected [3], semi-connected [4] using different kind of closed sets. In this order, we define a new kind of closed sets called  $RL$ -closed sets using the concept of locally closed sets and define the connectedness called  $RL$ -connectedness and compactness called  $RL$ -compactness.

## 2 Preliminaries

Throughout this paper, we represent  $X$ ,  $Y$  and  $Z$  as the topological spaces  $(X, \tau)$  and  $(Y, \sigma)$  respectively on which no separation axioms are assumed unless otherwise stated. For a subset  $A$  of  $X$ ,  $cl(A)$  denotes the closure of  $A$  and  $int(A)$  denotes the interior of  $A$ .

We recall the following definitions in the topological space  $X$ .

---

**Definition 2.1.** (5) A subset  $A$  of a topological space  $X$  is called a regular open set if  $A = \text{int}(cl(A))$ . The complement of the regular open set is called the regular closed set.

**Definition 2.2.** (6) A subset  $A$  of a topological space  $X$  is called locally closed set (briefly  $lc$  set) if  $A = U \cap V$ , where  $U$  is open and  $V$  is closed in  $(X, \tau)$ .

**Definition 2.3.** A map  $f : X \rightarrow Y$  is called,

- (a)  $LC$ -continuous (6)  $f^{-1}(V)$  is locally closed set in  $X$  for each open set  $V$  in  $Y$ .
- (b)  $LC$ -irresolute (6) if  $f^{-1}(V)$  is locally closed set in  $X$  for each locally closed set  $V$  in  $Y$ .

### 3 Regular locally closed sets ( $RL$ -closed sets)

**Definition 3.1.** A subset  $A$  of a topological space  $X$  is called a regular locally closed set ( $RL$ -closed set) if  $A = U \cap V$ , where  $U$  is a regular open set and  $V$  is a closed set in  $(X, \tau)$ . The collection of all  $RL$ -closed sets of  $X$  is denoted by  $RLC(X)$ .

**Example 3.1.** Let  $X = \{a, b, c, d\}$  and  $\tau = \{X, \emptyset, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$ . Then, the closed sets are  $X, \emptyset, \{d\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}$  and  $RLC(X) = \{X, \emptyset, \{a\}, \{b\}, \{d\}, \{c, d\}, \{a, c, d\}, \{b, c, d\}\}$ .

*Remark 3.1.* It is clear that every  $RL$ -closed set is a closed set.

**Lemma 3.2.** Every  $RL$ -closed set is a locally closed set.

*Proof.* Let  $A$  be a  $RL$ -closed set in a topological space  $(X, \tau)$ . Then,  $A = U \cap V$ , where  $U$  is regular open set and  $V$  is closed set in  $(X, \tau)$ . As every regular open set is open, we have  $A$  is a locally closed set.  $\square$

**Proposition 3.1.** If both  $A$  and  $B$  are  $RL$ -closed sets in a topological space  $(X, \tau)$ , then  $A \cap B$  is  $RL$ -closed set.

*Proof.* Let  $A$  and  $B$  be two  $RL$ -closed sets in a topological space  $X$ . Then,  $A = U_1 \cap V_1$  and  $B = U_2 \cap V_2$ , where  $U_1, U_2$  are regular open sets and  $V_1, V_2$  are closed sets. Now,  $A \cap B = U_3 \cap V_3$ , where  $U_3 = U_1 \cap U_2$  and  $V_3 = V_1 \cap V_2$ . Since the intersection of any two regular open sets is regular open set, so we get  $A \cap B$  is a  $RL$ -closed set.  $\square$

*Remark 3.2.* If both  $A$  and  $B$  are  $RL$ -closed sets in a topological space  $X$ , then in general,  $A \cup B$  need not be a  $RL$ -closed set. This can be seen from the following example:

**Example 3.3.** Let  $X = \{a, b, c\}$  and  $\tau = \{X, \emptyset, \{a\}, \{b\}, \{a, b\}\}$ . Then,  $RLC(X) = \{X, \emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$ . Let  $A = \{a\}$  and  $B = \{b\}$  be  $RL$ -closed sets in  $X$ . Then,  $A \cup B = \{a, b\}$  is not a  $RL$ -closed set in  $X$ .

The regular locally open sets are defined to be the complement of the regular locally closed sets.

**Definition 3.2.** Let  $X$  be a topological space and  $A$  be a subset of  $X$ . Then,  $A$  is called regular locally open set (briefly  $RL$ -open set) if  $A = U \cup V$ , where  $U$  is a regular closed set and  $V$  is an open set in  $X$ . The collection of all regular locally open sets of  $X$  is denoted by  $RLO(X)$ .

*Remark 3.3.* It is clear that every  $RL$ -open set is an open set.

**Definition 3.3.** Let  $X$  be a topological space and let  $A \subseteq X$ . The union of all  $RL$ -open sets contained in  $A$  is called regular locally interior of  $A$  and is denoted by  $\text{int}_{RL}(A)$ .

---

**Definition 3.4.** Let  $X$  be a topological space and let  $A \subseteq X$ . The intersection of all  $RL$ -closed supersets of  $A$  is called regular locally closure of  $A$  and is denoted by  $cl_{RL}(A)$ . That is,  $cl_{RL}(A) = \bigcap_{i \in I} F_i$  where  $A \subseteq F_i$ ,  $F_i$  is  $RL$ -closed set for each  $i$ .

**Proposition 3.2.** If both  $A$  and  $B$  are  $RL$ -open sets, then  $A \cup B$  is a  $RL$ -open set in a topological space  $X$ .

*Proof.* Let  $A$  and  $B$  be two  $RL$ -open sets in a topological space  $(X, \tau)$ . Then,  $A = U_1 \cup V_1$  and  $B = U_2 \cup V_2$ , where  $U_1, U_2$  are regular closed sets and  $V_1, V_2$  are open sets. Now  $A \cup B = U_3 \cup V_3$ , where  $U_3 = U_1 \cup U_2$  and  $V_3 = V_1 \cup V_2$ . Since the finite union of regular closed sets is a regular closed set, we get  $A \cup B$  is a  $RL$ -open set in  $X$ .  $\square$

**Definition 3.5.** A mapping  $f : X \rightarrow Y$  is called a regular locally closed continuous map (briefly  $RLC$ -continuous map) if  $f^{-1}(V)$  is a  $RL$ -closed set in  $X$  for every closed set  $V$  in  $Y$ .

**Example 3.4.** let  $X = Y = \{a, b, c, d\}$ ,  $\tau = \{X, \emptyset, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$  and  $\sigma = \{Y, \emptyset, \{a, b\}, \{a, b, c\}, \{a, b, d\}\}$ . Define  $f : (X, \tau) \rightarrow (Y, \sigma)$  by  $f(a) = c, f(b) = b, f(c) = f(d) = d$ . Then,  $f$  is a  $RLC$ -continuous map.

**Theorem 3.5.** A mapping  $f : X \rightarrow Y$  is called a  $RLC$ -continuous map if and only if the inverse image of every open set in  $Y$  is a  $RL$ -open set in  $X$ .

*Proof.* Suppose that  $f : X \rightarrow Y$  is a  $RLC$ -continuous map. Let  $U$  be an open set in  $Y$ . Then,  $Y - U$  is closed in  $Y$ . Since  $f$  is  $RLC$ -continuous,  $f^{-1}(Y - U) = X - f^{-1}(U)$  is a  $RL$ -closed set in  $X$ . Hence  $f^{-1}(U)$  is a  $RL$ -open set in  $X$ .

Conversely, suppose that  $f^{-1}(U)$  is a  $RL$ -open set in  $X$  for every open set  $U$  in  $Y$ . Let  $V$  be a closed set in  $Y$ . Then,  $Y - V = U$  is an open set in  $Y$  so  $f^{-1}(U) = X - f^{-1}(V)$  is a  $RL$ -open set in  $X$ . That is,  $f^{-1}(V)$  is a  $RL$ -closed set in  $(X, \tau)$ . Hence  $f$  is a  $RLC$ -continuous map.  $\square$

**Definition 3.6.** A mapping  $f : X \rightarrow Y$  is called regular locally closed irresolute map (briefly  $RLC$ -irresolute map) if  $f^{-1}(V)$  is a  $RL$ -closed set in  $X$  for each  $RL$ -closed set  $V$  in  $Y$ .

**Example 3.6.** Let  $X = \{a, b, c\}$  and  $\tau = \{X, \emptyset, \{a\}, \{c\}, \{a, c\}\}$  and let  $Y = \{a, b, c, d\}$  and  $\sigma = \{Y, \emptyset, \{a\}, \{b\}, \{a, b\}, \{a, b, c\}\}$ . Define  $f : X \rightarrow Y$  by  $f(a) = b, f(b) = c$  and  $f(c) = a$ . Then,  $f$  is a  $RLC$ -irresolute map.

**Theorem 3.7.** A mapping  $f : X \rightarrow Y$  is  $RLC$ -irresolute if and only if the inverse image of every  $RL$ -open set in  $Y$  is a  $RL$ -open set in  $X$ .

*Proof.* Suppose that  $f : X \rightarrow Y$  is  $RLC$ -irresolute map. Let  $U$  be a  $RL$ -open set in  $Y$ . Then,  $Y - U$  is  $RL$ -closed set in  $Y$  so  $f^{-1}(Y - U) = X - f^{-1}(U)$  is a  $RL$ -closed set in  $X$ . Hence  $f^{-1}(U)$  is a  $RL$ -open set in  $X$ .

Conversely, suppose that  $f^{-1}(U)$  is a  $RL$ -open set in  $X$  for every  $RL$ -open set  $U$  in  $Y$ . Let  $V$  be a  $RL$ -closed set in  $Y$ . Then,  $(Y - V) = U$  is a  $RL$ -open set in  $Y$  so  $f^{-1}(U) = X - f^{-1}(V)$  is a  $RL$ -open set in  $X$ . That is,  $f^{-1}(V)$  is a  $RL$ -closed set in  $X$ . Hence  $f$  is a  $RLC$ -irresolute map.  $\square$

## 4 Regular Locally Connected Spaces

**Definition 4.1.** Let  $A$  and  $B$  be subsets of a topological space  $X$ . Then,  $A$  and  $B$  are called,  $RL$ -separated if  $A \cap cl_{RL}(B) = \emptyset = cl_{RL}(A) \cap B$ .

---

**Definition 4.2.** A topological space  $X$  is said to be regular locally connected (briefly  $RL$ -connected) if  $X$  cannot be written as the union of two non-empty disjoint  $RL$ -open sets.

**Example 4.1.** Let  $X = \{a, b, c\}$  and  $\tau = \{X, \phi, \{a\}\}$ . Then, the topological space  $(X, \tau)$  is  $RL$ -connected.

**Theorem 4.2.** The following statements are equivalent for a topological space  $X$ :

- (a)  $X$  is  $RL$ -connected,
- (b) the only subsets of  $X$  which are both  $RL$ -open and  $RL$ -closed are  $X$  and the empty set,
- (c)  $X$  cannot be expressed as the union of two disjoint non-empty  $RL$ -open sets,
- (d) there is no  $RLC$ -continuous function from  $X$  onto a discrete two-point space  $\{a, b\}$ .

*Proof.* First we prove (a)  $\Rightarrow$  (b). Suppose  $X$  is  $RL$ -connected and let  $A$  be a non-empty  $RL$ -open and  $RL$ -closed subset of  $X$ . Then,  $X - A$  is both  $RL$ -open and  $RL$ -closed. Since  $X$  is the disjoint union of  $RL$ -open set  $A$  and  $X - A$ , one of these must be empty. That is,  $A = \emptyset$  or  $X - A = \emptyset$ . Now we prove (b)  $\Rightarrow$  (c). Suppose that the only subsets of  $X$  which are both  $RL$ -open and  $RL$ -closed are  $X$  and the empty set. Let  $X = A \cup B$ , where  $A$  and  $B$  are two disjoint non-empty  $RL$ -open sets.  $B = X - A$ , which is a  $RL$ -closed set. But this means that  $B$  is both  $RL$ -open and  $RL$ -closed which contradicts our hypothesis. Therefore,  $X$  cannot be expressed as the union of two disjoint non-empty  $RL$ -open sets. We prove (c)  $\Rightarrow$  (d). Suppose that  $X$  cannot be expressed as the union of two disjoint non-empty  $RL$ -open sets. Let  $Y$  be a discrete space with more than one point and let  $f : X \rightarrow Y$  be an onto  $RL$ -continuous function. Define the non-empty  $RL$ -open sets  $U, V$  such that  $Y = U \cup V$ . Since  $f$  is  $RLC$ -continuous,  $X = f^{-1}(U) \cup f^{-1}(V)$ . But this is a contradiction to the hypothesis. Therefore,  $Y$  can't be a discrete space with more than one point. That is, there is no  $RLC$ -continuous function from  $X$  onto a discrete two-point space  $\{a, b\}$ . Finally, we prove (d)  $\Rightarrow$  (a). If  $f : X \rightarrow Y$  be an onto  $RLC$ -continuous, then  $f^{-1}(a), f^{-1}(b)$  are disjoint open subsets of  $X$  whose union is  $X$  and we have both  $f^{-1}(a), f^{-1}(b)$  are non-empty. Then,  $X$  is not  $RL$ -connected.  $\square$

**Theorem 4.3.** If  $f : X \rightarrow Y$  is a  $RLC$ -continuous surjective map and  $X$  is  $RL$ -connected, then  $Y$  is connected.

*Proof.* Suppose that  $X$  is  $RL$ -connected and assume that  $Y$  is not connected. Then,  $Y = A \cup B$ , where  $A$  and  $B$  are non-empty disjoint open sets in  $Y$ . Since  $f$  is a  $RLC$ -continuous surjective map,  $X = f^{-1}(A) \cup f^{-1}(B)$ , where  $f^{-1}(A)$  and  $f^{-1}(B)$  are non-empty disjoint  $RL$ -open sets. This is a contradiction to that  $X$  is  $RL$ -connected. Hence  $Y$  is connected.  $\square$

**Theorem 4.4.** If  $f : X \rightarrow Y$  is a  $RLC$ -irresolute surjective map and  $X$  is  $RL$ -connected, then  $Y$  is  $RL$ -connected.

*Proof.* Suppose that  $X$  is  $RL$ -connected and assume that  $Y$  is not  $RL$ -connected. Then,  $Y = A \cup B$ , where  $A$  and  $B$  are non-empty disjoint  $RL$ -open sets in  $Y$ . Since  $f$  is a  $RLC$ -irresolute surjective map,  $X = f^{-1}(A) \cup f^{-1}(B)$ , where  $f^{-1}(A)$  and  $f^{-1}(B)$  are non-empty disjoint  $RL$ -open sets. This is a contradiction to that  $X$  is  $RL$ -connected. Hence  $Y$  is  $RL$ -connected.  $\square$

**Definition 4.3.** A subset  $Y$  of a topological space  $X$  is called the  $RL$ -subspace of  $X$  if  $Y \cap U$  is  $RL$ -open, when  $U$  is  $RL$ -open in  $X$ .

**Definition 4.4.** A  $RL$ -subspace  $Y$  of a topological space  $X$  is  $RL$ -disconnected if there exist  $RL$ -open subsets  $U$  and  $V$  of  $X$  such that  $Y \cap U$  and  $Y \cap V$  are disjoint non-empty  $RL$ -open sets whose union is  $Y$ . The  $RL$ -subspace is  $RL$ -connected if it is not  $RL$ -disconnected.

**Lemma 4.5.** If  $Y$  is a  $RL$ -connected subspace of  $X$  and if the sets  $U$  and  $V$  form a  $RL$ -separation of  $X$ , then  $Y \subset U$  or  $Y \subset V$ .

---

*Proof.* Since  $U$  and  $V$  are both  $RL$ -open in  $X$ , the sets  $Y \cap U$  and  $Y \cap V$  are  $RL$ -open in  $Y$ . We have,  $(Y \cap U) \cup (Y \cap V) = Y$  and  $(Y \cap U) \cap (Y \cap V) = \emptyset$ . If  $Y \cap U$  and  $Y \cap V$  are non-empty, then  $Y$  is  $RL$ -separated. But  $Y$  is  $RL$ -connected. Then  $Y \cap U = \emptyset$  or  $Y \cap V = \emptyset$ . Therefore,  $Y \subset U$  or  $Y \subset V$   $\square$

**Theorem 4.6.** *Let  $A$  and  $B$  be subspaces of a topological space  $X$ . If  $A$  and  $B$  are  $RL$ -connected and not  $RL$ -separated, then  $A \cup B$  is  $RL$ -connected.*

*Proof.* Assume that  $A \cup B$  is not  $RL$ -connected. Then,  $A \cup B = U \cup V$ , where  $U$  and  $V$  are disjoint non-empty  $RL$ -open sets in  $X$ . Since  $A$  and  $B$  are  $RL$ -connected, then by Lemma 4.6, either  $A \subset U$  or  $A \subset V$  and  $B \subset U$  or  $B \subset V$ . If  $A \subset U$  and  $B \subset U$ , then  $A \cup B \subset U$  and  $V = \emptyset$ . This is a contradiction to that  $V$  is non-empty. Therefore,  $A \cup B$  is  $RL$ -connected.  $\square$

**Theorem 4.7.** *If  $\{A_\alpha : \alpha \in I\}$  is non-empty collection of  $RL$ -connected subspaces of a topological space  $X$  such that  $\bigcap_{\alpha \in I} A_\alpha \neq \emptyset$ , then  $\bigcup_{\alpha \in I} A_\alpha$  is  $RL$ -connected.*

*Proof.* Assume that  $Y = \bigcup_{\alpha \in I} A_\alpha$  is not  $RL$ -connected. Then  $Y = U \cup V$ , where  $U$  and  $V$  are non-empty disjoint  $RL$ -open sets in  $X$ . Since  $\bigcap_{\alpha \in I} A_\alpha \neq \emptyset$ , there is a point  $p$  of  $\bigcap_{\alpha \in I} A_\alpha$ . Since  $p \in Y$ , either  $p \in U$  or  $p \in V$ . Suppose that  $p \in U$ . Since  $A_\alpha$  is  $RL$ -connected,  $A_\alpha \subset U$  or  $A_\alpha \subset V$ . Since  $p \in A_\alpha$ ,  $A_\alpha \not\subset V$ . Hence,  $A_\alpha \subset U$  for every  $\alpha$ . Then  $Y = \bigcup_{\alpha \in I} A_\alpha \subset U$ . This is a contradiction to that  $V$  is non-empty. Therefore,  $\bigcup_{\alpha \in I} A_\alpha$  is  $RL$ -connected.  $\square$

**Theorem 4.8.** *Let  $A$  be a  $RL$ -connected subspace of  $X$ . If  $A \subset B \subset cl_{RL}(A)$ , then  $B$  is also  $RL$ -connected.*

*Proof.* Assume that  $B$  is not  $RL$ -connected. Then,  $B = U \cup V$ , where  $U$  and  $V$  are disjoint non-empty  $RL$ -open sets in  $B$ . Since  $A$  is  $RL$ -connected, then by Lemma 4.6, either  $A \subset U$  or  $A \subset V$ . Suppose that  $A \subset U$ . Then  $cl_{RL}(A) \subset cl_{RL}(U)$ . Since  $cl_{RL}(U)$  and  $V$  are disjoint,  $B$  cannot intersect  $V$ . This contradicts the fact that  $V$  is a non-empty subset of  $B$ . Therefore,  $B$  is  $RL$ -connected.  $\square$

**Corollary 4.9.** *Let  $\{A_\alpha : \alpha \in I\}$  be a non-empty collection of  $RL$ -connected subspaces of a topological space  $X$  and  $A$  be a  $RL$ -connected subspace of  $X$ . If  $A \cap A_\alpha \neq \emptyset$  for all  $\alpha$ , then  $A \cup (\bigcup_{\alpha \in I} A_\alpha)$  is  $RL$ -connected.*

*Proof.* By Theorem 4.7, each set  $A \cap A_\alpha$ ,  $\alpha \in I$  is  $RL$ -connected and  $\bigcap_{\alpha \in I} (A \cap A_\alpha) \neq \emptyset$  since it contains  $A$ . Thus,  $A \cup (\bigcup_{\alpha \in I} A_\alpha) = \bigcup_{\alpha \in I} (A \cup A_\alpha)$  is  $RL$ -connected.  $\square$

## 5 Regular Locally Compact Spaces

**Definition 5.1.** A collection  $\{G_i : i \in I\}$  of  $RL$ -open sets of  $X$  is said to be  $RL$ -open cover for the space  $X$  if  $X = \bigcup_{i \in I} G_i$ .

---

**Definition 5.2.** A collection  $\{G_i : i \in I\}$  of *RL*-open sets is said to be *RL*-open cover for a subset  $A$  of the space  $X$  if  $A \subseteq \bigcup_{i \in I} G_i$ .

**Definition 5.3.** A topological space  $X$  is said to be regular locally compact space (briefly *RL*-compact space) if for every *RL*-open cover of  $X$  has a finite subcover.

**Definition 5.4.** A subset  $A$  of a topological space is said to be regular locally compact set (briefly *RL*-compact set) if for every *RL*-open cover of  $A$  has a finite subcover.

**Theorem 5.1.** Every *RL*-closed subset of a *RL*-compact space is a *RL*-compact space.

*Proof.* Let  $A$  be a *RL*-closed set of the *RL*-compact space  $X$  and let  $\mathcal{A} = \{G_i : i \in I\}$  be a covering of  $A$  by *RL*-open sets in  $X$ . Let  $\mathcal{B}$  be a *RL*-open cover of  $X$ .  $\mathcal{B} = \mathcal{A} \cup \{X - A\}$ . Since  $X$  is *RL*-compact,  $\mathcal{B}$  has a finite subcover  $\mathcal{B}_{finite}$  of  $X$ . If  $\mathcal{B}_{finite}$  contains the set  $X - A$ , discard  $X - A$ . Otherwise, leave  $\mathcal{B}_{finite}$  alone. Then  $\mathcal{B}_{finite}$  is a finite sub collection of  $\mathcal{A}$  that covers  $A$ .  $\square$

**Theorem 5.2.** The image of a *RL*-compact space under a *RLC*-continuous map is compact.

*Proof.* Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be *RLC*-continuous map from  $X$  onto  $Y$ . Let  $\{G_i : i \in I\}$  be an open cover for  $Y$ . Then,  $\{f^{-1}(G_i) : i \in I\}$  is a *RL*-open cover for  $X$ . Since  $X$  is *RL*-compact, this *RL*-open cover has a finite subcover  $\{f^{-1}(G_1), f^{-1}(G_2) \dots f^{-1}(G_n)\}$ . Since  $f$  is onto,  $\{G_1, G_2 \dots G_n\}$  is the finite open cover for  $Y$ . Therefore,  $Y$  is compact.  $\square$

**Theorem 5.3.** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a *RLC*-irresolute map and  $A \subseteq X$  be a *RL*-compact relative to  $X$ , then the image  $f(A)$  is *RL*-compact relative to  $Y$ .

*Proof.* Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a *RLC*-irresolute map from topological spaces  $X$  onto  $Y$ . Let  $\{G_i : i \in I\}$  be *RL*-open cover of  $f(A)$  relative to  $Y$ . Then,  $\{f^{-1}(G_i) : i \in I\}$  is *RL*-open cover for  $A$  relative to  $X$ . Since  $A$  is *RL*-compact relative to  $X$ , this *RL*-open cover has a finite subcover  $\{f^{-1}(G_1), f^{-1}(G_2) \dots f^{-1}(G_n)\}$ . Since  $f$  is onto,  $\{G_1, G_2 \dots G_n\}$  is a finite *RL*-open cover for  $f(A)$ . Therefore,  $f(A)$  is *RL*-compact.  $\square$

## 6 Conclusions

In this paper, we defined a new class of locally closed sets namely, *RL*-closed sets and investigated some of their properties using the *RL*-closed sets. The *RLC*-continuous maps and *RLC*-irresolute maps were also defined and investigated some of their properties. Finally, the concept of *RL*-connectedness and *RL*-compactness were introduced and their properties were established.

---

## References

- [1] Aho, T. and Nieminen, T. (1994). Spaces in which preopen subsets are semi-open. *Ricerche Mat.*, 43, 55-59.
- [2] Popa, V. and Noiri, T. (1994). Weakly  $\beta$ -continuous functions. *An. Univ. Timisoara Ser. Mat. Inform.*, 32, 83-92.
- [3] Popa, V. (1987). Properties of  $H$ -almost continuous functions. *Bull. Math. Soc. Sci. Math. R. S. Roumanie(N. S)*, 31(79), 163-168.
- [4] Pipitone, V. and Russo, G. (1975). Spazi semiconnessi e spazi semiaperti. *Rend. Circ. Mat. Palermo*, (2)24, 273-285.
- [5] Stone, M. (1937). Applications of the theory of Boolean rings to general topology. *Trans. Amer. Math. Soc.*, 41, 375-381.
- [6] Ganster, M. and Reilly, I. L. (1989). Locally closed sets and  $LC$ -continuous functions. *Internat. J. Math. Sci.*, 12, 417-424.