

## **Evaluation of the morphological quality of hybrid seeds and the vegetative development of plants from the two reproductive systems of elite *Duragenitors* from oil palm (*Elaeis guineensis* Jacq.) Seed fields in Côte d'Ivoire**

### **Abstract**

The production of quality oil palm plant material (*Elaeis guineensis* Jacq.) necessarily involves the creation of seed fields made up of elite *Dura* parents. Two methods of reproduction of these elite *Duragenitors* were adopted for the establishment of current seed fields by the National Center for Agronomic Research (CNRA). The first group is made up of elite first-cycle parents who have undergone two successive generations of [(G1) AF] AF type self-fertilization, noted AFAF. The second group is composed of genitors having undergone a prior self-fertilization followed by AFSIB-type recombination cycle of recombination between genitors of the type (G1 × G2) AF, noted AFSIB. This study consisted in evaluating the morphological quality of seeds from the two breeding systems of the parents and their quality of vegetative development. For this, the seeds from 553 *Dura* parents distributed between eight (8) descendants and grouped within the category C1001 F were used. The results which derive from this study have shown that the seeds obtained from the parents obtained by a double self-fertilization of AFAF-type parents are characterized by seeds with a thin shell and a high percentage of germination. After the elimination of the abnormal plants at the end of the nursery, the measurements carried out (circumference at the crown, height of the plants and the number of leaves emitted from the plants) on the healthy plants showed that the plants resulting from the seeds obtained from self-fertilization followed by AFSIB-type recombination showed good vegetative development and vigor compared to those of AFAF type

Keywords: Oil palm, hybrid seeds, reciprocal recurrent selection, self-fertilization.

### **INTRODUCTION**

To exploit the heterosis existing in the crosses made between certain geographically different origins and the constraints due to inbreeding depression, the IRHO adopted in 1957 a reciprocal recurrent selection scheme inspired by work carried out on maize [2]. This selection scheme exploits two groups of populations (A and B) of different geographical origin and with production components of complementary diets. The two (2) populations are intercrossed to obtain A x B hybrids and are evaluated in the field in descendants tests to look

for General and Specific Aptitudes to the combination. The best crosses are subsequently reproduced from self-fertilization of the parents. A subsequent cycle is initiated from recombination of the parents and introductions are made to enrich the genetic basis of the material.

Overall in 70 years, two (2) complete selection cycles have been carried out and have made it possible to produce plant material for producers.

If the hybrids of the 1st cycle were obtained from self-fertilization of parents, this is not the case for the 2nd round of selection. Hybrid seeds of the 2nd cycle are obtained from two (2) breeding systems of genitors. This is a group using two (2) successive AFAF-type self-fertilization cycles and a second group obtained from self-fertilization followed by recombination of related AFSIB-type parents. These two groups of Dura genitors coexist in female seed fields for oil palm seed production. The Tenera type (D x P) hybrid seeds produced from crosses between these two groups of elite Dura female parents and pisifera male parents are currently distributed as a mixture by the National Center for Agronomic Research (CNRA) and yield 24 tons / ha / year of bunches and 6 tons of palm oil per hectare in plantation in Côte d'Ivoire under the best growing conditions.

[12; 11;10] have shown during their work that *Dura* progenitors from self-fertilization followed by AFSIB-type recombination have a faster height growth rate, a performance more important agronomic and a better quality of the fruit and bunches than those obtained from a double self-fertilization of AFAF type genitors.

It is in the face of this observation that it seems interesting to us to assess the morphological quality of the seeds and the vegetative development of the plants resulting from the two breeding systems of the parents in order to choose the best one to optimize the yield of the plots.

## **MATERIAL AND METHODS**

### **Plant material**

The plant material is composed of five hundred and fifty-three (553) genitors of the *Dura* variety used as female parents in oil palm seed fields in Côte d'Ivoire. *Duragenitors* were obtained either by a double self-fertilization of AFAF-type genitors, or by self-fertilization

followed by recombination between related *Dura* palms of the AFSIB type. All these evaluated parents are divided between eight (8) descendants and grouped within the seed category C1001F. Category C1001 F was chosen for its tolerance to Fusarium wilt, therefore used for replanting. All of these palms have the original parents of DA115D x DA115D (Table 1). Within this category, four (4) AFAF-type descendants and four AFSIB-type descendants were evaluated. The parents used are between 21 and 24 years old

### **Experimental dispositive**

The seed fields from which the evaluated genitors originate were established between 1994 and 1997 at the La Mé research station, Côte d'Ivoire. They were planted according to a system in total randomization, at a density of 143 trees per hectare.

### **Variables evaluated**

#### ***Morphological qualities of Tenera hybrid seeds from categories***

For the determination of the morphological quality of the selected seeds, the seeds of six (6) descendants out of the eight (8) from this category were analyzed. These are seeds from descendants LM19016, LM18805, LM19121, LM18801, LM17114 and LM18783. Seeds from descendants LM19175 and LM18503 were not evaluated here as they were not available during the period because they were not available during the study. Analyzes focused on the constituent parts of the palm seed, in particular those involved in its germination. These are the weight of the seeds, the weight of the shell, the thickness of the shell, the weight of the kernel and the percentage of normal embryos of the seeds.

Within the framework of this study, we randomly selected by descent, five (5) parents. From each of them, one (1) bunch is taken and 20 seeds are evaluated. All these seeds had the same shelf life in the seed preparation service and subjected to the same storage temperature which oscillates between 18 and 24 ° C. In total, one hundred (100) seeds per descendants, therefore 600 dry seeds distributed among the six (6) descendants were treated.

#### **- Seed weight (g)**

In each descendant, the 100 seed samples collected were weighed individually using an electronic balance. The average seed weight is obtained from the ratio of the total seed weight of the descendants to the total number of seeds.

$$P. \text{ seed} = (\text{PR seed}) / (\text{Number of seeds})$$

P seed: Average seed weight (g)

PR seed: Total seed weight

#### **- Shell thickness (mm)**

Once weighed, the 100 seed samples from each descendant are cut crosswise. The almond is then separated from the shell to facilitate thickness measurement. The almond was extracted with a pair of pliers. The thickness of the shell was measured using a digital caliper. Seed thickness was grouped by descendants as before. The shell thickness evaluated in mm was determined from the ratio between the sums of the measurements made on the total number of seeds

Ecoque =  $\Sigma mE / (\text{Number of seeds})$

Ecoque: Shell thickness (mm)

$\Sigma mE$ : Sum of measurements taken

#### **-Hull weight (g)**

After the thickness measurement, all the shells were weighed finally to determine the average shell weight of the seeds of the different descendants. The hulls were weighed using an electronic scale.

The weight of the hull was determined from the following relationship:

$Phull = (\text{PR hull}) / (\text{Number of seeds})$

P. hull: Weight hull (g)

PR hull: Total hull weight (g)

#### **-Kernel weight (g)**

The kernels extracted from each shell were weighed to assess the average kernel weight of the seeds of the different descendants. The weight of the almonds is expressed in grams (g). Weightings were performed using an electronic scale. The average seed kernel weight was determined from the following relationship:

Kernel weight =  $(\text{PR Almond}) / (\text{Total number of kernel})$

Kernel weight = Weight kernel

PR kernel = Total weight kernel

#### **-Percentage of normal embryos (%)**

After weighing the almond, an opening is made slightly towards the bulge of the operculum to locate the embryo. The almond is then slashed perpendicular to the embryo just below the operculum to expose it. The embryo is obtained by extracting it with a mounted pin. Embryos

are arranged in rows of 10 (grouping doubles or triples per nut) in a petri dish with a bottom lined with moistened kraft paper, observed with a magnifying glass. Normal embryos are characterized by swelling and yellowish coloration at one end and homogeneous size while abnormal embryos are entirely white and very heterogeneous in size. The percentage of normal embryos was obtained from the following relationship:

$$\% \text{ EN} = (\text{Total Number Seeds} - (\text{MF} + \text{Ab} + \text{NV})) / (\text{Total Number Seeds}) \times 100$$

% EN = Percentage Normal Embryos

MF = Malformed Embryos

Ab = Embryos absent in the almond

NV = Empty seed (without almond)

### ***Evaluation of the germination percentage of Tenera hybrid seeds***

The seed germination percentage was measured in the eight (8) descendants of the C1001F category. Four (4) descendants from the AFSIB type genitor's reproduction system comprising 41390 seeds and four (4) other descendants from the AFAF type genitors reproduction system comprising 40345 seeds.

The percentage of germination was determined by making the ratio between the number of seeds germinated on the total number of seeds put in germination.

$$\% \text{ Germination} = (\text{Number of germinated seeds}) / (\text{Total number of seeds germinated}) \times 100$$

% Germination: Percentage of germination

### ***Evaluation of the growth characteristics of plants in the nursery from the two descendant's reproduction systems in category C1001F***

For the determination of the growth characteristics of the plants in the nursery of the descendants from the two breeding systems of genitors, the observations were made on the plants from the seeds of the genitors of the C1001F category. The assessment covered 22,981 12 month old plants ready for planting. All these plants come from 8 descendants, 4 of which were obtained from the reproduction system of AFSIB type genitors with 10,529 plants and 4 other descendants from the AFAF type genitor's reproduction system comprising 12,452 plants. The bags containing the plants in the nursery were placed in an equilateral triangle device (60 cm between the rows and 70 cm between the plants on the same row). At the end of the nursery, non-vigorous plants showing symptoms of disease were identified and eliminated in the offspring of each reproductive system of the parents. The study focused on the following variables:

**-Plant height (cm)**

Following removal, the length of all healthy and normal plants (21,484 12 month old plants) was measured from the crown to the end of the longest leaf. The average height of the plants expressed in centimeters (cm) was obtained from the following relationship:

Height = (Sum of measurements) / (Number of plants)

#### **-Circumference at the root collar of the plants (cm)**

The method of measuring the circumference at the collar was to go around the collar with a tape measure so that it grips the tree. The determination of the mean circumference at the crown of the offspring plants was obtained from the following formula:

Average circumference at crown = (Sum of measurements) / (Number of plants)

#### **-Number of leaves emitted from plants**

Finally, counting all the functional leaves of each plant made it possible to determine the number of leaves obtained per descendants from the following relationship:

Number of leaves = (Total number of leaves) / (Number of plants)

#### **Statistical analyzes**

Descriptive analyzes (calculations of means, standard deviations, coefficients of variation, confidence intervals) were carried out to describe the different descendants evaluated.

Levene's test was used to test the assumptions of normality and homogeneity of variance variances required for the use of parametric tests. The effect of the reproductive system was evaluated using Student's t-test to verify whether the mode of reproduction of seed-field genitors impacted seed quality and plant development. All of these analyzes were performed using Statistica 7.1 software.

## **RESULTS**

### ***Comparison of the morphological quality of seeds from the two breeding systems of genitors in category C1001 F***

Statistical analyzes revealed that in the majority of cases, no significant difference exists between the morphological quality of the seeds of the two breeding systems of genitors in category C1001 F. The weight of the seeds, the weight of the hulls, the weight of the seeds.

Almonds and the percentage of normal embryos did not differentiate the seeds of the two (2) reproductive systems in this category. On the other hand, the seeds obtained from the genitors of the AFSIB system gave hulls of thickness significantly wider (2.71 mm) than the thickness of the hulls of the seeds of the genitors derived from reproduction of type AFAF with 2.60 mm ( $P = 0.001$ ). The recorded variability was high in the morphological traits of the seeds in the two reproduction systems at the level of the two categories (Table 2).

***Germination performance of hybrid seeds from the descendants of the two breeding systems of genitors in category C1001F***

The results obtained on the seed germination percentage of the two breeding systems of genitors in category C1001F showed that, the seeds from genitors obtained by a double self-pollination of AFAF type genitors recorded a germination percentage significantly higher than those of descendants derived from self-fertilization followed by recombination of related AFSIB-type genitors ( $P = 0.001$ ). Seeds from AFAF type genitors gave 78.11% germination with a coefficient of variation of 14.80% against 70.67% for seeds from AFSIB type genitors with a coefficient of variation of 29.08%. The recorded variability was low within seeds from AFAF-type reproduction systems than in seeds obtained from AFSIB-type reproduction system (Figure 1).

***Development of seedlings in the nursery from the descendants of the two reproductive systems of genitors in the category C1001 F of oil palm***

Significant differences were observed between crown circumference and the number of leaves emitted from nursery plants from the two breeding systems of genitors ( $P = 0.001$ ). On the other hand, the length of the plants in the nursery did not allow to distinguish the plants produced from the seeds of the two reproductive systems of the parents. At the level of the circumference at the collar, the plants resulting from the reproduction seeds of the AFSIB type genitors gave 12.51 cm against 11.49 cm for the plants resulting from the AFAF type seeds. Regarding the number of leaves emitted, the plants obtained from seeds obtained from reproduction of AFSIB type genitors produced 7.86 leaves against 7.45 leaves for plants obtained from reproduction seeds of AFAF type genitors (Figure 2, 3 and 4).

**DISCUSSION**

Studies carried out on the quality of hybrid seeds from the two (2) reproduction systems of oil palm genitors have shown that in category C1001F, the seeds obtained from genitors originating from double self-fertilization of the AFAF type are characterized by a significantly higher germination percentage than that of the descendants resulting from self-

fertilization followed by AFSIB-type recombination ( $P = 0.001$ ). Seeds with a large shell thickness appear to be characterized by a low germination potential. On the germination of oil palm seeds, a process comprising a seed dehydration phase favored by a significant gas exchange has been developed by the CNRA. Studies by [1] have shown that the percentage of germination of oil palm seeds varies from one origin to another. The work of [5] confirmed these results. This author has shown that certain origins of oil palm give only low germination rates regardless of the quality of the pollen. According to him, the percentage of germination of oil palm seeds is linked to the maternal origin of the parents.

But in our case, within the same category C1001F, the germination percentage of seeds from AFAF type genitors was higher than that of AFSIB type genitor's seeds. The low percentage of seed germination of AFSIB type genitors can be explained by formulating several hypotheses:

-the morphological quality of the seeds. Indeed, the wider thickness of the seed shell from AFSIB type genitors could cause a weak gas exchange which must have prevented a good dehydration of the seeds and a more difficult passage of heat to break the dormancy of the embryo. The effect of heat on improving the germination capacity of seeds has been demonstrated by several authors such as [7; 8; 6]. Particularly for oil palm, [9] has shown for fifty years that heat treatment is essential to break dormancy in seeds.

-the biochemical quality of seeds: in fact, the germination process is always linked to biochemical and metabolic manifestations which allow the embryo to develop into a whole plant [3]. Thus, the low germination percentage of seeds from AFSIB type genitors could be explained by the poor quality of the biochemical constituents of their embryos due to the nutrient deficiency of these genitors high producers of bunch [1].

The germination percentage of seeds from AFSIB type genitors is low, after the elimination of abnormal plants at the end of the nursery, the measurements carried out on healthy plants revealed that the circumference at the neck and the number of leaves emitted are significantly more important in plants derived from seeds obtained from AFSIB type genitors compared to those from AFAF type genitors. This result shows that the plants obtained from the seeds of AFSIB type genitors expressed good vegetative development and better vigor. The good vegetative development of these AFSIB plants could be explained by their good health status [4],

The presence of Tenera trees from the seeds of parents obtained by double self-fertilization of the AFAF type in palm groves could therefore cause a drop in yield due to genetic defects such as disruption of plant vigor (dwarfism) and flower abortion. [4]. On the other hand, the



use of Tenera hybrids derived from the seeds of parents originating from self-fertilization followed by AFSIB-type recombination could generate more homogeneous plantations with palm trees that are more genetically compliant. These healthy trees would express greater production potential.

## **CONCLUSION**

The studies undertaken on the evaluation of the morphological quality of seeds obtained from the two reproductive systems of the parents and their quality of vegetative development revealed that in category C1001 F, the seeds obtained from the parents obtained by a double self-fertilization of parents of type AFAP are characterized by thin-shelled seeds and a high percentage of germination. However, the weight of the seeds, shell, almond and the percentage of normal embryos did not allow differentiation of seeds from genitors obtained from the two brood breeding systems. Also, after the elimination of the abnormal plants at the end of the nursery, the measurements carried out on the healthy plants showed that the plants resulting from the seeds obtained from self-fertilization followed by AFSIB-type recombination presented good vegetative development and good force compared to those of type AFAP, namely: the average circumference at the wider crown, the number of leaves emitted more important with plants of larger sizes. Finally, we can remember that the double self-fertilization of AFAP-type parents does not affect the plants during the first months, that is to say germination and their growth in height, but appears later in production.

## **REFERENCES**

1. Beugré MM, Kouakou TH, KonéM, KouaméKR, Konan KE, KouadioYJ. Effect of the hydrogen peroxide treatments on germination of oil palm (*Elaeis guineensis* Jacq.) seeds. *International Journal of Agricultural and Food Science*. 2011; 1 (3): 58-65

2. Comstock RE, Robinson HF, Harvey PH. A breeding procedure designed to make maximum use of both general and specific combining ability, *Agronomy Journal*. 1949; 41: 360-367.
3. Duval B, Shetty K. The stimulation of phenolics and antioxidant in Pea (*Pisum sativum*) elicited by genetically transformed anise root extract. *Food Biochemical*. 2001 ; 25(16): 361-377.
4. Gascon JP, Noiret JM, Meunier J. Effet de la consanguinité chez *Elaeis guineensis* Jacq. *Oléagineux*. 1969 ;24 (11): 603-607.
5. Henry P. La germination des graines d'*Elaeis*. *Rev. Int. Bot. Appliquée et Agriculture Tropicale*. 1951 ; no 349 (350): 565 - 591 et no 351 (352) : 66 - 77.
6. Herrero C, San Martin R, Bravo F. Effect of heat and ash treatments on germination of *Pinus pinaster* and *Cistus laurifolius*. *Journal of Arid Environment*. 2007; 70 : 540-548.
7. Moussa H, Margolis HA, Dube PA, Odongo J. Factors affecting the germination of doum palm (*Hyphaene thebaica*; Mart.) seeds from the semi-arid zone of Niger, West Africa. *Forest Ecology and Management*. 1998;104: 27-41.
8. Periasamy A, Gopapal K, Soh AC. Productivity improvements in seed processing techniques for commercial oil palm seed production. *The Planter*. 2002; 78: 429-442.
9. Rees AR. High-temperature pre-treatment and the germination of seed of the oil palm *Elaeis guineensis* Jacq. *Annals of Botany*. 1962 ; 26 : 569-581
10. Tano EK, Alla O, Konan JN, Konan KE, Nguetta ASP. Effet du système de reproduction des géniteurs sur les caractéristiques de qualité du régime et du fruit des descendances Dura utilisées en production de semences de palmier à huile (*Eleais Guineensis* Jacq). *Agronomie Africaine*. 2020 ; 32 (1) : 101-112.
11. Tano EK, Konan JN, Alla O, Akanvou R, Nguetta ASP, Konan KE. Etude des performances génétiques des descendances parentales issues de deux systèmes de reproduction de géniteurs utilisés en production de semences sélectionnées de palmier à huile (*Elaeis guineensis* Jacq.). *Int. J. Biol. Chem. Sci.* 2019; 13 (3) : 1800-1816.
12. Tano EK, Allou D, Alla O, Konan KE, Nguetta ASP. 2017. Evaluation of the height of some genitors of seed field *Deli Dura* of Oil Palm (*Eleais Guineensis* Jacq) planted in 1998 and 1999 at La Me Station. *Agricultural Science Research Journal*. 2017; 7(10): 318 – 32

Table 1: Characteristics of *Dura* oil palm genitors studied

Plantation	Category	Original parents	Crosses between descendants of the first generation of categories	Reproductive system	Descendants from the second generation of categories	Number of <i>Dura</i> genitors assessed
1997			( <b>LM2509D</b> ) AFAF	AFAF	LM19175	71
1996			( <b>LM2509D</b> X LM2523D) AF	AFSIB	LM18503	22
1994			( <b>LM3394D</b> ) AFAF	AFAF	LM17114	61
1996	C1001F	DA115D AF	( <b>LM3394D</b> X LM2519D) AF	AFSIB	LM18783	66
1997			( <b>LM2531D</b> ) AFAF	AFAF	LM19016	69
1997			(LM2515D X <b>LM2531D</b> ) AF	AFSIB	LM18805	100
1997			( <b>LM3005D</b> ) AFAF	AFAF	LM19121	79
1997			( <b>LM3394D</b> x <b>LM3005D</b> ) AF	AFSIB	LM18801	85

*AFSIB* = Self-fertilization followed by recombination; *AFAF* = Double self-fertilization

Table 2: Comparison of the morphological quality of the seeds of the two breeding systems of genitors in category C1001 F of oil palm

Catégorie	Reproductive system	P. seeds	CV (%)	<b>E. shell</b>	CV (%)	P. thickness	CV (%)	P. kernel	CV (%)	% EN	CV (%)
C1001F	AFAF	3,13 <sup>a</sup>	40,62	<b>2,60<sup>a</sup></b>	21,24	2,24 <sup>a</sup>	41,98	0,76 <sup>a</sup>	46,90	95,00 <sup>a</sup>	30,46
	AFSIB	3,10 <sup>a</sup>	37,18	<b>2,71<sup>b</sup></b>	20,45	2,29 <sup>a</sup>	34,89	0,72 <sup>a</sup>	38,15	94,33 <sup>a</sup>	22,47

*The means followed by the same letter in the same column are not statistically different at the threshold of 0.05 by Student's t test; Cv = Coefficient of variation expressed as a percentage; P = weight (g); E = thickness (mm); EN = normal embryos (%); AFAF: Double self-fertilization; AFSIB: Self-fertilization followed by recombination*




Figure 1: Percentage of seed germination of the two reproductive systems of genitors of category C1001F of oil palm

*AFAF: Double self-fertilization; AFSIB: Self-fertilization followed by recombination*




Figure 2: Length of oil palm seedlings in the nursery of the two breeding systems of genitors in category C1001F

*AFAF = Double self-fertilization; AFSIB = Self-fertilization followed by recombination*




Figure 3: Crown circumference of oil palm plants in the nursery of the two breeding systems of genitors in category C1001F

*AFAF = Double self-fertilization; AFSIB = Self-fertilization followed by recombination*




Figure 4: Number of leaves emitted from oil palm plants in nurseries of hybrids of the two reproduction systems in category C1001F

*AFAF = Double self-fertilization; AFSIB = Self-fertilization followed by recombination*