

DEVELOPMENT AND VALIDATION OF ANALYTICAL METHOD FOR DETERMINATION OF AN IMPORTANT FLAVONOID IN *MALVA NEGLECTA*.

Abstract:

Development of genuine and dependable analytical methods which profile the marker phytoconstituent in an extract containing a mixture of several component is a challenging task. A simple, rapid, precise, and reliable HPLC method was developed for quantification of quercetin from the methanolic extract of *Malva neglecta*. The estimation was carried out using Phenomenex Gemini-NX-5 μm C18 (2), mobile phase consisting of Methanol: Phosphate buffer pH 3 (70:30); the flow rate of 1 mL/min and ultraviolet detection at 280 nm. The parameters for validation were accuracy, precision, linearity and robustness. The calibration curve was found to be linear in a concentration range of 20–100 $\mu\text{g/ml}$. The correlation coefficient was $r^2=0.9996$. The % Average recovery of quercetin was found to be in the range of 99.82 to 100.52 % which was within the acceptance criterion indicating the accuracy of the method. The results of the robustness study indicated that there is no influence of minor changes. The developed and validated method can be successfully used for the determination of quercetin in *Malva neglecta*; thereby helping in authentication and quality control of this plant

Key words: Development, validation, flavonoid, *Malva neglecta*.

38
39
40
41
42

43 1. INTRODUCTION:

44 *Malva neglecta* which is referred to as Khebaiz or Khobbeiza in Arabic is a plant belonging
45 to the Malvaceae family and is wildy grown in the Northern Border Province, Saudi Arabia.
46 It has been traditionally used for insect bites, bladder infection, burns, inflammation, ulcers,
47 wounds , as astringent, demulcent, diuretic, expectorant and laxative. Some of the
48 phytoconstituents reported in the literature are Quinic acid ,Aconitic acid , Chlorogenic acid ,
49 caffeic acid ,Coumaric acid , Rutin, Hyperoside, Myricetin ,Fisetin ,Coumarin, Quercetin
50 Naringenin , Luteolin ,Kaempferol , Apigenin, Rhamnetin and Chysin[1-5].

51 Flavonoids are an important class naturally occurring compound consisting of a wide
52 spectrum of biological activities ,some of the being anti-oxidative, anti-inflammatory, anti-
53 mutagenic, wound healing and other medicinal properties [6].Quercetin is a flavonoid and
54 review reports its various biological activities like antioxidant, anti-inflammatory,anti-cancer,
55 anti-ulcer, antibacterial , antiviral activity and in allergies [7]. WHO has published guidelines
56 to ensure the reliability and repeatability of research on herbal medicines to specify the
57 identity, purity, strength and manufacturing practices [8] Review reports the method
58 development and validation of quercitin in several other plants [9-12].However there are no
59 reports pertaining to quantification and validation of the phytoconstituents of this plant.
60 Hence a HPLC method has been developed in the present work for quantification and
61 validation of quercitin from methanolic extract of dried leaves of *Malva neglecta*.

62 2.METHODOLOGY:

63 (2a)Collection and extraction: The leaves of *Malva neglecta* was collected from Northern
64 Border Province, Saudi Arabia. Shade dried and powdered. 250 gm of the powder was
65 extracted with methanol. The residue was concentrated, dried and stored in the desiccator
66 until further experiment and analysis.

67 (2b). Determination of flavonoids [13]:

68 The total flavonoids content was estimated spectrophotometrically using Aluminum
69 chloride. The plant extracts in methanol were mixed with 1.5ml of methanol, 1M potassium
70 acetate, 10% aluminum chloride and distilled water. The absorbance of this mixture was
71 measured at 415 nm . Different concentrations of the quercitin i.e. 100 µg/ml,200 µg/ml,300
72 µg/ml,400 µg/ml and 500 µg/ml were taken to obtain a calibration curve. The total flavonoid
73 content was calculated as quercitin equivalent from the calibration curve.

74
75 (2c)Method development and Validation [14,15]:

76
77 Mobile phase preparation:

78 Mobile phase was prepared by mixing 70% of HPLC grade Methanol and 30% of 0.005M
79 Phosphate buffer of pH 3. This solution was filtered using a 0.45 micron Millipore filter
80 paper and was sonicated for 10mins. The total volume of the mobile phase prepared was
81 1500ml.

82 Standard preparation:

83 10mg of quercetin was taken in 10ml volumetric flask and make up the volume to 10 ml with
84 methanol. From this solution; 1ml was pipetted into 10ml volumetric flask and volume was
85 made up to the mark with acetonitrile. This is a working solution. Further different
86 concentration ranging from 20 µg/ml, 40 µg/ml, 60µg/ml, 80µg/ml and 100µg/ml was
87 prepared by transferring required aliquots of solution to 10ml volumetric flask and made the
88 volume up to the mark by methanol. This was sonicated for 8 mins then the solution was
89 filtered using 0.45 micron Millipore filters.

90 Sample preparation for Assay

91 1mg of the extract was weighed and dissolved in 1ml of methanol. This solution was filtered
92 using a 0.45 micron Millipore filter paper and was sonicated for 10mins. 20µl of this solution
93 was injected.

94 (2d)HPLC method development:

95 Quercetin in the sample was analyzed by HPLC technique using the conditions as shown in
96 table 1.
97

98 **Table 1: Optimized Chromatographic conditions:**
99

Parameters	Description
Detector	Shimadzu spd10A uv-vis, Japan
Pump	Shimadzu LC-10ATVP, Japan
Software	Baseline chromatography Data System N2000
Injection valve	7725i Rheodyne 20µl, USA
Column	Phenomenex Gemini-NX-5 µm C18(2) , LC Column 250 x 4.6 mm
Elution Type	Isocratic
Elution A	Methanol
Elution B	Methanol: Phosphate buffer pH 3 (70:30)
Flow Rate	1mL/min
Col. Temp	Ambient
Detection	UV-Vis Abs.-Variable Wave. (UV) at 280nm

100

101 HPLC method validation

102 The developed method was validated according to ICH guidelines. The parameters used for
103 validation were Specificity, Precision, Robustness Accuracy, Linearity, limit of quantitation
104 and detection

105 **Accuracy:** was performed in triplicate for various concentrations of quercetin to determine
106 the accuracy of the proposed method. Amount equivalent to 80%, 100% and 120% of the

107 standard amount was injected into the HPLC system in accordance with the procedure.
108 Accuracy was assessed as the percentage accuracy and mean % recovery

109 **Precision:** Interday and intraday were studied to determine the intermediate precision of the
110 proposed analytical method. Quercetin in the concentration of 60 µg/mL was analyzed for
111 intraday and interday variation. The results were expressed as %RSD (Relative Standard
112 Deviation).

113 **Specificity:** The specificity of the method was ascertained by analyzing the standard drug
114 and extract. Quercetin in the sample was confirmed by comparing the R_f values with that of
115 the standard.

116 **Detection limit:** The Limit of detection LOD and limit of quantitation LOQ values were
117 calculated from the calibration curves as per the protocol

118 **Linearity:** Linearity was established by triplicate injections of solutions containing standard
119 quercetin. The linearity range maintained was 20 µg/ml, 40 µg/ml, 60µg/ml, 80µg/ml and 100
120 µg/ml

121 **Robustness:** Robustness was tested by deliberately introducing small changes during the
122 development of the analytical procedure and examining the effect in a particular aspect of its
123 performance, normally its accuracy. The influence of independent variables on the response value
124 was investigated using this method .The variables used in the study were temperature, pH and
125 flow rate of mobile phase. 60µg/ml concentration is selected for carrying out robustness .The
126 parameters varied is as shown in table 2. The experimental runs were performed in triplicate.
127

128 **Table 2: Parameters varied for robustness:**

129

Sl.No	Parameters Varied	I	II	III
1	pH of the mobile phase	2.9	3.0	3.1
2	Temperature of the column	28 °C	30 °C	33 °C
3	Flow rate	0.98ml/min	1.00ml/min	1.02ml/min

130

131

132 **System suitability :** System suitability was established by injecting six replicate injections into
133 HPLC system as per test procedure. The system suitability parameters like retention time , peak
134 area ,peak height, Resolution , tailing factor, Asymmetry, Theoretical Plates were evaluated from
135 standard Chromatograms obtained, by calculating the % RSD .
136

137 **3.RESULTS AND DISCUSSION:**

138 (3a)The leaves of *Malva neglecta* were collected powdered and extracted with methanol as
139 described in the experimental section


140 3(b)Determination of flavonoid content : The total flavonoid content was evaluated by the
141 Aluminum chloride colorimetric method and was calculated as quercetin equivalent. The
142 amount of total flavonoids in the methanolic extract was found to be 292 µg/ml .
143

Fig 1: Calibration curve for quercetin for estimation of total flavonoids.

144

145

Calibration curve for Quercetin for estimation of total flavonoids



146

147

148

149

150 3(c)HPLC method development and validation

151 An HPLC method was developed and validated for the determination of quercetin in the
152 methanolic extract of *Malva neglecta*. Review reports the development and validation
153 methods for quercetin by various group of researchers in various plants using different
154 chromatographic conditions [16-18]. However there are no reports for method development
155 for quercetin from this plant. The Chromatographic conditions were optimized to provide a
156 good performance. The parameters used for validation of the method were accuracy,
157 linearity, precision and robustness. The best and satisfactory results were obtained using
158 Phenomenex Gemini-NX-5 µm C18(2) , LC Column 250 x 4.6 mm and themobile phase
159 consisting of Methanol: Phosphate buffer pH 3 at a ratio of 70:30 and a flow flow rate of
160 1.0ml/min .The retention time for quercetin was 2.848. The method produced linear
161 responses in the concentration range of 20-100%. LOD and LOQ were found to be
162 0.544µg/mL and 1.3/ µg mL respectively The amount of of quercetin was found to be
163 0.23µg/gm. Thus, the HPLC method was found to be selective. The method was performed
164 and validated for the various parameter as per ICH guidelines. The results of the Method
165 development and validation of quercetin are as shown in the following chromatograms and
166 tables .

167


168 **Fig 2: HPLC chromatogram for quercetin in the sample**

169

170

171

172



173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188

Linearity of the developed method:


The concentration, peak area and retention time for linearity of quercetin and the regression line relating standard concentrations of drug using regression analysis were evaluated. The calibration curves were linear in the studied range and equations of the regression analysis were obtained i.e. $R_2 = 0.9996$ for quercetin. Linearity concentration was in the range of 20/ $\mu\text{g/ml}$ to 100/ $\mu\text{g/ml}$. Good linearity was observed over the above-mentioned range indicating that the method is linear over the concentration range studied table 3 depicts the linearity data for quercetin.

Table 3: Linearity data for quercetin

Concen($\mu\text{g/ml}$)	Conc as % of analyte Target	Peak are (mean of three injections)	Peak Area % RSD
20	20	572676.9587	1.96386303
40	40	1088280.208	0.229418599
60	60	1652236.333	1.72731281
80	80	2155012.167	1.21857
100	100	2749456.833	1.725641898
Equation for regression line $Y=27,226.0656x + 8,307.1370$		Correlation coefficient (r^2) =0.9996	

189
190
191
192

Fig 3: Linearity graph for quercetin



193
194
195
196
197
198
199
200
201

Table 4: Data for system suitability for quercitin

System suitability parameter	Acceptance Criteria	Result	Criteria met/not met
Injection precision for retentiontime(min)	RSD \leq 2%	0.559734318	Met
Injection precision for peak area (n = 6)	RSD \leq 2%	1.085541137	Met
Injection precision for peak height	RSD \leq 2%	0.915565447	Met
Resolution (Rs)	Rs \geq 2.0	4.406	Met
USP tailing factor (T)	T = \leq 2.0	1.165	Met
Asymmetry	K = \leq 2.0	1.336	MET
Theoretical Plates (N)	N \geq 2000	2443.14	Met

202
203
204
205
206
207
208
209

Accuracy:

The accuracy study was performed by adding known amounts of rutin in three replicates. The recovery range for rutin was found to be 99.82 to 100.52 %. The accuracy data is as depicted in table 5

Table 5: Data showing % recovery

210

Sample	Percentage nominal (mean of 3 inj)	Amount of standard(μg)		Recovery (%)
		Spike	Found	
1	80	48	47.9179	99.828901
2	100	60	59.4443	99.073856
3	120	72	72.3801	100.52795

211

212

213

214 Precision of the developed method: Repeatability (Intra-day and intermediate precision
215 data) is as shown in table 6 and 7. The quantity used was $60\mu\text{g/ml}$ The RSD values
216 obtained for intraday & intermediate precision rutin were 0.5597 & 0.3959 respectively
217

217

218

Table 6: Intraday Precision –Data sheet

219

220

Sno.	Peak area	Retention Time
1	1600036.125	4.648
2	1587093.375	4.632
3	1609695.5	4.623
4	1585969.5	4.598
5	1633015.5	4.582
6	1598850.625	4.642
Mean	1602443.438	4.620833333
StDev	17395.1827	0.02586439
%RSD	1.085541137	0.559734318

221

222

223

Table 7 : Intermediate Precision – Data sheet showing the peak areas

225

226

Sl no	Day-1	Day-2	Day-3

1	1571014.25	1615860	1607930.25
2	1569665.75	1613946.25	1596367
3	1568828.625	1615205.5	1606692.5
Mean	1569836.208	1615003.917	1603663.25
StDev	1102.738044	972.6698546	6348.972697
%RSD	0.07024542	0.06022709	0.395904358

227
228
229
230
231
232
233
234
235
236

The robustness of the proposed method was evaluated by deliberately changing the chromatographic conditions such as flow rate, temperature and pH. The results showed that varying the chromatographic conditions had no appreciable effects on the chromatographic parameters.

Table 8: Robustness study of the proposed HPLC method

Parameter	Conditions	Retention Time
Temp	28	4.40
	31	4.43
	33	4.42
pH	2.9	4.40
	3.0	4.42
	3.1	4.41
Flow Rate (mL/min)	0.98	4.97
	1.0	4.50
	1.02	4.06

237
238
239
240
241
242
243
244
245

4. CONCLUSION:

ICH guidelines were used for the development and validation of a new HPLC method for determination of quercetin in *Malva neglecta*. The method was found to be simple, specific, accurate and precise. The linearity of the method was determined from the correlation coefficient and the method was found to be linear and within the range of 20/ µg/ml to 100/ µg/ml. The accuracy of the method was calculated by recovery study the results of the proposed method was found to be accurate as all the parameters were in compliance with the

246 acceptance criteria This method can be adopted for the routine quantification and quality
247 control of quercetin in *Malva neglecta* and traditional drugs and formulations containing
248 quercetin.
249

250
251

252 6. REFERENCES:

- 253 1. *Malva neglecta* – Wikipedia https://en.wikipedia.org/wiki/Malva_neglecta retrived on
254 25.11.2017
- 255 2. Wildlife Database Conserving what matters! - Jabal Moussa. [jabalmoussa.org/wildlife-](http://jabalmoussa.org/wildlife-database?page=6)
256 [database?page=6](http://jabalmoussa.org/wildlife-database?page=6)) retrived on 25.11.2017
- 257 3. Flowers in the Bible: Wood Mallow - Flowers in Israel. [www. flowersinisrael. com/](http://www.flowersinisrael.com/Malvasylvestris_page.htm)
258 [Malvasylvestris _page.htm](http://www.flowersinisrael.com/Malvasylvestris_page.htm)
- 259 4. Nesrin H, Abdulselam E, Elif VO, Hüseyin A, Mehmet B, Mustafa A Y, İsmail Y
260 etal Chemical Profile of *Malva Neglecta* and *Malvella Sherardiana* by LCMS/MS,
261 GC/MS and Their Anticholinesterase, Antimicrobial and Antioxidant Properties With
262 Aflatoxin-Contents Marmara Pharma J. 2017;21(3): 471-484.
- 263 5. Güder A, Korkmaz H. Evaluation of in-vitro Antioxidant Properties of
264 Hydroalcoholic Solution Extracts *Urtica dioica* L., *Malva neglecta* Wallr. and Their
265 Mixture .Iranian J Pharma Res. 2012; 11(3): 913-923.
- 266 6. Karak P. Biological activities of flavonoids: an overview. Inter J Pharma Sci
267 Res. 2019; 10(4):1567-74
- 268 7. David A V, Arulmoli R, Parasuraman S. Overviews of Biological Importance of
269 Quercetin: A Bioactive Flavonoid. Pharmacogn Rev. 2016; 10(20): 84–89.
- 270 8. Rudolf B. Quality criteria and standardization of phyto pharmaceuticals. Drug Info J.
271 1998; 6(32): 101
- 272 9. Sanghavi N, Bhosale SD, Malod Y RP-HPLC method development and validation of
273 Quercetin isolated from the plant *Tridax procumbens*. J Sci Innov Res. 2014; 3(6):
274 594-597.
- 275 10. Pawanpreet K , Singh B Analytical Method Development and Validation of
276 Quercetin: A Review . Inter J Pharma Clin Res. 2019; 11(2): 49-56
- 277 11. Savic IM, Nikolic VD, Nikolic LB, Stankovic M Z. Development and validation of a
278 new RP-HPLC method for determination of quercetin in green tea. J Anal Chem.
279 2013; 68(10): 906–911
- 280 12. Girme AS, Saste GB, Shengule SA, Kunkulol RR and Hingorani LL. Method
281 development, optimization and validation of RP-UFLC method for bioactive
282 flavonoids from *Cassia auriculata* . J Pharmacog Phytochem. 2019; 8(1): 77-81
- 283 13. Marinova D, Ribarova F, Atanasova M. Total phenolics and flavonoids in Bulgarian
284 fruits and vegetables. J Univ ChemTech Metall, 2005; 40: 255–260.
- 285 14. Guideline on Validation of Analytical Procedure-Methodology. International
286 conference on Harmonization, Geneva, Switzerland, 1996
- 287 15. Krishna VB, Naira Nayeem. Method Development and Validation for Quantification
288 of Betaine in *Achyranthes aspera* by HPLC Am. J. PharmTech Res. 2013; 3(4)681-
289 87.
- 290 16. Subramanian G, Meyyanathan SN, Karthik Y, Karunakaranair A, Palanisamy DS.
291 Development and Validation of HPLC Method for the Simultaneous Estimation of
292 Quercetin and Rutin in *Aganosma dichotoma* [Roth] K. Schum. Inter J Pharma
293 Pharmaceu Sci. 2014; 6(2): 606-608.
- 294 17. Shah R, Sharma N, Patel V, Savai J, Saraswathy N. Validated HPLC fingerprint

295 analysis for Simultaneous Determination of Quercetin and Kaempferol in Methanolic
296 Extract of *Tridax procumbens*. Inter J Pharma Bio Res. 2012; 3(4): 166-175.
297 18. Movaliya V, Zaveri MN. HPTLC Method Development and Estimation of Quercetin
298 in the Alcoholic Extract of *Aerva javanica* Root. Adv Res Pharma Bio. 2012; 2:
299 222-228

300

301

302

UNDER PEER REVIEW