

Original Research Article

ANTIMICROBIAL ACTIVITY OF LEMON GRASS ON ENTEROCOCCI

ABSTRACT

Aim :The aim of this study was to investigate antimicrobial activity against Enterococcus bacteria.

Materials and methods : Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) were determined by the Broth Dilution Method. The antibiotic susceptibility test against the test organisms was performed by the Disc Diffusion Method.

Results and discussion :Antimicrobial activity of LG was examined against common Enterococcus bacteria using broth microdilution assay and the disc diffusion method. Zone of inhibition was seen when LG was kept with enterococcus bacteria, this proves to be bacteriostatic.

Conclusion : Lemon Grass oil proved to be bacteriostatic against Enterococcus, further research to be done to implement LemonGrass oil as an antibacterial agent.

Keywords: lemongrass oil, Enterococci, antimicrobial.

Running Title - Antibacterial activity of lemongrass

INTRODUCTION

There are roughly 55 unique sorts of lemongrass species—an enduring grass that flourishes in heat and humidities. Two of the species utilized in oils are *Cymbopogon citratus* (West Indian grass) and *Cymbopogon flexuosus* (East Indian grass). *Cymbopogon flexuosus* is the sort utilized in Cliganic's lemongrass oil. Lemongrass oil is removed through steam refining from the new and half dried lemongrass leaves and stems. The fragrance of lemongrass oil is amazingly solid—a smidgen goes far. It can rapidly overwhelm different oils if not utilized in the perfect sums. The aroma is depicted as new, lemony, verdant, earthy and sweet. The vast majority discover the smell alluring. (Honey bees additionally think that it's appealing on the grounds that the oil impersonates the pheromone created by bumble bees. Utilizing lemongrass oil can urge honey bee settlements to move into a specific spot.

In vitro and in vivo considerations have indicated that lemongrass EO and its volatiles can repress a scope of microbes and growths known to be the major causative creatures for a few stockpiling and food-borne microorganisms(1)(2). Lemongrass EO has been accounted for to impede waste of pastry shop items, cheddar, natural products, and chocolates, among others(3)(3,4). The oil has likewise been demonstrated to be a successful fumigant for putting away food spoilage because of its bioactivity in the fume stage (5). Lemongrass EO was found to essentially decrease settlement improvement against key postharvest microorganisms: *Colletotrichum coccodes*, *Botrytis cinerea*, *Cladosporium herbarum*, *Rhizopus stolonifer*, and *Aspergillus niger* in vitro(6). The oil was found to increase the shelf life of realistic usability and tactile properties of refrigerated mussels and vegetables(7). Reports propose that lemongrass oil is a protected common flavor perplexing, additive, and food deterioration inhibitor fit for decreasing the danger of sicknesses related with tainted items(8).

Citratus is generally utilized in medication for treatment of apprehensive and gastrointestinal disturbances, and as antispasmodic, pain relieving, calming, antipyretic, diuretic and narcotic. Studies on concentrates from *Citratus* leaves have shown oxidant, hostile to microbial and hostile to contagious exercises.

The development of bacterial resistance to most of the presently available antibiotics has necessitated the search for new and efficient antibacterial agents. Our team has extensive knowledge and research experience that has translate into high quality publications(18–29),(30–34). Hence the present study was carried out to find out the antibacterial activity of lemongrass oil against *Enterococcus* bacteria. The aim of this study was to investigate antimicrobial activity against *Enterococcus* bacteria.

MATERIALS AND METHODS

Study setting: Biomedical and Environmental Health Research Lab - microbiology Lab, Saveetha dental College.

Procurement of lemongrass oil

The essential oil of lemongrass (*Cymbopogon citratus*) was collected from a local medical shop in Chennai, India.

Test organisms/Bacterial organisms.

The test organisms used in this study were obtained from the Culture Collections of the Department of Microbiology and Microbial Technology, Saveetha institute of medical and technical sciences-Deemed University. The organism used in this study was *Enterococcus*.

Propagation and maintenance of test organisms

The test organisms were streaked on the Nutrient Agar slants and were incubated overnight at (37 +1) °C. The cultures were kept under refrigerated conditions and were subcultured after every day(9).

Preparation of concentrations of lemongrass oil

The different concentrations (v/v) of lemongrass oil viz 5%, 10%, 15%, 20% were prepared aseptically in sterile conditions(10).

Antibacterial activity

The testing of the bacterial cultures for the inhibitory effect of essential oil of lemon grass for different concentrations (5 %, 10 %, 15%, 20%) were performed by using disc diffusion method as described by Southwell et.al (11). 1 □L of active cell suspension of organisms was spread with the help of sterilized long swabs on the agar surface uniformly. The measured quantity of 5-20 □ L of each concentration was pipetted out with a sterilized pipette and the discs were soaked with various concentrations of LG oil and then placed on the agar plate. The plates were incubated at (37+1) °C for 24-48 hours. The zone of inhibition (mm) was measured with a measured scale after the period of incubation.

RESULTS




Figure 1



Figure 2

Figure 1:- Coating of the nutrient agar plate with enterococcus bacteria

Figure 2:- placing the disc containing lemongrass oil




Figure 3




Figure 4

Figure 3: represents the partition made according to the different concentrations of lemon grass oil.

Figure 4: represents the plates kept inside the incubator for 24 hours.




Figure 5

Figure 5:- represents the different zones of inhibitions for different concentrations of lemongrass oil.

The result of the present study was given in Table 1 which depicts the zone of inhibition for different concentrations for the clinical isolated species of enterococci

Table 1:- representing the different zones of inhibition for different concentrations of lemongrass oil added.

Concentration	DISC 1	DISC 2	DISC 3	Mean Value
5 micro L	11 mm	10 mm	8 mm	9.6 mm
10 micro L	13 mm	13.5 mm	15 mm	13.8 mm
15 micro L	15 mm	14 mm	14 mm	14.3 mm
20 micro L	15 mm	14.5 mm	14 mm	14.5 mm




Figure 6:- graph representing the concentration of lemongrass oil versus zone of inhibition. X axis represents the concentration in μ /ml, Y axis represents the zone of inhibition in micrometers. 20 μ l concentration of lemongrass showed the highest zone of inhibition when exposed to enterococci bacteria.

DISCUSSION

The antibacterial activity was found progressively increasing with the increase in concentration of oil. The maximum effect was found at 20% concentration and minimum effect was observed at 5% concentration of oil (Table 1). In the Broth Dilution Method the test organisms were found to be inhibited by lemongrass oil at very low concentration(12).

After incubating for 24 hours, zones of inhibitions were calculated for each concentration. For 5 μ L of lemongrass oil concentration, 9.6 mm of zone of inhibition was present, For 10 μ L of lemongrass oil concentration, 13.6 mm of zone of inhibition was present, For 15 μ L of lemongrass oil concentration, 14.3 mm of zone of inhibition was present, similarly for 20 μ L of lemongrass oil concentration, 14.5 mm of zone of inhibition was seen (Table 1).

Limitations of the study are: Less number of sample size, Homogeneous population, Restriction of sample to specific species of the organism. Our discoveries may, notwithstanding, be summed up to comparative gatherings, with a great view of alternatives to regular medicines. Future research on lemongrass oil can be done on a wide basis to know the exact components present which are responsible for the antimicrobial activity.

Thus, we conclude that in the present era of emerging multidrug resistance among gram positive and gram negative organisms, lemongrass oil will be helpful in treating such infections.

CONCLUSION

Thus, we conclude that Lemon Grass oil proved to be bacteriostatic against Enterococcus, further research to be done to implement LemonGrass oil as an antibacterial agent, lemongrass oil will be helpful in treating infections in the present era of emerging multidrug resistance among gram positive and gram negative organisms.

REFERENCES

1. Eissa HA, Saad SMM, Abd El-Aleem IM, Ibrahim WA, Abd Elmoniem GM, Helmy AM. EFFECT OF NATURAL CABBAGE and TARO EXTRACTS ON OXIDATIVE ENZYMES ACTIVITY OF FROZEN AND DRIED BANANA PRODUCTS [Internet]. Vol. 34, Journal of Food and Dairy Sciences. 2009. p. 11131–46. Available from: <http://dx.doi.org/10.21608/jfds.2009.115827>
2. Abu-Seif F, Abdel- Fattah SM, h. Abo Sreia Y, Shaban H, Ramadan M.

ANTIFUNGAL PROPERTIES OF SOME MEDICINAL PLANTS AGAINST UNDESIRABLE AND MYCOTOXIN-PRODUCING FUNGI [Internet]. Vol. 34, Journal of Food and Dairy Sciences. 2009. p. 1745–56. Available from: <http://dx.doi.org/10.21608/jfds.2009.112162>

3. الربيعي بعل, عبید هح, صالح ته. Antimicrobial Activity For Crud EWatery Extract Of Seeds Of Citrus Aurantifolia (Lime Fruit) Against Gram Positive And Negative Bacteria In Vitro [Internet]. مجلة كلية المأمون. 2016. p. 404. Available from: <http://dx.doi.org/10.36458/1253-000-027-020>
4. Alam K, Agua T, Maven H, Taie R, Rao KS, Burrows I, et al. Preliminary Screening of Seaweeds, Seagrass and Lemongrass Oil from Papua New Guinea for Antimicrobial and Antifungal Activity [Internet]. Vol. 32, International Journal of Pharmacognosy. 1994. p. 396–9. Available from: <http://dx.doi.org/10.3109/13880209409083022>
5. Von Frisch K. The Dance Language and Orientation of Bees. Belknap Press; 2013. 592 p.
6. Ali-Shtayeh MS, -R Yaghmour RM, Faidi YR, Salem K, Al-Nuri MA. Antimicrobial activity of 20 plants used in folkloric medicine in the Palestinian area [Internet]. Vol. 60, Journal of Ethnopharmacology. 1998. p. 265–71. Available from: [http://dx.doi.org/10.1016/s0378-8741\(97\)00153-0](http://dx.doi.org/10.1016/s0378-8741(97)00153-0)
7. Schaneberg BT, Khan IA. Comparison of Extraction Methods for Marker Compounds in the Essential Oil of Lemon Grass by GC [Internet]. Vol. 50, Journal of Agricultural and Food Chemistry. 2002. p. 1345–9. Available from: <http://dx.doi.org/10.1021/jf011078h>
8. Chao SC, Gary Young D, Oberg CJ. Screening for Inhibitory Activity of Essential Oils on Selected Bacteria, Fungi and Viruses [Internet]. Vol. 12, Journal of Essential Oil Research. 2000. p. 639–49. Available from: <http://dx.doi.org/10.1080/10412905.2000.9712177>
9. CDC. Performance Standards for Antimicrobial Susceptibility Testing. 2017. 250 p.
10. Bell JM, Turnidge JD, Jones RN. Development of a Disk Diffusion Method for Testing *Moraxella catarrhalis* Susceptibility Using Clinical and Laboratory Standards Institute Methods: a SENTRY Antimicrobial Surveillance Program Report [Internet]. Vol. 47, Journal of Clinical Microbiology. 2009. p. 2187–93. Available from: <http://dx.doi.org/10.1128/jcm.00304-09>
11. Southwell IA, Hayes AJ, Markham J, Leach DN. THE SEARCH FOR OPTIMALLY BIOACTIVE AUSTRALIAN TEA TREE OIL [Internet]. Acta Horticulturae. 1993. p. 256–65. Available from: <http://dx.doi.org/10.17660/actahortic.1993.344.30>
12. Pereira RS, Sumita TC, Furlan MR, Jorge AOC, Ueno M. Atividade antibacteriana de óleos essenciais em cepas isoladas de infecção urinária [Internet]. Vol. 38, Revista de Saúde Pública. 2004. p. 326–8. Available from:

<http://dx.doi.org/10.1590/s0034-89102004000200025>

13. Onawunmi GO, Yisak W-A, Ogunlana EO. Antibacterial constituents in the essential oil of *Cymbopogon citratus* (DC.) Stapf [Internet]. Vol. 12, Journal of Ethnopharmacology. 1984. p. 279–86. Available from: [http://dx.doi.org/10.1016/0378-8741\(84\)90057-6](http://dx.doi.org/10.1016/0378-8741(84)90057-6)
14. Cimanga K, Kambu K, Tona L, Apers S, De Bruyne T, Hermans N, et al. Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo [Internet]. Vol. 79, Journal of Ethnopharmacology. 2002. p. 213–20. Available from: [http://dx.doi.org/10.1016/s0378-8741\(01\)00384-1](http://dx.doi.org/10.1016/s0378-8741(01)00384-1)
15. Janssen A, Scheffer J, Svendsen A. Antimicrobial Activity of Essential Oils: A 1976-1986 Literature Review. Aspects of the Test Methods [Internet]. Vol. 53, Planta Medica. 1987. p. 395–8. Available from: <http://dx.doi.org/10.1055/s-2006-962755>
16. Hili P, Evans CS, Veness RG. Antimicrobial action of essential oils : the effect of dimethylsulphoxide on the activity of cinnamon oil [Internet]. Vol. 24, Letters in Applied Microbiology. 1997. p. 269–75. Available from: <http://dx.doi.org/10.1046/j.1472-765x.1997.00073.x>
17. Tortorano AM, Viviani MA, Barchiesi F, Arzeni D, Rigoni AL, Cogliati M, et al. Comparison of Three Methods for Testing Azole Susceptibilities of *Candida albicans* Strains Isolated Sequentially from Oral Cavities of AIDS Patients [Internet]. Vol. 36, Journal of Clinical Microbiology. 1998. p. 1578–83. Available from: <http://dx.doi.org/10.1128/jcm.36.6.1578-1583.1998>
18. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species [Internet]. Vol. 94, Archives of Oral Biology. 2018. p. 93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>
19. Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. J Periodontol. 2019 Dec;90(12):1441–8.
20. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. Hypertens Res. 2020 Feb;43(2):153–4.
21. Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. An insight into the emergence of *Acinetobacter baumannii* as an oro-dental pathogen and its drug resistance gene profile - An in silico approach. Heliyon. 2018 Dec;4(12):e01051.
22. Paramasivam A, Vijayashree Priyadharsini J. Novel insights into m6A modification in circular RNA and implications for immunity. Cell Mol Immunol. 2020 Jun;17(6):668–9.
23. Paramasivam A, Priyadharsini JV, Raghunandhakumar S. Implications of m6A modification in autoimmune disorders. Cell Mol Immunol. 2020 May;17(5):550–1.

24. Girija ASS, Shankar EM, Larsson M. Could SARS-CoV-2-Induced Hyperinflammation Magnify the Severity of Coronavirus Disease (CoViD-19) Leading to Acute Respiratory Distress Syndrome? *Front Immunol.* 2020 May 27;11:1206.
25. Jayaseelan VP, Arumugam P. Exosomal microRNAs as a promising theragnostic tool for essential hypertension. *Hypertens Res.* 2020 Jan;43(1):74–5.
26. Ushanthika T, Smiline Girija AS, Paramasivam A, Priyadharsini JV. An in silico approach towards identification of virulence factors in red complex pathogens targeted by reserpine. *Nat Prod Res.* 2021 Jun;35(11):1893–8.
27. Ramalingam AK, Selvi SGA, Jayaseelan VP. Targeting prolyl tripeptidyl peptidase from *Porphyromonas gingivalis* with the bioactive compounds from *Rosmarinus officinalis*. *Asian Biomed.* 2019 Oct 1;13(5):197–203.
28. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A computational study. *pharmaceutical-sciences* [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1-mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>
29. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with *Murraya koengii* bio-compounds: An in-silico approach. *Acta Virol.* 2020;64(1):93–9.
30. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent.* 2021 May;31(3):436–41.
31. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent.* 2021 Mar;31(2):285–6.
32. Barma MD, Muthupandiyan I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol.* 2021 Jun;126:105132.
33. Teja KV, Ramesh S. Is a filled lateral canal - A sign of superiority? *J Dent Sci.* 2020 Dec;15(4):562–3.
34. Reddy P, Krithikadatta J, Srinivasan V, Raghu S, Velumurugan N. Dental Caries Profile and Associated Risk Factors Among Adolescent School Children in an Urban South-Indian City. *Oral Health Prev Dent.* 2020 Apr 1;18(1):379–86.