

Original Research Article

A COHORT-STUDY TO DETERMINE THE PREVALENCE OF PULMONARY MANIFESTATIONS IN RHEUMATOID ARTHRITIS

ABSTRACT

Background: Rheumatoid arthritis is an ailment that has an effect on the lungs in cases of pleural inflammation, it affects the lower as well as upper lung airways. Other effects of the disease can be seen in interstitial lung problem (parenchyma) and pulmonary vasculature.

Aim: Evaluating the occurrence of pulmonary manifestations in RA patients was the objective of this study.

Methods: An observational method which was cross-section in nature was employed in this study which was held in the Rheumatology and Medicine department, Jinnah Medical and Dental College Karachi Pakistan for one-year duration from June 2020 to June 2021. Eighty subjects were included in this study, and all of them underwent a general examination, their medical history was taken into account. A number of lab tests were performed on the patients, ESR, BUN, CRP, HRCT, ALT, radiological investigations and chest X-rays were included.

Results: In this study 42 percent patients presented with pulmonary manifestations. About 45.70 percent of the patients presented with abnormalities in the spirometry test, 42.85 percent showed HRCT abnormalities. The HRCT scores vary with age, TJC (Tender Joint Count), ESR and SJC. FEV₁, HRCT and FVC showed a negative correlation among each other. However other variables did not show any statistically significant correlation.

Conclusion: RA showed prevalence of pulmonary affection and this can be observed in pulmonary and radiological functions. There was an association of age, pulmonary functions and ESR.

Keywords: Rheumatoid arthritis, Pulmonary manifestation, chest infection

INTRODUCTION

Among the leading autoimmune diseases which cause inflammation, Rheumatoid arthritis is the most prevalent. The main affected areas are the joints but there are extra articular effects of the disease which involve other systems such as the lungs which are most affected by the disease¹⁻². The vasculature, pleura and airways of the lungs are affected by the disease and can cause lung parenchyma³⁻⁴. Evaluating the occurrence of pulmonary ailments in Egyptian patients presenting with RA was the main objective of this study.

METHODOLOGY

An observational method which was cross-section in nature was employed in this study which was held in the Rheumatology and Medicine department, Jinnah Medical and Dental College Karachi Pakistan for one-year duration from June 2020 to June 2021. The study consisted of 70 patients and criteria of American College of Rheumatology / European League Against

Rheumatism (ACR/EULAR) 2010 classification was followed.

The following information was obtained from the patients:

- A detailed relevant data collection was performed including age, Medical history, articular symptoms, smoking history and a complete general examination.
- Lab tests were performed on patients, ESR, Complete Blood Count, ALT, CRP-titer, AST, Creatinine and BUN.
- A thorough Musculoskeletal examination was performed on all patients.
- A latex agglutination test kit was used to find Rheumatoid factor titer (RF).
- ELISA was used for Anti-CCP antibody titer

Radiographic evaluation:

1. An x-ray of the Chest (plain) for both the posterior and anterior view
2. Pulmonary high resolution computed tomography scan (HRCT): It was done using both techniques GE Light Speed Plus MSCT 4 channels set and Toshiba Aquilion MSCT 64 channels set. HRCT films were done in supine position with full inspiration without contrast enhancement.

Spirometric pulmonary function test: Spirometry was performed according to American Thoracic Society guidelines (ATS) with assessment of vital capacity, forced vital capacity (FVC), Forced expiratory volume in 1 second (FEV1), FEV1/FVC ratio.

The study was conducted after obtaining an ethical approval from the Ethical committee. SPSS version 22.0 was used to evaluate all statistical data. Standard deviation, range, mean were the qualitative variables, number was described as (no) & (%) as percentage, to compare all variables a qualitative Chi-square test was employed. We used T-test to independently compare two quantitative variables. To rank the variable as negative or positive the Pearson correlation ranking test was performed. P-value > 0.05 was seen as significant.

RESULTS

Out of the 80 participants, 12 subjects were male and the remaining 68 were female 15 percent and 85 percent respectively. The range of age in patients included was 25-77 years in this study, with mean age of 45 ± 11.1 years. The duration of being ill was 0.8 to 24 years with the mean range of 7.4 ± 6.1 years. 9 patients were smokers that makes 11.25% of the patient number included in the study.

Table-1 shows the Demographic Data of the patients

Males	12	15%
Females	68	85%
Age Range	25-77 Years	
Mean Age	45 ± 11.1 years	
Mean Duration of Illness	0.8 to 24 years	

Arthralgia was found to be the most common element, 73.75%, active arthritis 77.5%. Deformed joints were found in 16.25%, dyspnea 31.25% and cough in 41.25% patients.

Table 2. Abnormal spirometry (n=42) and its percentage

Parameter	Pts with abnormal spirometry (n=42)	Percent
Abnormal spirometric patterns	32	45.70%
Mild restrictive	8	19.05%
Moderate restrictive	10	23.81%
Severe restrictive	5	11.90%

Mild obstructive	3	7.14%
Small airway (early obstructive)	9	21.43%
Mixed	7	16.67%

Methotrexate was the dominant steroid used in 83.75% patients, hydroxychloroquine 71.25% and leflunomide in 46.25% and sulfasalazine in 21.25% patients. Abnormal spirometric results were found in 32(45.71%) patients. Severe airflow restriction was observed in 11.90%, moderate in 23.81% and mild in 19.05% patients. 7.14% patients showed mild obstructive pattern, mixed pattern was observed in 16.67% and early obstruction in 21.88% patients.

The HRCT findings are given in Table-3

Table 3. HRCT findings

HRCT finding	No. of pts (80)	Percent
Abnormal HRCT findings	30	42.85
Ground glass appearance	15	18.75%
Pleural irregularities	13	16.25%
Reticulation	21	26.25%
Honey combing	9	11.25%
Subpleural cyst	4	5.00%
Airspace consolidation	3	3.75%
Nodules	5	6.25%
Emphysema	8	10.00%
Bulla	5	6.25%
Bronchiectasis	15	18.75%
Mosaic perfusion	6	7.50%
Pleural effusion	6	7.50%
Air trapping	7	8.75%

More active arthritis $p=0.004$ and older patients $p=0.005$ showed pulmonary affection, swollen joint count ($p=0.002$), dyspnea ($P<0.001$), cough ($P=0.001$), crepitations ($P=0.002$), chest pain ($P=0.027$) and wheezing ($P=0.022$). RA patients with pulmonary affection by HRCT showed more affection regarding FVC ($P=0.0045$) and FEV1 ($P=0.022$), but there was no significant difference regarding FEV1/FVC ratio ($P=0.641$). (Table 3)

A positive statistical difference between age and score was found in total HRCT score and disease parameters ($r=0.380$, $P=0.002$), TJC showed ($r=0.459$, $P<0.001$), SJC difference was ($r=0.370$, $P=0.001$) while ESR showed ($r=0.252$, $P=0.033$) as a positive correlation. However, HRCT and AST showed negative difference ($r=-0.242$, $P=0.041$). FVC and FEV1 showed noteworthy negative correlation ($r=-0.521$, $P<0.001$), ($r=-0.438$, $P<0.001$) respectively. Table 3 shows other parameter results where no significant effect was observed.

DISCUSSION

The etiology of Rheumatoid arthritis (RA) is unknown and this disease is responsible for inflammation of numerous organs and tissues⁶⁻⁷. This disease is known to target synovial joints primarily by causing symmetric affection in peripheral joints. An extra articular effect of the disease is on the lungs in RA patients. RA affects respiratory system holistically including, vasculature, parenchyma, pleura and airways. RA lung manifests in a progressive

disease of parenchyma of the lungs ILD (Interstitial lung disease), clinically 10 percent RA subjects suffer from this condition and 30 percent never go to clinics. Even if articular disease is not present ILD can occur⁸⁻⁹.

We included 80 patients in our study out of which 12 were male and 68 were female subjects 15 percent and 85 percent, respectively. These patients showed mean age \pm SD 45 ± 11.1 years and the age was between 25-77 years, spanning between 0.8-24yr of progressive disease, mean \pm SD 7.4 ± 6.10 yr. Zurouret *al* had involved 12 males and 63 females 16 and 84 percent, respectively. Age of the subjects ranged between 20 and 84 years and 48 years was the mean age of these patients, duration of the disease ranging between 60 days to 27 years. 11.25 percent patients were smokers in our study compared to Zurouret *al* who had 11 smokers included in his study making up 14.6 percent of the study population¹⁰⁻¹¹. Smoking patient percentage is low due to a small number of male subjects as compared to females due to cultural trends.

37 (46.25%) patients of our study showed symptoms of respiration dyspnea being present in 25 (31.25%), cough 41.25%, 9 (12.8) wheezing and 11 chest pain patients making up 15.7%. When compared to Zurouret *al* 32% showed respiratory ailments where wheezing was least common and dyspnea was most occurring symptom, 13.3% cough patients, chest discomfort in 10.7 %, wheezing in 6.7 % and 25.3% Dyspnea¹²⁻¹³.

Forty-two patients had HRCT that makes up 52.5% and 30 (42.85%) patients had abnormalities, ILD was in 24 (30%) patients. 13 (16.25%) subjects showed pleural irregularities, 4 (5%) subjects had subpleural cysts, ground glass in (15) 18.75% patients, septal lines in 23% and honey comb appearance in 11.25 %, 9 patients. The less observed airway issues included, 8.75% air trapping subjects, mosaic perfusion 7.50%, bulla 6.25% patients, emphysema 10%, 2 patient showed airway consolidation, and 18.75% patients showed bronchiectasis. Pulmonary nodule and pleural effusion were found in 6.25% and 7.50% patients respectively.

Our results are consistent with the study done by Youssef *et al* where RA patients 47% showed ILD as most occurrent in 38.9%. Reticular opacities were found in 8 patients 22.2%, while 4 (11.1%) showed ground-glass opacity while 2 (5.6%) patients presented with honeycombing¹⁴⁻¹⁵.

13 patients presented with bronchial abnormality that is 36.1%, 2 patients with emphysema 5.6%, pleural effusion patients were 5.6% 2, and 4 patients with rheumatoid pulmonary nodules 11.1%.

Cortet *et al* found in his study that abnormal HRCT was present in 80.9% patients out of total 68 patients, most common being bronchitis which was present in 30.5%, nodules in patients 28%, ground glass 17.1%, air trapping in 25% and honeycombing in 2.9 percent, only 1.5% pleural effusion was found in this cohort¹⁶⁻¹⁷. The variable results indicate differences in the definition of the disease or the characteristics, another reason could be differences in methodology. PFTs in 52.5% of the study population were found to be abnormal in our study, they showed an obstructive pattern in small airway and restrictive pattern. This result was consistent with the results of a study undertaken by Farraget *al* where he found that 37% patients showed Abnormal PFT's. The obstructive pattern was detected to be 31.4 percent while obstructive was present in 12.8 percent¹⁸.

In another study conducted by Joshi *et al*, 31.4% patients showed the obstructive PFT outline and 12.8% patients showed obstructive pattern. Conflicting results were found in a study undertaken by Biomdoet *al* where the total abnormality percentage of PFT was 38.5% however, the obstructive pattern was found dominant, which stood at 20.4% compared to 16.9% restrictive pattern, the authors of that study found environmental and climate reasons to be responsible for unusual results¹⁹.

In our study a significant association between HRVT score (P value <0.0001) and FVC and

FEV1 was found which is consistent with Zurouret *al*. However, this result was inconsistent with Bilgicet *al*; substantial correlation was not present. Severity score of HRCT and age were correlated in our study. Zhang *et al* found that age is correlated with RA-ILD. That makes it important to ILD screen elder patients²⁰. No correlation was found by Sakreet *al* regarding age of the patients.

Consistent with Yilmazer *et al* and Bilgicet *al*; No association among disease duration and disease severity was found in our study. Hence, suggesting that the severity of disease is not dependent on the duration²¹. In this regard another study conducted by Joshi *et al* found ILD to be more probable in extended disease period. It is safe to say that duration of disease in his cohort may have been longer.

We could not establish a connection among disease and smoking history. There are other studies who also found same results and couldn't establish relationship between smoking and pulmonary issue, Sakreet *al* and Bilgicet *al* in RA²². However, Zhang *et al* found conflicting results and established smoking as a significant risk factor of ILD.

No significant relation between medication and HRCT can be established in our study, Sakreet *al* agrees with our results in his own findings where no association of medication could be established.

Methotrexate however, was found to be a small risk factor which was attributed to increase in pulmonary abnormality in RA patients in a study conclusion of Conway *et al*. Rheumatoid had no association with severity of the disease and there was no significant difference found in groups for anti CCP. Two studies however found its correlation with RF but they could not establish it with anti CCP by Zhang *et al* and Joshi *et al*, no connotation was found. Skareet *al*, Zrouret *al* however, found it with RF but no evaluation of anti CCP was undertaken in this study²³.

A study conducted by Doyle however, demonstrated that titers of RF and anti-CCP antibodies were advanced in RA-ILD subjects. RF and anti-CCP are found to be noteworthy ILD predictors. Anti-citrullinated peptide antibody positive suggests presence of ILD, and more severe ILD is associated with advanced titers of ACPA. The variance however is thought to be linked with lesser male participants, duration and less smokers.

CONCLUSION

RA affected subjects showed more occurrence than expected and can be a dominant risk of more age, ESR, PFT irregularities.

REFERENCES

1. Vakharia AM, Cohen-Levy WB, Vakharia RM, Sodhi N, Mont MA, Roche MW. Perioperative complications in patients with rheumatoid arthritis following primary total knee arthroplasty: an analysis of 102,898 patients. *The journal of knee surgery*. 2019 Nov;32(11):1075-80.
2. ElSherbiny DA. Frequency and Predictors of Extra-articular Manifestations in Patients with Rheumatoid Arthritis. *The Egyptian Journal of Hospital Medicine*. 2019 Jul 1;76(5):4062-7.
3. Salaffi F, Carotti M, Di Carlo M, Tardella M, Giovagnoni A. High-resolution computed tomography of the lung in patients with rheumatoid arthritis: Prevalence of interstitial lung disease involvement and determinants of abnormalities. *Medicine*. 2019 Sep;98(38).
4. Mena-Vázquez N, Albaladejo LP, Manrique-Arija S, Barco CM, Cano CG, Garnica IU, Fernández-Nebro A. Analysis of clinical-analytical characteristics in patients with rheumatoid arthritis and interstitial lung disease: Case-control study. *Reumatología Clínica (English Edition)*. 2020 Mar 12.
5. Li L, Gao S, Fu Q, Liu R, Zhang Y, Dong X, Li Y, Li M, Zheng Y. A preliminary

- study of lung abnormalities on HRCT in patients of rheumatoid arthritis-associated interstitial lung disease with progressive fibrosis. *Clinical rheumatology*. 2019 Nov 1;38(11):3169-78.
6. Kaeley N, Ahmad S, Pathania M, Kakkar R. Prevalence and patterns of peripheral neuropathy in patients of rheumatoid arthritis. *Journal of family medicine and primary care*. 2019 Jan;8(1):22.
 7. Cho SK, Doyle TJ, Lee H, Jin Y, Tong AY, Ortiz AJ, Sparks JA, Kim SC. Validation of Claims-based Algorithms to Identify Interstitial Lung Disease in Patients with Rheumatoid Arthritis. In *Seminars in Arthritis and Rheumatism 2020* May 20. WB Saunders.
 8. Mira-Avendano I, Abril A, Burger CD, Dellaripa PF, Fischer A, Gotway MB, Lee AS, Lee JS, Matteson EL, Eunhee SY, Ryu JH. Interstitial lung disease and other pulmonary manifestations in connective tissue diseases. In *Mayo Clinic Proceedings 2019* Feb 1 (Vol. 94, No. 2, pp. 309-325). Elsevier.
 9. Kang EH, Jin Y, Desai RJ, Liu J, Sparks JA, Kim SC. Risk of exacerbation of pulmonary comorbidities in patients with rheumatoid arthritis after initiation of abatacept versus TNF inhibitors: a cohort study. In *Seminars in arthritis and rheumatism 2020* Jun 1 (Vol. 50, No. 3, pp. 401-408). WB Saunders.
 10. Farquhar H, Vassallo R, Edwards AL, Matteson EL. Pulmonary complications of rheumatoid arthritis. In *Seminars in respiratory and critical care medicine 2019* Apr (Vol. 40, No. 02, pp. 194-207). Thieme Medical Publishers.
 11. Yu M, Guo Y, Zhang P, Xue J, Yang J, Cai Q, You X, Ma J, Yang D, Jia Y, Wang Y. Increased circulating Wnt5a protein in patients with rheumatoid arthritis-associated interstitial pneumonia (RA-ILD). *Immunobiology*. 2019 Jul 1;224(4):551-9.
 12. Fragoulis GE, Chatzidionysiou K. Lung cancer in rheumatoid arthritis. Is there a need for better risk assessment and screening?. *Clinical Rheumatology*. 2020 Mar;39(3):957-61.
 13. Gupta S, Ferrada MA, Hasni SA. Pulmonary Manifestations of Primary Sjögren's Syndrome: Underlying Immunological Mechanisms, Clinical Presentation, and Management. *Frontiers in Immunology*. 2019;10.
 14. Malakar S, Sharma KN, Rana A, Sharma T. Rheumatoid arthritis 1st time presenting with bilateral pleural effusion-A rare presentation. *Indian Journal of Case Reports*. 2019:404-6.
 15. Kameda H, Fujii T, Nakajima A, Koike R, Sagawa A, Kanbe K, Tomita T, Harigai M, Suzuki Y, Japan College of Rheumatology subcommittee on the guideline for the use of methotrexate in patients with rheumatoid arthritis. *Japan College of Rheumatology guideline for the use of methotrexate in patients with rheumatoid arthritis*. *Modern rheumatology*. 2019 Jan 2;29(1):31-40.
 16. Wang D, Zhang J, Lau J, Wang S, Taneja V, Matteson EL, Vassallo R. Mechanisms of lung disease development in rheumatoid arthritis. *Nature Reviews Rheumatology*. 2019 Oct;15(10):581-96.
 17. Kurata I, Tsuboi H, Terasaki M, Shimizu M, Toko H, Honda F, Ohyama A, Yagishita M, Osada A, Ebe H, Kawaguchi H. Effect of biological disease-modifying anti-rheumatic drugs on airway and interstitial lung disease in patients with rheumatoid arthritis. *Internal Medicine*. 2019:2226-18.
 18. Keerio NH, Shah GA, Afzal T, Khanzada AA, Joyo MR, Ahmed N, Noor SS. Surgical Site Infection Incidence Following Hip Surgery: A Cross-Sectional Study. *JPRI*. 14Aug.2021;33(41A):66-74.
 19. Sparks JA, Doyle TJ, He X, Pan B, Iannaccone C, Frits ML, Dellaripa PF, Rosas IO,

- Lu B, Weinblatt ME, Shadick NA. Incidence and Predictors of Dyspnea on Exertion in a Prospective Cohort of Patients With Rheumatoid Arthritis. *ACR open rheumatology*. 2019 Mar;1(1):4-15.
20. Duarte AC, Porter JC, Leandro MJ. The lung in a cohort of rheumatoid arthritis patients—an overview of different types of involvement and treatment. *Rheumatology*. 2019 Nov 1;58(11):2031-8.
21. Esposito AJ, Chu SG, Madan R, Doyle TJ, Dellaripa PF. Thoracic manifestations of rheumatoid arthritis. *Clinics in chest medicine*. 2019 Sep 1;40(3):545-60.
22. Hussain N, Iqbal M, Aamir N. Retrospective analysis of recurrent hospital admissions due to gout and the associated comorbidities at a tertiary care hospital. *Indo american journal of pharmaceutical sciences*. 2017 Jun 1;4(6):1476-80.
23. Vial G, De Pouilly A, Scoupe L, Pereira B, Daien C, Lukas C, Ruysen-Witrand A, Vergne-Salle P, Richez C, Tournadre A. Factors influencing the choice of biologic therapy following Rituximab in patients with rheumatoid arthritis: A retrospective study using propensity score. *Joint Bone Spine*. 2020 Jan 1;87(1):43-8.

UNDER PEER REVIEW