

# The Effects of Chicken Manure Application Rates on Growth and Yield of Swiss Chard

(*Beta vulgaris* var. *cicla* L.)

## ABSTRACT

Swiss chard (*Beta vulgaris* var. *cicla*) is a leafy vegetable that belongs to the *Chenopodiaceae* family. The leaves are cooked, if still tender they are used in salads. Over the years, Swazis have adopted the use of inorganic ~~fertilisers~~ fertilizers as they are easy to apply and come with recommended application rates. However, their main drawback is that they are environmentally unfriendly especially when washed into rivers, streams and other water bodies. For this cause, the use of animal manures has been promoted. ~~Four~~ Four-week old Swiss chard seedlings were transplanted on the 4<sup>th</sup> of February, 2016 in 1.5 x 1.5 m plots with an inter and intra row spacing of 45 cm and they were irrigated twice a day during the first week and every second day from the second week until the end of the experiment. The experiment was conducted at the Horticulture Department Farm, Faculty of Agriculture and Consumer Sciences, Luyengo Campus of the University of Swaziland to determine the effects of chicken manure application rates on growth, yield and quality of Swiss chard. Four chicken manure application rates (10, 20, 40 and 80 t/ha) and a recommended 900kg/ha, inorganic basal ~~fertiliser~~ fertilizer with a 125kg/ha LAN top dressing fertilizer used as a control. A Randomised Complete Block Design (RCBD) with four replicates was used. The study showed that the application of 80 t/ha of chicken manure improved the growth and yield of Swiss chard. It is recommended that farmers may use 80 t/ha of chicken manure because it gave the best results compared to the other treatments.

**Comment [a1]:** In the previous evaluation, some numerical data were requested but not added. It should be added. The reader wants to see high yield. How much?

23

24

25

## 26 1.0. INTRODUCTION

27 Swiss chard is a leafy vegetable that belongs to the *Chenopodiaceae* family and is scientifically  
28 known as *Beta vulgaris* var. *cicla*. It belongs to the same family as beetroot and mangel-wurzel.

Formatted: Font: Not Italic

29 Unlike beetroot and mangel-wurzel, Swiss chard lacks the large bulbous tap root. It is one of the  
30 most nutritious vegetable crops in the world. Swiss chard may be grown in Eswatini/ Swaziland  
31 all year round, in all the ecological zones.

32 Swiss chard forms part of the several leafy green vegetables that are known as 'greens.' It is a  
33 biennial plant with large dark leaves. The leaves are large, glossy and crispy and can grow up to  
34 37 cm long and 25 cm wide [1]. Stalks of Swiss chard come in a variety of colours depending on  
35 the cultivar, they are usually white, yellow, orange or red [2]. The first records of cultivation  
36 place the origin of Swiss chard in the Mediterranean region particularly Italy and was first  
37 written about by the Greek philosopher Aristotle in 4 B.C [1]. Swiss chard is a short day (SD)  
38 plant with critical day length of 12 hours. It grows best at temperatures ranging from 7 to 24 °C.  
39 Swiss chard can withstand light frosts but an extended exposure to temperatures less than 5°C  
40 induces bolting. In hot weather, the leaves remain small and are of inferior quality [3]. Leaves of  
41 Swiss chard are harvested usually within eight weeks from sowing and once they are in good size  
42 [2]. Harvesting is done continuously so that the leaves do not stay long and lose their colour or  
43 become tough.

44

45 | The use of inorganic ~~fertilisers~~fertilizers has resulted in residual toxicities and degradation of the  
46 | soil structure. These inorganic ~~fertilisers~~fertilizers become an environmental threat to aqua life  
47 | when washed into rivers, streams and other water bodies. They are relatively expensive such that  
48 | not all farmers afford them. As a result, some farmers produce Swiss chard below the expected  
49 | optimum level.

**Comment [a2]:** Despite the fact that the introduction is suggested to be improved in terms of use of other organic fertilizers, no improvement has been made.

50 | The main objective of this study was to improve the production of Swiss chard and to contribute  
51 | towards food security and income generation in Eswatini/Swaziland. The specific objective was  
52 | to determine the optimum level of chicken manure application on growth, yield and quality of  
53 | Swiss chard.

54

55

## 56 | 2.0 MATERIAL AND METHODS

### 57 | 2.1 Experimental site

58 | The experiment was conducted at the Horticulture Department Farm, Faculty of Agriculture and  
59 | Consumer Sciences, Luyengo Campus of the University of Swaziland. The farm is located at  
60 | Luyengo, Manzini region, in the Middleveld agro-ecological zone. Luyengo is located at latitude  
61 | 26°4' S and longitude 31°4' E. The average altitude of this area is 750 m above sea level. The  
62 | mean annual precipitation is 980 mm with most of the rain falling between October and April.  
63 | Drought hazard is about 40%. The average summer temperature is 27°C and the winter  
64 | temperature is about 15°C. The soils of Luyengo are classified under Malkerns series. They are  
65 | ferrasolic or merely a ferralitic soil integrated to fersialitic soils or typical ultisols. The soil in the  
66 | experimental area was a sandy loam [4].

## 67 2.2 Plant Materials

68 ~~Four-Four-week~~ old Swiss chard seedlings were obtained from Greenhouse Seedlings, Ezulwini.  
 69 They were transplanted on the 4<sup>th</sup> of February, 2016 in 1.5 x1.5 m plots with an inter and intra  
 70 row spacing of 45 cm and they were irrigated twice a day during the first week and every second  
 71 day from the second week until the end of the experiment.

## 72 2.3 Experimental Design

73 Four chicken manure application rates (10, 20, 40 and 80 t/ha) and a recommended 900kg/ha,  
 74 inorganic basal ~~fertiliser-fertilizer~~ with a 125kg/ha LAN top dressing ~~fertiliser-fertilizer~~ was used  
 75 as a control (Table 1). A Randomised Complete Block Design (RCBD) with four replicates was  
 76 used. Each plot had four rows and there were four plants in each row which gave a total of 320  
 77 plants used in the experiment.

78 Table 1: Treatment descriptions.

Treatment code	Treatment
1	80 t/ha
2	40 t/ha
3	20 t/ha
4	10 t/ha
5	900 kg of 2:3:2 (22) and 125 kg of LAN (28)

Comment [a3]:

Comment [a4]: What is content of the inorganic fertilizer?

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Font:

79

## 80 2.4. Soil analysis

81 Soil chemical properties were ~~analysed-analyzed~~ at the Soil Chemistry laboratory of the  
 82 University of Swaziland, Luyengo Campus.

## 83 2.5. Manure analysis

84 | Chicken manure chemical properties were ~~analysed-analyzed~~ at the soil Chemistry laboratory of  
85 | the University of Swaziland, Luyengo Campus.

## 86 2.6. Data collection

87 | Data was collected weekly, from the second week after transplanting. Five plants were randomly  
88 | selected in each plot for data recording. Data was collected on the following growth parameters:  
89 | plant height, number of leaves and leaf area while leaf area index was calculated. The fresh mass  
90 | and dry mass of the Swiss chard were measured after harvesting.

## 91 2.7. Growth parameters

### 92 2.7.1. Plant height unit

93 | Five plants were randomly selected per plot and plant height was measured from the base of the  
94 | plant to the leaf apex (tip) ~~using a 30 cm ruler.~~

### 95 2.7.2. Number of leaves unit

96 | The number of leaves per plant was determined by physically counting all the leaves on each  
97 | selected plant. Five plants were selected per plot and it was done on a weekly basis, which was at  
98 | week 3, 4, and 5 after transplanting.

### 99 2.7.3. Leaf area unit

100 | The leaf area of the Swiss chard was determined by multiplying the leaf width and leaf length  
101 | and then multiplying the product by 0.75 (correction factor) ~~[5]. (Edje and Ossom, )~~ It was  
102 | expressed in  $\text{cm}^2$ .

Comment [a5]: not the appropriate literature citing.

### 103 2.7.4. Leaf area index

104 | The leaf area index was determined by dividing the leaf area in  $\text{cm}^2$  by the area occupied by a  
105 | single plant in  $\text{cm}^2$ : ~~[5]. Edje and Ossom~~

106 **2.7.5. Fresh and dry mass unit**

107 This was determined at the end of the cropping season by weighing the harvested leaves per plot.  
 108 Five plants per plot were used to determine the fresh and dry mass in this experiment. The plants  
 109 were randomly selected per plot and their shoot fresh mass was measured using a digital scale  
 110 balance. They were then oven dried at a temperature of 72°C for 72 hours to determine their  
 111 shoot dry mass[5].

112 **2.8. Data analysis**

113 The data collected was subjected to analysis of variance (ANOVA) using MSTAT-C statistical  
 114 package, Version 1.4[6]. Where significant differences were detected mean separation ~~were~~ was  
 115 performed using Duncan's New Multiple Range Test (DNMRT) at 5 % probability level [7].

116 **3.0 RESULTS**

117 **3.1. Soil analysis**

118 Soil chemical properties were ~~analysed~~ analyzed at the Chemistry Laboratory of the University of  
 119 Swaziland, Luyengo Campus. The results of the soil chemical properties are shown in Table 2.

120 Table 2: Soil analysis

Soil parameter	Value
Soil pH	5.8
Phosphorus (mgP/kg)	39.56
Potassium (cmolc/kg)	1.54

121

122 **3.2. Manure analysis**

123 Chemical properties of the chicken manure were analysed at the Chemistry Laboratory of the  
 124 University of Swaziland, Luyengo Campus. The results of the chemical properties of chicken  
 125 manure are shown in Table 3.

126 Table 3: Chicken manure analysis

Manure parameter	Value
pH	7.2
Phosphorus(mgP/kg)	17
Potassium(cmolc/kg)	1 895
Magnesium	Not determined-


127 **3.3. Plant height**

128 ~~Plant~~ The plant height of spinach was significantly ( $P < 0.05$ ) different among the different  
 129 treatments. The highest plant height (34.6 cm) was obtained in spinach treated with 80 t/ha of  
 130 chicken manure while the lowest plant height (26.0 cm) was obtained in Swiss chard plants  
 131 treated with 10 t/ha of chicken manure (Figure 1). The plant height of Swiss chard plants treated  
 132 with inorganic fertilizers was higher (28.1 cm) but not significantly ( $P > 0.05$ ) different from those  
 133 treated with 10 t/ha of chicken manure (26.0 cm) (Figure 1).

Comment [a6]: ???

Comment [a7]: ???

Formatted: Highlight



134

135 Figure 1: Effects of chicken manure on **Swiss chard** plant height. Vertical bars are standard

136 error (se) below and above the mean.

### 137 3.4. Number of leaves


138 The number of leaves per plant ~~were~~ was not significantly ( $P>0.05$ ) different among the **Swiss**

139 **chard** plants. The highest number of leaves (7.6) was obtained in plants treated with 80 t/ha of

140 chicken manure while the lowest number of leaves (6.4) was obtained in plants treated with 40

141 t/ha of chicken manure (Figure 2).





142


143 Figure 2: Effects of chicken manure on number of leaves of **Swiss chard**. Vertical bars are  
 144 standard error (se) below and above the mean.

145

### 146 3.5. Leaf area

147 The leaf area of **Swiss chard** plants was significantly ( $P < 0.05$ ) different among treatments. The  
 148 highest leaf area ( $291.9 \text{ cm}^2$ ) was obtained in plants treated with 80 t/ha of chicken manure while  
 149 the lowest leaf area ( $169.8 \text{ cm}^2$ ) was obtained in plants treated with 10 t/ha of chicken manure at

150 5 WAT (Figure 3). The leaf area of **Swiss chard** increased with increasing application rates of  
 151 chicken manure.



152  
 153 Figure 3: Effects of chicken manure on the leaf area per plant of **Swiss chard**. Vertical bars are  
 154 standard error (se) below and above the mean.

### 155 3.6. Leaf area index


156 The leaf area index (LAI) was significantly ( $P < 0.05$ ) different among treatments. The highest LAI  
 157 (1.1) was obtained in plants treated with 80 t/ha of chicken manure while the lowest LAI

158

159

160 (0.5) was obtained in plants treated with 10 t/ha of chicken manure at 5 WAT (Figure 4). The

161 leaf area of **Swiss chard** increased with increasing application rates of chicken manure.



162

163 Figure 4: Effects of chicken manure on the LAI per plant of **Swiss chard**. Vertical bars are

164 standard error (se) below and above the mean.

165


### 166 3.7. Fresh mass and dry mass

167 There was a significant ( $P < 0.05$ ) difference in the fresh shoot mass of **Swiss chard** plants among

168 treatments (Figure 5). The highest fresh shoot mass (237.5 g) was obtained in plants treated with

169 80 t/ha of chicken manure while the lowest fresh shoot mass (100.0 g) was obtained in plants  
 170 ~~fertilised-fertilized~~ with inorganic ~~fertilisersfertilizers~~. There was no significant difference in  
 171 Swiss chard fresh maas amended with 40t/ha or 80 t/ha chicken manure.

172 There was a significant ( $P < 0.05$ ) difference in the dry shoot mass of Swiss chard plants among  
 173 the different treatments (Figure 5). The highest dry shoot mass (20.4 g) was achieved at 80 t/ha  
 174 of chicken manure while the lowest dry shoot mass (10.1 g) was obtained in Swiss chard plants  
 175 treated with inorganic ~~fertilisersfertilizers~~.



176  
 177 Figure 5: Effects of chicken manure on fresh and dry shoot mass of Swiss chard at week 5 after  
 178 transplanting. Bars followed by the same alphabet are not significantly different from  
 179 one another at  $P = 0.05$ . Mean separation by Duncan's New Multiple Range Test.

180

## 181 4.0. DISCUSSION

182 Different application rates of chicken manure had varying effects on growth, yield and quality of  
 183 **Swiss chard**. Plants treated with 80 t/ha of chicken manure performed better in terms of growth  
 184 in comparison with the other treatments. These **Swiss chard** plants had the highest plant height,  
 185 number of leaves, fresh shoot mass, dry shoot mass, leaf area and leaf area index compared to  
 186 spinach treated with 10, 20, 40 t/ha of **chicken** manure and application of inorganic fertilisers  
 187 recommended for **Swiss chard** production. **Swiss chard** plants treated with 10 t/ha had the lowest  
 188 plant height, leaf area and leaf area index. The highest number of leaves of **Swiss chard** plants  
 189 from the highest application rate of chicken manure must have been as a result of relatively high  
 190 amounts of nitrogen [8]. It was also noted that plant height, number of leaves, leaf area, leaf area  
 191 index, fresh and dry shoot mass increased with increasing levels of chicken manure. These  
 192 results are in agreement with those of [8]-who studied the effects of an organic fertilizer (cattle  
 193 manure) on *Zea mays*. As chicken manure application rate was increased, the availability of plant  
 194 nutrients in the soil also increased. **This resulted in the increase of growth and yield.**

Formatted: Highlight

195 Chicken manure at 80 t/ha performed better in comparison with inorganic fertilizers. These  
 196 findings do not deviate much from those obtained by [9] Owen (2008) who reported that  
 197 synthetic ~~fertilisers-fertilizers~~ do not have good characteristics in aggregating soil particles. The  
 198 plants treated with inorganic ~~fertilisers-fertilizers~~ gave a lower yield than those treated with 80  
 199 t/ha of chicken manure. Animal manures have beneficial effects on the physical and chemical  
 200 properties ~~of soil~~ and therefore have the ability to retain water, supply macro- and trace elements  
 201 absent in inorganic ~~fertilisers-fertilizers~~. Increased vegetable yield with the use of manure ~~have~~  
 202 has been previously reported for okra [10]. The benefits of organic fertilizer use in  
 203 vegetable production ~~has-have~~ previously been reported [11,12,13] and very recently reported  
 204 [14,15,16] in the Kingdom.

Comment [a8]: Which one??

## 205 5.0. CONCLUSION AND RECOMMENDATION

206

207 The study showed that the application of 80 t/ha of chicken manure improved the growth and  
208 yield of **Swiss chard**. From these findings, it can ~~therefore~~, therefore, be concluded that 80 t/ha  
209 was best for [8]-under the conditions of this study. **If fresh mass at the end is the most important**  
210 **parameter farmers could as well use 40 t/ha because there was no significant difference in plants**  
211 **amended with 80 t/ha.**

212 It is recommended that farmers may use 80 t/ha of chicken manure because it gave the best  
213 results compared to the other treatments.

214

## 215 REFERENCES

- 216 1.Pierce LC. Vegetables: Characteristics, Production, and Marketing.John Wiley and Sons.  
217 New York, New York, USA. . 1987.
- 218 2.Hadfield J. Vegetable Gardening in Central Africa.Purnell and Sons. Johannesburg, South  
219 Africa.1960.
- 220 3. Gilber, Z, Hadfield J. Down-to- Earth Fruit and Vegetable Gardening in South Africa. Struik  
221 Publishers. Cape Town, South Africa.1987.
- 222 4.Murdoch G. Soils and Land Capacity in Swaziland. Swaziland Ministry of Agriculture.  
223 Mbabane, Swaziland. 1970.
- 224 5.Edje OT,Ossom EM. **Crop Science Handbook**.Blue Moon Publishers.Manzini,  
225 Swaziland.2009.

- 226 6.Nissen,O. MSTAT-C. A micro Computer Program Design for Management and  
227 Analysis of Agronomic Research Experiments.Michigan State University. East  
228 Lansing, Michigan, USA. 1989.
- 229 7.Gomez KA, Gomez A A. Statistical Procedures for Agricultural Research. John Wiley  
230 and Sons 2<sup>nd</sup> Edition. Singapore.(1984).
- 231 8.Lungu O I,Chirwa B. Effect of lime and farmyard manure on soil acidity from and maize  
232 growth on alfisols from Zambia.Tropical Agric(Trinidad)1993; 70; 309-315.
- 233 9.Owen P. Origin and distribution of lettuce.  
234 (2008).<http://www.calettuceresearchboard.org/Origin.html>. 10/04/2016
- 235 10.Ogunlela VB, Masarirambi MT,Makuza SM. Effects of cattle manure application on  
236 pod yield and yield indices of Okra (*Abelmoschus esculentus* L. Moench) semi-arid and  
237 subtropical environment. J Food, Agric Environ. 2005;3 (1):125-129
- 238 11. Masarirambi MT, Hlawe MM, Oseni TO,Sibiya TE.Effects of organic fertilizers on growth,  
239 yield, quality and sensory evaluation of red lettuce (*Lactuca sativa*L.) ‘VenezuelaRoxa’. Agric and  
240 Biol J North America2010;1(6):1319-1324
- 241 12. Masarirambi M T, Dlamini P., Wahome PK,Oseni TO.Effects of chicken manure on growth,  
242 yield and quality of lettuce (*Lactuca sativa* L.) ‘Taina’ under a lath house in a semi-arid sub-  
243 tropical environment. American Eurasian J Agric EnvironSci,2012;12 (3): 399-406
- 244 13. Msibi, B.M., Mukabwe WO,Manyatsi AM, MhazoNMasarirambi MT. Effects of liquid  
245 manure on growth and yield of spinach (*Beta vulgaris* var*ciela*) in a sub-tropical environment in  
246 Swaziland.Asian J AgricSci,2014;6(2): 40-47

247 14. Kunene TR, Masarirambi MT, Wahome PK, Oseni TO. Influence of kraal manure, chicken  
248 manure and inorganic fertilizer on growth, yields and post-harvest quality of pepper (*Capsicum*  
249 *annuum* L.) in a sub-tropical environment. *Asian J Advances Agric Res*, 2019;11(1):1-11

250 15. Mabuza TZ, Masarirambi M.T. Nxumalo KA, Wahome PK. Effects of different rates of cattle  
251 manure on growth, yield and quality of pepper (*Capsicum annuum* L.) in a sub-tropical  
252 environment of Eswatini (Swaziland). *Asian J Advances Agric Res*, 2019;11(1):1-7

253 16. Dlamini SN, Masarirambi MT, Wahome P K, Oseni T.O. The effects of organic fertilizers  
254 on the growth and yield of amaranth (*Amaranthus hybridus* L.) grown in a lath house. *Asian J*  
255 *Advances Agric Res*, 2020;12(1):1-10

256

257

258

259

260

UNDER PEER REVIEW