

1
2
3
4
5
6
7
8
9

Original Research Article

**Efficacy of excised-bud (EB) and half-corm (HC)
at four physiological growth stages on plantlet
regeneration of *Musa* genotypes**

UNDER PEER REVIEW

10 **ABSTRACT**

Aims: This study was conducted to determine which of excised buds (EB) and half corms (HC) from 3 *Musa* genotypes at four growth stages of mother plants would produce the most plantlets and also how scarification affects the number of plantlets regenerated.

Study design: Treatments comprised three *Musa* genotypes at four growth stages and two macro-propagation methods – excised bud and half-corm in a randomized complete block design with 4 replications.

Place and Duration of Study: International Institute of Tropical Agriculture (IITA) High Rainfall Station, Onne (4°51'N, 7° 03'E, 10m above sea level), Rivers State, Nigeria for eighteen months.

Methodology: Propagules, excised buds and half corms from a tetraploid cooking banana hybrid BITA 3; tetraploid plantain hybrid PITA 14, and a cooking banana landrace Cardaba, at 6-month vegetative, pre-flowering, post-flowering and bunch harvest stages were planted to regenerate plantlets. At bunch harvest growth stage, additional excised buds and half corms were scarified to find out the effect on regeneration of plantlets.

Results: Excised buds and half corms did not differ significantly ($P = .05$) in number of plantlets produced in PITA 14 irrespective of growth stage but bunch harvest stage was best. In BITA 3, excised buds produced significantly more plantlets than half corms at the 6-month vegetative and bunch harvest stages. However, at the pre-flowering stage, half corms produced significantly more plantlets than excised buds. In Cardaba, half corms were significantly better at all growth stages especially bunch harvest stage. In all *Musa* genotypes, scarification increased significantly the number of plantlets.

Conclusion: Excised buds or half corms at any growth stage or at bunch harvest stage for PITA 14; and excised buds at 6-month vegetative or bunch harvest stage for BITA 3 with half corm at pre-flowering stage are best. For Cardaba, half corm at any growth stage or bunch harvest stage was best.

Keywords: [*Musa* genotype, Macro-propagation, scarification, excised bud, half corm]

11

12

13

1. INTRODUCTION

14

Musa is vegetatively propagated and planting materials can be produced either by micro-propagation or by macro-propagation. Natural replacement of suckers through regeneration of landraces, hybrids or clones, the method preferred by rural farmers [1,2] is very slow because apical dominance causes “shy suckering” which prevents buds from developing into suckers until the reproductive phase of the mother plant has commenced, ~~causes slow regeneration~~ [3]. Apical dominance is controlled by a growth hormone that is produced in the terminal bud and inhibits growth of the lateral shoots [4]. Besides being slow, natural suckering does not yield enough suckers of the desired varieties and pest and disease susceptibility can be quite high in the event of outbreaks [1,5,6] which can easily wipe-out whole plantations. The result is a serious shortage of clean planting materials and this shortage of planting materials is considered a serious constraint to the rapid expansion of *Musa* production [7]. While micro-propagation methods can provide large quantities and high quality planting materials [8], the tender plantlets require great care in the first 2 months of planting. Also the equipment, technical skills, cost and highly controlled environment required are beyond the reach of resource poor farmers [9]. Therefore, macro-propagation has remained an effective alternative method which requires less capital and skills to produce large numbers of better-quality *Musa* planting material by farmers. However, some problems associated with macro-propagation include use of large numbers of parent materials, large space required for multiplication, and lack of uniform size of plantlets. Macro-propagation techniques include traditional methods that use whole suckers or relatively large pieces of the parent plants to produce planting materials; these are usually bulky and difficult to transport. Common methods of macro-propagation include decapitation and false decapitation. Decapitation is the destruction of the terminal bud to increase the sprouting and development of suckers [10]. False decapitation also destroys the main apex in order to remove apical dominance, but it maintains the entire plant [11]. The rate of

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

Comment [DS1]: Do bananas have lateral shoots? Or, What is meant by lateral shoots in bananas?

Comment [DS2]: What is the relationship between "pest and disease susceptibility" and "natural suckering"?

40 suckering using the above methods range from nine to fourteen suckers per annum
41 [7,12]. Stripping of older sheaths to expose buds as well as mulching and earthing of the
42 exposed buds have also been used to increase the number of suckers obtained from a
43 mother plant [13,14,15]. Whole plants (peepers or sword) have also been used to
44 produce planting materials [16,17]. The whole corm and corm-bits are used to produce
45 few plantlets of uniform size [18]. The study [19] showed that the corm method could
46 produce about five hundred suckers within eight months. More recent macro-propagation
47 techniques involve methods that employ whole suckers or relatively large pieces of corm
48 tissue to produce planting material in a propagator [20]. Other methods of macro-
49 propagation utilize the whole corm, split-corm, split-bud and corm-bit techniques [21,22].
50 Depending on variety, one corm can yield an average of 10 seedlings, which can be
51 increased by a factor of 3–4 by removal of the apical meristem of emerging lateral buds
52 [23]. Hence, alternative methods based on bud excision are being investigated. The
53 method requires that buds be removed from the mother corm, and incubated in the pre-
54 nursery to generate shoots. Prior to transplanting, the shoots obtained could be further
55 multiplied by making incisions-scarification, which could yield a higher number of uniform
56 size plantlets. Different banana propagation techniques can give different number of
57 shoots [24]; while number of shoots produced is also influenced by the banana's
58 genotype [25,26]. To the best of our knowledge no studies have investigated the use of
59 excised buds (EB) and half corms (HC) obtained at four physiological growth stages of
60 different *Musa* species as propagules for production of planting material. It is important
61 that such a study be conducted in order to provide critical information on the ideal
62 physiological growth stage that can provide the maximum number of propagules for rapid
63 multiplication of each *Musa* spp. This study was therefore conducted specifically to:

- 64 1. Assess and compare the rate of regeneration of excised buds (EB) and half corms
65 (HC) obtained at four physiological growth stages as viable macro-propagation
66 materials in 3 *Musa* genotypes
- 67 2. Find out how scarification of excised buds and half corms affect the rate of
68 regeneration of plantlets of 3 *Musa* genotypes

70 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / 71 METHODOLOGY

72 This study was carried out at the International Institute of Tropical Agriculture (IITA) High
73 Rainfall Station, Onne (4°51'N, 7° 03'E, 10m above sea level), in Rivers State, south-
74 eastern Nigeria. The rainfall pattern is monomodal, distributed over a 10month period
75 from February through December, with an annual average of 2400mm. Relative humidity
76 remains high all year round with mean values of 78% in February, increasing to 89% in
77 the months of July and September. The mean annual minimum and maximum
78 temperatures are 25°C and 27°C, respectively, while solar radiation / sunshine lasts an
79 average of 4hours daily [27]. The soil is derived from coastal sediments of the Niger
80 Delta, freely drained and acidic (pH 4.3), and made up of mainly Kaolinite. Onne soils
81 are also high in phosphorus 60mg kg⁻¹, manganese 0.2mmol kg⁻¹, but low in nitrogen
82 [28,29].

83 Preparation of Macro-propagation Materials

84 Three *Musa* genotypes comprising one tetraploid cooking banana hybrid BITA-3 (TMBx
85 5295-1) that is resistant to black Sigatoka disease; one tetraploid plantain hybrid PITA 14
86 (TMPx 7152-2) which is high yielding, short cycling, and resistant to black Sigatoka and
87 to Banana streak virus diseases; and a cooking banana landrace (Cardaba) resistant to
88 black Sigatoka disease were the source of the macro-propagation materials. Corms were
89 harvested from 5 field-grown plants of each of these genotype source materials at each
90 of the four physiological stages of growth as follows:

- 91 (i) At 6-month vegetative growth stage,
- 92 (ii) At onset of flowering growth stage,
- 93 (iii) At end of flowering growth stage, and
- 94 (iv) At bunch harvest growth stage

95 The harvested corms were immediately washed under a running tap. Roots were
96 trimmed off and plant debris was removed to expose all buds on the corm, after which

Comment [DS3]: Only 9-12 plantlets per year?

Comment [DS4]: This is so much different from 9-12 suckers (line 40)

97 each corm was split into two equal halves. One part was used as half-corm while buds
98 were excised from the other half. Buds of about 150kg?? each were excised from the
99 corms with a locally fabricated mechanical extractor to ensure uniformity in size of buds.
100 These two macro-propagation methods, excised-bud (EB) and half-corm (HC) were used
101 in multiplication of plantlets in order to determine which technique produced the highest
102 number of healthy plantlets. [Please add with the figure of methods](#),

Formatted: English (United States)

103 Treatment Applications and Experimental Design

104 Treatments were the three *Musa* genotypes and four physiological stages described
105 earlier and two macro-propagation methods – excised bud (EB) and half-corm (HC)
106 giving a 3 X 4 X 2 factorial combination in a randomized complete block design with 4
107 replications. The excised buds were initially surface sterilized with 20% solution of
108 Sodium hypochlorite, and allowed to stand for 5minutes in a solution of 6g copper-
109 oxochloride in one litre of water to prevent decay, after which they were allowed to air-dry
110 for 4hours. The treated materials were planted at a spacing of 20cm by 20cm in a
111 germination chamber consisting of a concrete basin filled with a mixture of sawdust and
112 poultry manure at a ratio of 3:1 and watering was done as required. At the bunch harvest
113 stage of physiological growth, an additional set of excised buds and half corms were
114 scarified. Scarification was by making 2 incisions on the excised buds and on the
115 growing point of the half corms in order to find out how scarification would affect the
116 number of regenerated plantlets compared to non scarified ones. [Illustration of the
117 methods is very essential](#),

Formatted: English (United States)


118 Data Collection and Statistical Analyses

119 Sprouting was considered to have occurred when the buds grew about 5cm above the
120 soil level. The final number of regenerated plantlets was recorded. The data were
121 subjected to square-root transformation, prior to analysis of variance (ANOVA) to test
122 treatment effects. All data were analysed using the general linear model procedure of
123 Statistical Analyses Software [30] and any effects found to be significant have been
124 tested at a significance level of 5% while means were compared using the Least
125 Significant Difference (LSD) at $P = .05$.

126 3. RESULTS

127 Plantlet Regeneration at 6-month Vegetative Growth Stage

128 At the 6-month vegetative stage of growth, excised buds (EB) from the cooking banana
129 hybrid (BITA 3) produced significantly ($P = .05$) more (333% more) plantlets than its half
130 corm (HC) counterpart (Fig 1). However, there was no significant difference ($P = .05$) in
131 the number of plantlets produced by the excised buds and the half corms in the plantain
132 hybrid (PITA 14). In the cooking banana (Cardaba) the half corms produced significantly
133 ($P = .05$) more (600% more) plantlets. In fact, excised buds did not produce any plantlets
134 in Cardaba.
135
136




137

138
139
140
141
142
143
144
145
146
147
148
149

Fig. 1. Number of regenerated plantlets from excised buds and half corms obtained at the 6-month vegetative stage of growth in 3 *Musa* genotypes

Plantlet Regeneration at Pre-Flowering Growth Stage

In Figure 2, at the pre-flowering growth stage, half corms (HC) produced significantly ($P = .05$) more (120% more) plantlets than the excised buds (EB) in the cooking banana hybrid (BITA 3). There was no significant difference ($P = .05$) in the number of plantlets produced by the half corms and the excised buds in the plantain hybrid (PITA 14). Again at this stage of growth, half corms produced significantly ($P = .05$) more (900% more) plantlets than excised buds which did not produce any plantlets in the cooking banana landrace Cardaba




150
151
152
153
154
155
156
157
158
159
160

Fig. 2. Number of regenerated plantlets from excised buds and half corms obtained at the pre-flowering stage of growth in 3 *Musa* genotypes

Plantlet Regeneration at Post Flowering Growth Stage

At post flowering growth stage, there was no significant difference ($P = .05$) in the number of plantlets produced by the excised buds and the half corms in the cooking banana hybrid (BITA 3) (Fig 3) as well as in the plantain hybrid (PITA 14). However there was a significant difference ($P = .05$) in the number of plantlets produced by the excised buds and the half corms in the cooking banana Cardaba




161
162
163
164
165

Fig. 3. Number of regenerated plantlets from excised buds and half corms obtained at the post-flowering stage of growth in 3 *Musa* genotypes

166 **Plantlet Regeneration at Bunch Harvest Growth Stage**

167 Excised buds (EB) obtained at bunch harvest stage in the cooking banana hybrid (BITA
168 3) produced significantly ($P = .05$) more (86% more) plantlets than its half corm (HC)
169 counterpart (Fig.4). There was no significant difference ($P = .05$) in the number of
170 plantlets produced by excised buds and half corm at this stage of growth in the plantain
171 hybrid (PITA 14). The half corms produced significantly ($P = .05$) more (550% more)
172 plantlets than excised buds in the cooking banana landrace Cardaba. This was the most
173 productive physiological stage for using the half corm propagule in the cooking banana
174 Cardaba
175



176
177 **Fig. 4. Number of regenerated plantlets from excised buds and half corms**
178 **obtained**
179 **at the bunch harvest stage of growth in 3 Musa genotypes**
180

181 **Effects of Scarification on Excised Buds (EB) and Half Corms (HC)**

182 **Excised Buds**

183 Scarification increased significantly ($P = .05$) by more than 4 times, the number of
184 plantlets produced by excised buds (EB) in the cooking banana hybrid (BITA 3) and by
185 approximately 3 times in the plantain hybrid (PITA 14) (Fig 5). In the cooking banana
186 Cardaba, scarification resulted in a significant ($P = .05$) increase by doubling the number
187 of plantlets produced compared to non scarified buds.

188 **Half Corms**

189 Scarification increased significantly ($P = .05$) by more than double, the number of
190 plantlets produced by half corms (HC) in the cooking banana hybrid (BITA 3) and by
191 50% in the plantain hybrid (PITA 14) compared to non scarified half corms (Fig 5). In the
192 cooking banana Cardaba, scarification increased significantly ($P = .05$) the number of
193 plantlets by 91% compared to non scarified half corms.
194




Fig. 5. Number of regenerated plantlets from scarified and non scarified excised buds and half corms obtained at the bunch harvest stage of growth in 3 *Musa* genotypes

4. DISCUSSION

Musa genotypes and Macro-propagation methods/ propagules

Generally, hybrid cooking banana (BITA 3) produced significantly the highest number of plantlets from excised buds (EB) at both 6-month vegetative and bunch harvest stages of growth. However, at the pre-flowering stage, half corm produced the highest number of plantlets indicating at which stage to use each propagation method/ propagule. For hybrid plantain (PITA 14), excised buds and half corms produced the highest number of plantlets at bunch harvest stage of growth. Cooking banana, Cardaba, produced the highest number of plantlets from half corms obtained at harvest, followed by those obtained at pre-flowering and 6-month vegetative stages in that order. Of the 3 genotypes, significantly higher numbers of plantlets were obtained from the hybrids than from the cooking banana Cardaba. Generally excised buds were best for the hybrid cooking banana and half corm for cooking banana Cardaba while either of the propagules could be used for hybrid plantain. The higher number of plantlets obtained from hybrids suggests genetic improvement of the hybrids over the banana landrace Cardaba. Higher suckering of the hybrids over their plantain parents has been attributed to their ability to overcome apical dominance [31]. According to [11] sucker production and development are influenced by growth hormones produced by the mother plant, which is regulated by the Ad gene [31]. It could also be from hormonal changes which occur during the lifespan of any plant [32]. Besides the action of hormones, apical dominance may be influenced by the physiological stage of the plant which depends upon the source-sink relationship. The rate of regeneration is determined by the amount of assimilates from leaves to sink which in turn depend upon age and vigour of the plant [33]. The higher regeneration of the hybrids over the cooking banana landrace may also be due to the higher ploidy level of the hybrids. Higher plant ploidy could confer higher vigour resulting from higher sink accumulation and consequently result in a higher number of plantlets [34,35].

Scarification

Scarification of excised buds and half corms may have (a) triggered hormones that induced cell division, callus formation and elongation, (b) increased efficiency of uptake and translocation within the propagules and accumulation at the active sites and (c) may have removed any anatomical barrier limiting formation of plantlets causing higher regeneration of plantlets in both propagules [22,36,37,38].

4. CONCLUSION

This study found that macro-propagation of the hybrid plantain PITA 14 could be done using either excised buds or half corms at any physiological growth stage but ideally at

237 bunch harvest stage for best results. In the cooking banana hybrid BITA 3, excised buds
238 at the 6-month vegetative or bunch harvest stage proved optimal, while use of half corms
239 is best at pre-flowering stage. In the cooking banana Cardaba, half corms at all
240 physiological growth stages could be used although bunch harvest stage was the most
241 productive. Scarification of excised buds and half corms increased number of plantlets in
242 all genotypes.

243

244 **COMPETING INTERESTS**

245 Authors have declared that no competing interests exist.

246

247

REFERENCES

- 248 1. Kasyoka MR, Mwangi M, Kori N, Gitonga N, Muasya R. Evaluating the
249 macropropagation efficiency of banana varieties preferred by farmers in eastern and
250 central Kenya. Second RUFORUM Biennial Meeting, Entebbe. Uganda, 2010:449–
251 503.
- 252 2. Ocimati W, Karamura D, Rutikanga A, Sivirihauma C, Ndung V, Ntamwira J, Kamira
253 M, Kanyaruguru JP, Blomme G. Agronomic practices used by farmers in the
254 management of *Musa* across different agro-ecological zones in Burundi, Eastern
255 Democratic Republic of Congo and Rwanda. In: Blomme G, Van Asten P, Vanlauwe
256 B. (eds.). *Banana Systems in the Humid Highlands of Sub-Saharan Africa: Enhancing Resilience and Productivity*, (Wallingford, UK: CAB International). pp.
257 2013:175–190. <https://doi.org/10.1079/9781780642314.0175>.
- 258 3. Braide J, Wilson GF. Plantain decline: a look at possible causes. *Paradisiaca*.
259 1980;4:3-7.
- 260 4. De Langhe EA, Swennen R, Wilson G. Aspects hormonaux du rejetonnage des
261 bananiers plantains. *Fruits*. 1983;38: 318-325.
- 262 5. Manzur MD. *In situ* mass propagation of the FHIA-20 banana hybrid using
263 benzylaminopurine. *Infomusa* 2001;10(1), 3–4.
- 264 6. Butler D. Fungus threatens top banana, *Nature*. 2013;504:195-196
- 265 7. Wilson GF, Vuylsteke D, Swennen R. Rapid multiplication of plantain: an improved
266 field technique. In: International Cooperation for Effective Plantain and Banana
267 Research, Proceedings, 3rd IARPB/INIBAPA, Montpellier, France. 1987:24-26.
- 268 8. Singh TD, Singh CH, Nongalleima K, Moirangthem S, Devi HS. Analysis of growth,
269 yield potential and horticultural performance of conventional vs. micropropagated
270 plants of *Curcuma longa* var. Lakadong. *Afr. J. Biotech.* 2013;12:1604-1608
- 271 9. George P, Manuel J. Low cost tissue technology for the regeneration of some
272 economically important plants for developing countries. *Inter. J. Agric, Environ.*
273 *Biotec.* 2013;6,703- 711
- 274 10. Green-Ortiz JJ, Fierro C. Removal of the apex as a means of propagating plantains.
275 In: Proceedings of the 24th Annual Congress of the American Society for Horticultural
276 Science. Tropical Region University of Mayaguez, Puerto Rico. 1976:223-225.
- 277 11. Swennen RA. Physiological Study of the Suckering Behaviour in Plantain (*Musa* cv.
278 AAB). Faculty of Agriculture, Catholic Uni. Louv., Belgium. Ph.D. Thesis.
279 1984:132:80p.
- 280 12. Noupadja P. Study of three field multiplication techniques for generating planting
281 material of *in vitro* propagated plantain (*Musa* cv. AAB). *MUSAFRICA*. ISSN
282 1995;117-2266.
- 283 13. Barker GW. A system of maximum multiplication of banana plants. *Tropical*
284 *Agriculture*, Trinidad. 1959;36:275-284.
- 285 14. De Langhe EA. Multiplication végétative accélérée, en plantation, du bananier plantain
286 (Bosua). Bulletin Information l'INEAC. 1961;10 (2): 69-90.
- 287 15. Charpentier JM. La remontée du meristème central du bananier. *Fruits*. 1966;21
288 (3):103-119.
- 289 16. Strivastava RP. Selection of planting materials for banana. Allahabad Farm-
290 Management.. 1963;37:18-20.
- 291 17. Baiyeri KP, Aba SC. A review of protocols for macropropagation in *Musa* species.
292 *Fruit, Veg. Cereal Sci. Biotech.* 2007;1:110-15.
- 293

- 294 18. Shanmugavelu KG, Aravindakshan K, Sathiamoorthy S. Banana, Taxonomy,
295 Breeding and Production Technology. Metropolitan Book Co. PVT. LTD. New Delhi.
296 1992:459p
- 297 19. Adelaja BA. Rapid on-farm multiplication technique for plantain and banana. Annual
298 Reports. National Horticultural Research Institute. 1983. ISSN 0795-4115.
- 299 20. Tenkouano A, Hauser S, Coyne D, Coulibaly O. Clean planting materials and
300 management practices for sustained production of banana and plantain in Africa.
301 *Chronica Horticulturae*. 2006;46:14–18
- 302 21. Faturoti B, Tenkouano A, Lemchi J, Nnaji N. Rapid Multiplication of plantain and
303 banana: Macropropagation techniques, IITA Report; 2002
- 304 22. Tumuhimbise R, Talengera D. Improved Propagation Techniques to Enhance the
305 Productivity of Banana (*Musa* spp.). *Open Agriculture*. 2018;3:138–145.
306 <https://doi.org/10.1515/opag-2018-0014>
- 307 23. Njeri N, Mwangi M, Gathu R, Mbaka J, Kori N, Muasya R. Assessing effectiveness of
308 macro-propagation technology to produce healthy seedlings of banana varieties with
309 high market demand in eastern and central provinces, Kenya. Second RUFORUM
310 Biennial Meeting 20–24 September 2010, Entebbe, Uganda. Research Application
311 Summary. 2010:531–533;
- 312 24. Karamura E, Staver C. Strategies for improving bananas and plantains seed
313 systems in Africa. BMGF Technical report. 2010.
- 314 25. Vuylsteke D, Swennen R, Wilson GF, De Langhe E. Phenotypic variation among *in*
315 *vitro* propagated plantain (*Musa* sp. Cultivar 'AAB'), *Sceintia Hort*. 1998;36:79-88.
- 316 26. Singh HP, Uma S, Selvarajan R, Karihaloo J.L. Micro-propagation for production of
317 quality banana planting material in Asia-Pacific, Asia-Pacific Consort. *Agric. Biotec.*,
318 NewDelhi, India; 2011.
- 319 27. Ortiz R, Austin PD, Vuylsteke D. IITA High Rainfall Station African humid forest.
320 *American Journal of Horticultural Science*. 1997;32:969-972.
- 321 28. Winslow MD. Silicon disease resistance and yield of rice genotypes under upland
322 cultural conditions. *Crop Science*. 1992;32:1208-1213. Retrieved from
323 <http://www.nal.usda.gov/>
- 324 29. Swennen R, Vuylsteke D, Ortiz R. Phenotypic diversity and patterns of variation in
325 West and Central African Plantains (*Musa* spp., AAB group *Musaceae*). *Economic*
326 *Botany*. 1995;49:320-327.
- 327 30. SAS Institute Inc. SAS User's Guide. Statistical Analysis Institute Inc., Cary, NC;
328 1992.
- 329 31. Ortiz R, Vuylsteke DR. Genetics of apical dominance in plantain (*Musa*, AAB group)
330 and improvement of suckering behaviour. *Journal of American Society of*
331 *Horticultural Science*. 1994;119 (5):1050-1053.
- 332 32. Peñarrubia L, Moreno J. Senescence in plants and crops. In: Pessaraki M. (ed.).
333 Handbook of plant and crop physiology. Marcel Dekker, Inc. New York. 1995:461-
334 481.
- 335 33. Eckstein K, Robinson JC, Davie SJ. Physiological responses of banana (*Musa* AAA;
336 Cavendish sub-group) in the subtropics 111. Gas exchange, growth analysis and
337 source-sink interaction over a complete crop cycle. *American Journal of Horticultural*
338 *Science*. 1995;70 (1):169-180.
- 339 34. Craenen K, Ortiz R. Effect of black Sigatoka resistance locus *bs₁* and ploidy level on
340 fruit and bunch traits of plantain-banana hybrids. *Euphytica*. 1996;87:97-101.
- 341 35. Ortiz R. Secondary polyploids heterosis and evolutionary crop breeding for further
342 improvement of the plantain and banana (*Musa* spp., L.) genome. *Theoretical and*
343 *Applied Genetics*. 1997;94:113-120.
- 344 36. Naqvi SSM. Plant growth hormones: Growth promoters and inhibitors. In: Pessaraki
345 M. (ed.). Handbook of plant and crop physiology. Marcel Dekker, Inc. New York.
346 1995:527-556
- 347 37. Mok DW, Mok MC. Cytokinin metabolism and action. *Ann. Rev. Plant*
348 *Physiol. Plant Mol. Biol.*, 2001;52:89-118
- 349 38. Kim EK, Hahn EJ, Murthy HN, Pack KY. High frequency of shoot
350 multiplication and bulbet formation of garlic in liquid cultures. *Plant Cell*
351 *Tiss. Org. Cult.*, 2003;73: 231-236.