

REVIEW ON THE ROOT, STEM AND LEAF INITIATIONS IN PLANTS

Abstract

The root and shoot apical meristem serve as sources of pluripotent cells and provide new cells for repetitive organ [initiation], they are the major meristematic regions on which plant development take place. New meristems are continuously formed when plants produce new branches or lateral roots thus making the Understanding of meristem function central to the understanding of how plants can establish so many different growth types, ranging from tiny herbs to huge trees. The sizes and numbers of meristems that are initiated during later development control the size and number of fruits and the generation of seeds. The development of a lateral root from a limited number of cells requires tightly coordinated asymmetric cell divisions to generate cell diversity and tissue patterns which characteristically involves the specification of founder cells, followed by a number of cellular changes until the cells divide and give rise to unequally sized daughter cells. This has been carefully discussed under corresponding headings. Leaf development exemplifies the dynamic nature and flexibility of plant development in response to internal and external cues which is evidenced in the fact that two plants, even if genetically identical, do not look the same, two leaves on the same plant are different, and the final shape of a leaf is not predetermined when it starts to form. Leaves evolved from lateral branches following the acquisition of determinate growth and a flat structure, thus the specification of organ initiation involves a complex network of genetic, hormonal and mechanical factors which has been discussed in this work.

Keywords: Initiation, Meristem, pluripotent, hormones.

INTRODUCTION

Plant development depends on the activity of two main meristems; the root meristem and the shoot apical meristem, which serve as a source of pluripotent stem cells and provide new cells for repetitive organ initiation [1]. Thus, plant meristems are the stem cell niches that allow stem cells to remain undifferentiated and to proliferate. Meristems are dynamic structures that can be generated de novo, for example during flower formation [2]. Furthermore, new meristems are continuously formed when plants produce new branches or lateral roots. Understanding meristem function is therefore central to understand how plants can establish so many different growth types, ranging from tiny herbs to huge trees. Furthermore, the sizes and numbers of meristems that are initiated during later development control the size and number of fruits and the generation of seeds [2].

The formation of a plant root system takes place post-embryonically and relies on de novo formation of organs. Typically, these lateral root organs are initiated close to the root tip and emerge in the differentiation zone. Over the past few years, knowledge about the regulatory mechanisms behind many aspects of lateral root formation has increased considerably [3, 4]. The

Comment [MM1]: Small letter for initiation

Comment [MM2]: Lower case u for understanding

Comment [MM3]: Delete these words

Comment [MM4]: Delete

Comment [MM5]: Abstract not relating to roots and stems as it is with the leaves yet it should be a summary of the title and body

Comment [MM6]: Why is this exactly like in abstract

development of a lateral root from a limited number of cells requires tightly coordinated asymmetric cell divisions to generate cell diversity and tissue patterns. This characteristically involves the specification of founder cells, followed by a number of cellular changes until the cells divide and give rise to unequally sized daughter cells [5].

The development of plant leaves follows a common basic program that is flexible and is adjusted according to species, developmental stage and environmental circumstances. Leaves initiate from the flanks of the shoot apical meristem and develop into flat structures of variable sizes and forms. This process is regulated by plant hormones, transcriptional regulators and mechanical properties of the tissue.

ROOTS DEVELOPMENT IN PLANT

Roots serve a multitude of functions such as anchorage, as the conduit to supply both nutrients and water to the plant from the soil, as a location for the synthesis and exchange of various plant hormones, as storage organs of plant resources [6]. Plant roots grow in a highly heterogeneous environment—the soil, and possess an ability to react to this heterogeneity and modify the form of their root system as a consequence. This is a “phenotypic plasticity” which is influenced by a genetic program and environmental factors and the ultimate configuration of the root system. To understand the morphogenesis of roots it is necessary to define the organization of the root meristem and attempt to determine the fate map of cells emerging from the root meristem [7]. As the majority of the root system consists of lateral roots, which is controlled by various, genetic, hormonal and nutritional factors. Primary root growth occurs from the root apical meristem (RAM) and is dependent on a stem cell niche or microenvironment being established giving rise to the quiescent Centre (QC) [8]. The RAMs are maintained by retaining a stem cell reservoir and a pool of undifferentiated initial cells.

ROOT MORPHOGENESIS

The Root Apex

Many of the important unanswered questions in root development involve events that occur at the root apex. For example, little is known about the nature of the stem cells, how cell files are established, how cell numbers and vascular patterning are determined, what controls the organization and size of the meristem, how root hair initials are formed, how cell expansion is regulated, and, perhaps most important, what controls the cell cycle and the planes of cell division. Many of these are general questions that apply equally well to morphogenesis in other parts of the plant. However, several aspects of root morphogenesis serve to simplify the study of these basic questions. First, the root apical meristem is easily accessible (not enclosed the way the shoot apical meristem is), essentially transparent (due to a lack of pigment), and lacks branching primordia. Second, the root as a whole is a simple organ that displays a radial symmetry in the external layers of cells. Third, root morphogenesis normally occurs in a reiterative and uniform fashion, without any major change in the organization of the root apex. Thus, all stages of root development are apparent at all times, and there is nothing analogous to the vegetative to floral conversion that occurs in shoot meristems. Fourth, roots have relatively few differentiated cell types. Finally, the various developmental processes are largely confined to classically defined “zones” along the length of the root, as indicated in Figure 1. These include the meristematic zone (site of cell divisions), elongation zone (cell expansion), and specialization

Comment [MM7]: Write in plural as Plants

Comment [MM8]: Delete

Comment [MM9]: Delete

Comment [MM10]: Delete as and replace with and

Comment [MM11]: Remove this, is this coping from some source as it is

Comment [MM12]: Sentence sounds hanging

Comment [MM13]: Any references to these thoughts

zone (cell differentiation). Although this zonal classification is probably too simplistic (there is overlap in the cellular processes occurring in the various zones), it nonetheless serves to emphasize the spatial separation of these processes in cell files of roots.

To fully understand the morphogenesis of roots, it is necessary to define the organization of the root meristem and determine the fate of cells that emerge from the meristem. One of the most revealing analyses of this type was performed on the root of the water fern *Azolla* [9]. The precise placement and timing of each cell division were determined and mapped, providing a complete cellular fate map of the root. In higher plants, the characterization of multicellular root meristems led to the discovery of a unique set of cells, the quiescent center, which is located at the center of the root apex but undergoes relatively infrequent cell divisions [10]. The precise function of the cells of the quiescent center and their relationship to the rest of the meristem still need to be defined.




Figure 1. Schematic Outline of Internal Processes and External Factors That Control Root Development [10].

The classically defined zones of cellular activities are indicated, as are the environmental stimuli that influence root morphogenesis.

Lateral Roots

Branching in roots differs from branching in stems. Lateral roots do not develop directly from cells in the apical meristem but rather develop from differentiated cells in a special layer (the pericycle) located just below the endodermal layer. Evidence that this process involves redifferentiation comes from an analysis of the expression pattern of the enzyme hyoscyamine 6- α -hydroxylase, which is localized to pericycle cells. Upon induction of lateral roots, expression of the enzyme decreases dramatically [11]. Not all pericycle cells have an equal probability of giving rise to lateral root primordia; lateral roots are not usually initiated near the root tip (suggesting the presence of an inhibitor diffusing from the tip) and normally are formed from pericycle cells opposite to the xylem elements [12].

The formation of lateral roots is a particularly intriguing aspect of root development because it represents the initiation of a new meristem and may provide clues as to how the primary root meristem arises during embryogenesis [12]. However, important questions regarding lateral root development remain unanswered. The initiation of lateral roots must involve some sort of signal that is perceived by pericycle cells. A single cell may be activated that then recruits others; alternatively, multiple cells may simultaneously perceive the signal. Once the meristem is formed, it must elaborate a root that somehow forces its way through the existing cortex and epidermal tissue and emerges into the external environment. It should be possible to address these issues by identifying genes specifically expressed in these cells and by isolating mutants blocked at different stages of lateral root development [12].

Plant Hormones

All aspects of root development are profoundly affected by plant hormones, with the strongest effects attributed to auxin, cytokinins, and ethylene [13, 14]. Because of the difficulty in interpreting the effect of exogenously applied hormones on internal hormone ratios, there is considerable controversy in the literature as to the relative importance of various growth regulators on root development [14]. Alternative approaches, such as the analysis of transgenic plants in which hormone ratios have been modified *in vivo* by expression of hormone biosynthetic enzymes [15, 16, 17] and the characterization of mutants with reduced hormone biosynthesis or altered sensitivity [18], may help resolve many of the outstanding questions.

Comment [MM14]: Why bold

Environmental influences

Although root morphology is guided by a genetic program, the ultimate configuration of a root system under natural conditions is largely determined by environmental factors. The effects of gravity on root growth have been explored most extensively; roots generally respond in a positive fashion to gravity, with the root cap cells playing a major role in perception [19].

Roots also respond to chemical gradients; they proliferate in regions of the soil that contain high concentrations of certain ions, such as nitrate or phosphate [20, 21]. In addition, root growth can be influenced by the soil moisture content, with roots penetrating deeper when soil moisture is low [22] and developing air spaces (aerenchyma) when the soil is waterlogged [23]. Although roots usually grow in a subterranean environment, light has been shown to affect root extension, gravitropism, and lateral root production in some species [24, 25, 26].

Furthermore, the growth of roots can be influenced by temperature gradients [27, 28], mechanical impedance [29], aeration [30], and the roots of adjacent plants [31]. The morphological plasticity of roots represents one of the most interesting aspects of root development, and there is a clear need for further exploration of the manner in which external stimuli affect the root's developmental program.

LEAF DEVELOPMENT

Leaf development exemplifies the dynamic nature and flexibility of plant development in response to internal and external cues. Just as two plants, even if genetically identical, do not look the same, two leaves on the same plant are different, and the final shape of a leaf is not predetermined when it starts to form. Leaves evolved from lateral branches following the acquisition of determinate growth and a flat structure [32, 33, 34, 35]. Leaves can be divided into

Comment [MM15]: As indicated on comment why emphasize leaves only in abstract, this is almost the same as in the abstract

two basic forms: simple and compound. A simple leaf has an entire, continuous lamina, whereas a compound leaf is composed of multiple subunits termed leaflets, each resembling a simple leaf [34]. On a developmental timescale, simple leaves differentiate and flatten relatively fast, whereas compound leaves are in some ways intermediate forms between lateral branches and simple leaves [35].

LEAF INITIATION

During initiation, a distinct domain within the shoot apical meristem (SAM), which is separated from the rest of the SAM by a boundary domain, is specified [36, 37]. According to the Hofmeister principle, leaf initiation occurs at the point most distant from existing primordia, leading to the hypothesis that existing primordia generate an inhibition field [38, 39]. The specification of organ initiation involves a complex network of genetic, hormonal and mechanical factors.

THE MECHANICS OF LEAF INITIATION

Accumulating evidences point to the potential role of mechanics in the regulation of leaf positioning and initiation, either as a signal or via differential tissue properties [40]. Tissue and cell geometry, mechanical stresses, cellulose and microtubule orientation and growth directions have long been proposed to be involved in morphogenesis, both in plants and animals [41]. Using atomic force microscopy (AFM), cell walls in the Central Zone were found to be stiffer and their stiffness more variable than that of cell walls in the Peripheral Zone [42]. In agreement, using osmotic manipulations, the Central Zone and the Peripheral Zone were shown to differ in their mechanical properties, and these differences correlated with increased growth in the Peripheral Zone [43]. Mechanical forces were also shown to affect microtubule orientation [44]. This effect is mediated by the microtubule-severing protein KATANIN that promotes growth variability between neighboring cells [45]. Thus, correlated mechanical properties, growth directions and microtubule orientation characterize the Central Zone, Peripheral Zone and the boundary region between them.

MECHANICAL FORCES IN LEAF INITIATION AND GROWTH

Organ initiation involves loosening of the cell wall by cell-wall modifiers, such as expansins and pectin methylesterases (PMEs) [46, 47]. Auxin induces these factors, and they thus partially mediate the effect of auxin on organ initiation [48]. Additionally, mechanical forces as well as the cell wall were shown to affect the levels and polar distribution of PIN1 within the cell [49, 50, 51]. However, mechanical stress appears to affect microtubule orientation and PIN1 polarization in parallel, as disruption of microtubule polymerization did not affect organ initiation in the short term [44, 49]. Together, these studies point to a scenario in which organ initiation is instructed in part by the geometry of the Shoot Apical Meristem and differential mechanical properties of distinct regions within the Shoot Apical Meristem. These properties affect the growth properties of the tissue as well as auxin distribution. Auxin, in turn, induces changes in cell wall properties and also interacts with transcription factors and additional hormones to specify leaf initiation and growth.

GENES THAT REGULATE INITIATION

Specification of the organ initiation domain is also accompanied by differential expression of genes that regulate the balance between meristematic and initiation fates. For example, class I Knotted-Like Homeobox (KNOXI) transcription factors, which promote Shoot Apical Meristem function, are expressed in the Central Zone of the Shoot Apical Meristem and are down regulated at the site of organ initiation [52]. KNOXI expression is down regulated at the site of leaf initiation by ARP [ASYMMETRIC LEAVES1 (AS1), ROUGH SHEATH2 (RS2), PHANTASTICA] transcription factors, together with the LBD protein AS2 and the chromatin remodeling factor HIRA, promoting specification of the organ initiation domain [53]. Several recent studies have established a role for chromatin remodeling factors in the repression of KNOXI genes by AS1-AS2 in *Arabidopsis*. For example, AS1 interacts with the histone deacetylase HDA6, and several KNOXI genes show increased acetylation in *hda6* mutants [54]. In addition, the AS1-AS2 complex has recently been shown to recruit POLYCOMB-REPRESSIVE COMPLEX 2 (PRC2), a complex involved in chromatin structure modification, to the promoters of two KNOXI genes, possibly enabling their stable repression at later stages of leaf development [55]. The expression of KNOXI genes is also regulated by BLADE ON PETIOLE (BOP) [56, 57], JAGGED LATERAL ORGANS (JLO) [58] and auxin [59]. KNOXI proteins, in turn, feedback to regulate the auxin response [60, 61, 62]. KNOXI proteins also regulate the balance between cytokinin, which promotes meristematic fate, and gibberellic acid (GA), which promotes differentiation [63, 64, 65, 66]. Thus, KNOXI proteins coordinate the activity of several plant hormones during the specification of the distinct domains in the Shoot Apical Meristem, enabling the balance between continuous Shoot Apical Meristem function and organ initiation.

Additional early markers of the leaf initiation domain include genes encoding transcription factors from the AINTEGUMENTA (ANT)-like (AIL)/PLT family, and genes from the YABBY (YAB) family of HMG-like proteins. AIL/PLT genes have been shown to promote organ initiation and growth in *Arabidopsis* [67, 68] and to partially mediate the effect of MP on organ initiation [69]. Recently, some AIL/PLT genes were suggested to affect phyllotaxis by promoting auxin biosynthesis in the Central Zone of the Shoot Apical Meristem [70]. Phenotypes resulting from mutations and overexpression of YABBY genes suggest that they are involved in the specification of organ fate and the suppression of meristem fate, in addition to their role in leaf polarity [71, 34].

THE BALANCE BETWEEN MORPHOGENESIS AND DIFFERENTIATION

Following initiation, the leaf primordia undergoes growth, morphogenesis and differentiation in a highly flexible process that ultimately gives rise to the final leaf shape. This flexibility is manifested in a continuum of leaf shapes, ranging from very simple to highly complex [34]. The flexibility of leaf development is achieved by modulating the overall rate of leaf maturation and the balance between morphogenesis and differentiation, as well as specific patterning events [72].

THE REGULATION OF LAMINA INITIATION AND GROWTH

One of the first events during primary morphogenesis is the initiation and growth of a lamina, leading to the formation of a flat rather than a radial structure. Lamina initiation and growth are thought to require the juxtaposition of abaxial and adaxial tissues [73], and a number of genes

have been implicated in this process. YABBY and AIL/PLT genes, for example, have been linked to the promotion of lamina outgrowth and expansion in *Arabidopsis*, maize and rice [74, 75, 76, 34]. In addition, JAGGED (JAG) and its paralog NUBBIN (NUB) are redundant, positive regulators of leaf blade growth in *Arabidopsis* [77, 78]. Accordingly, *jag nub* double mutants have a reduced leaf blade area, and combined *jag-1/fil/yab3* mutations result in a severe loss of blade development. Recently, JAG was shown to directly repress meristematic and cell cycle genes, thus promoting differentiation [79]. WOX transcription factors have also been linked to the promotion of blade outgrowth in several species. For example, the *Nicotiana sylvestris* WOX gene mutant *lam1* has vestigial lamina-less leaves that lack mesophyll differentiation [80, 81, 82]. It therefore appears that an overlapping set of genes is involved in lamina initiation and expansion and in leaf initiation, and that these processes require repression of meristematic fate. It remains to be seen how the activities of these different regulators of lamina initiation and growth are coordinated.

Lamina growth also requires coordination between the epidermis and the mesophyll layers, and it was recently shown that the transcriptional co-activator *Angustifolia3* (AN3) is produced only in mesophyll cells but moves into the epidermis to promote growth in both layers [83]. AN3 was subsequently shown to modulate transcription through interaction with chromatin-remodeling factors [84].

Several genes involved in basic cellular functions have also been shown to influence leaf lamina growth. In *Arabidopsis*, ribosomal protein mutants have pointed leaves with more prominent marginal serrations, possibly due to a decrease in the relative cellular growth rate [85, 86, 87]. Furthermore, the E3 ubiquitin-ligase BIG BROTHER (BB) can repress plant organ growth, probably by marking cellular proteins for degradation [88]. Recently, poly (A) polymerases (PAPS) have been shown to influence leaf size and shape, probably by affecting the expression of specific subsets of relevant genes [89]. In *Cardamine hirsuta*, the ribosome-associated protein SIMPLE LEAF3 also affects leaf growth and leaflet development [90]

ROLE AND MAINTENANCE OF THE MARGINAL BLASTOZONE

Leaf growth is mostly determinate. However, transient indeterminate growth is maintained in specific regions of the leaf. These include a growing region at the leaf base or the leaf tip, depending on the species [91], and regions in the leaf margin that possess organogenic potential, known as marginal blastozones (MBs) [92]. The marginal blastozone is responsible for lamina initiation and the organogenesis of marginal structures. Classic and recent research has shown that compound leaf development requires prolonged activity of the marginal blastozone during primary morphogenesis. Genetic and hormonal factors that regulate marginal blastozone activity were shown to partially overlap with those regulating SAM activity, in accordance with the evolutionary origin of a leaf as a modified shoot [93, 32]. The temporal and spatial length of the marginal blastozone activity determines the extent of the indeterminate phase in leaf growth and the consequent level of leaf complexity [92].

HORMONES THAT AFFECT THE BALANCE BETWEEN MORPHOGENESIS AND DIFFERENTIATION IN LEAF

The rate of leaf maturation is also regulated by several plant hormones, many of which interact with the transcription factors discussed above. For example, GA was found to regulate cell proliferation and expansion rate in *Arabidopsis* leaves [94]. Not surprisingly, GA negatively regulates leaf complexity in tomato. Upon increased GA levels or response, only primary leaflets with smooth margins are formed and the leaves mature faster than wild-type leaves do [95, 96, 97, 98]. Similarly, *Nicotiana solanifolia* mutants produce primary and intercalary leaflets only, with smooth margins, possibly due to elevated GA levels [95]. These findings suggest that GA promotes leaf maturation. However, in some species GA has the opposite effect of inducing more compound leaves [99, 100, 101]. For example, in pea, GA and auxin positively promote leaf dissection during leaf morphogenesis by prolonging the temporal window during which acropetally initiated leaflets are produced [101]. KNOXI and TCP proteins have also been linked to GA dynamics. KNOXI proteins negatively affect GA levels by repressing the GA biosynthesis gene *GA20ox* and activating the GA inactivation gene *GA2ox*. These effects on GA homeostasis mediate the function of KNOXI in tuning the SAM-leaf boundary and in modulating compound leaf development in *Arabidopsis*, maize, tobacco and tomato [102, 103]. By contrast, the TCP protein LA positively affects GA homeostasis in tomato [104]. Modulation of GA homeostasis therefore appears to be a common mechanism by which different transcription factors tune the rate of maturation and differentiation.

Cytokinin was also shown to affect the balance between morphogenesis and differentiation in leaf development. Increased cytokinin degradation in *Arabidopsis* leaf primordia accelerated cell expansion and early termination of cell proliferation, demonstrating that cytokinin delays the onset of cell differentiation [105, 106]. Interestingly, lettuce (*Lactuca sativa*) leaves that overexpress the *Arabidopsis* KNOXI gene *BP* acquire characteristics of indeterminate growth, which is associated with the accumulation of specific types of cytokinins [107]. Cytokinin was also shown to be involved in the maintenance of prolonged morphogenetic activity in the tomato leaf margin (Shani *et al.*, 2010). Genetic and molecular analysis indicated that cytokinin acts downstream of KNOXI activity in delaying leaf maturation. Conversely, promotion of leaf maturation by CIN-TCPs in *Arabidopsis* is mediated by reducing leaf sensitivity to cytokinin. TCP4 was shown to interact with the chromatin remodeler BRAHMA to directly activate the expression of *ARR16*, which encodes an inhibitor of cytokinin responses [108]. Interestingly, the class I TCPs TCP14 and TCP15, which are thought to act antagonistically with class II TCPs, positively regulate cytokinin response [109]. Thus, the antagonistic effect of KNOXI and TCP transcription factors on leaf maturation converges on the regulation of the GA/cytokinin homeostasis. It is interesting to see whether other factors affecting the rate of leaf maturation also affect this homeostasis. GA and cytokinin were also shown to antagonize the response of each other during tomato leaf development [97]. Leaves of some species, including tomato, maintain morphogenetic activity after leaf expansion, leading to further variability in leaf shape, as seen in the *cla* mutant. Interestingly, GA and cytokinin were both shown to modulate this late morphogenetic activity in tomato [110, 104]. Cumulatively, these studies suggest that the flexibility of leaf shape is achieved by tuning the balance between hormones that promote indeterminate state, such as cytokinin, and hormones that promote differentiation, such as GA.

Comment [MM16]: Attend to this reference

THE ROLE OF AUXIN DURING LEAF INITIATION

The plant hormone auxin has emerged as a central regulator of organ initiation. Points of auxin response maxima are observed prior to organ initiation. These are generated by auxin biosynthesis in the Shoot Apical Meristem and by directional auxin transport facilitated by the PIN-FORMED1 (PIN1) auxin transporter [111, 112, 113, 114]. Accordingly, inhibition of polar auxin transport or a mutation in *PIN1* inhibits organ initiation, whereas auxin application in the Peripheral Zone of meristems is sufficient to induce organ initiation. Mutations in auxin biosynthesis genes from the YUCCA family also inhibit organ initiation. Auxin gradients and/or flow are thought to direct PIN1 polarization in a positive-feedback loop, and auxin depletion by developing primordia is thought to comprise at least part of the hypothesized inhibitory field [38].

Comment [MM17]: Reference, is it 111 – 114 or 38 below

The response to auxin is mediated by transcription factors known as auxin response factors (ARFs). Mutations in the plant ARF gene *MONOPTEROS* (*MP*) lead to a wide variety of aberrant phenotypes, including reduced flower initiation. *MP* might therefore mediate the activity of auxin in organ initiation [115, 69]. However, it should be noted that much of the research to date on organ initiation in *Arabidopsis* has involved inflorescence meristems, which form flower meristems rather than leaf primordia. Flower meristems in *Arabidopsis* are derivatives of axillary meristems that form in the axils of cryptic bracts, which are miniature underdeveloped leaves [116]. Leaf and flower initiation are thus different processes and their regulation might, at least in part, involve different factors. This is exemplified by *Arabidopsis pin1* mutants: in *pin1* inflorescences, flower initiation is completely abolished, whereas leaf initiation is only partially compromised in *pin1* vegetative meristems, as well as when multiple *PIN* genes are mutated [117, 118].

Leaf initiation is closely correlated with the initiation of the midvein, a vascular strand in the middle of the leaf. The mid vein initiates from the auxin maxima at the leaf initiation site and gradually connects to the existing vasculature [119]. A strand of high auxin concentration marks the midvein initiation site and is correlated with a switch in PIN1 polarization, from polarization towards the convergence point in the outermost cell layer (L1) to basal localization towards the future midvein. This was hypothesized to be accompanied by a switch from auxin transport towards the highest auxin concentration to transport in the direction of auxin flow [119]. Distinct regulators of PIN1 localization were shown to be involved in these different phases, whereas the localization towards the convergence point is regulated in part by the serine/threonine kinase PINOID [120], which phosphorylates PIN1 [121] (Huang *et al.*, 2010), the switch to basal polarization is regulated by the MAB4 gene family [122, 123]. In angiosperm species other than the Brassicaceae, leaf initiation and vascular formation were suggested to be regulated by distinct members of the PIN family [124].

Comment [MM18]: Attend to this reference

THE BALANCE BETWEEN AUXIN AND CYTOKININ

In addition to auxin, leaf initiation involves the plant hormone cytokinin, which plays an important role in Shoot Apical Meristem maintenance [125, 126, 127]. As we discuss below, the specification of leaf initiation involves a delicate balance and complex feedback relationship between auxin and cytokinin.

Recently, light has been shown to be essential for leaf initiation in tomato, and this effect is mediated by both auxin and cytokinin [128]. In maize, the response regulator (RR) protein ABPHYL1 (ABPH1) is expressed at the site of future leaf initiation together with PIN1, and both are induced by cytokinin [129]. ABPH1 regulates Shoot Apical Meristem size and phyllotaxis, and belongs to a family of two-component RRs that are rapidly induced by cytokinin and are thought to act as negative regulators of the cytokinin response [130]. ABPH1 positively regulates organ initiation, perhaps by inhibiting the cytokinin response. In *Arabidopsis*, the RRs ARR7 and ARR15 are negatively regulated by MP, and mutants with elevated cytokinin levels suppress the flower initiation defect of *mp* mutants. This led to the hypothesis that auxin and cytokinin act synergistically in organ initiation in the *Arabidopsis* SAM, in contrast to their antagonistic action in the root [131, 132]. Thus, RRs are involved in balancing Shoot Apical Meristem size and organ initiation in both maize vegetative meristems and *Arabidopsis* inflorescences, but have opposing interactions with auxin in these two tissues. More recently, AHP6, another negative regulator of cytokinin signaling, was shown to regulate flower initiation downstream of auxin in a non-autonomous manner [133]. Together, these studies suggest that a fine coordination of local auxin and cytokinin responses regulates and stabilizes leaf initiation. However, whereas auxin is clearly a positive regulator of organ initiation, the exact effect of the cytokinin response on initiation is more complex, and its role appears to be dependent on species and developmental context. Furthermore, relative rather than absolute levels of cytokinin signaling, as well as the ratio between cytokinin and auxin and the tuning of hormone sensitivities, probably play a role.

CONTROLLING LEAF SIZE

Leaf size is largely dependent on the plant species, but is variable to a certain extent and is also tuned by environmental factors. Recent studies have shown that leaf size and the rate of leaf maturation are regulated by partially overlapping pathways, including those involving CIN-TCPs, ARP/AS2 and hormone dynamics. However, leaf size is not always correlated with leaf complexity or with the number of cells, pointing to partially independent regulation of these three processes [133, 134]. The issue of leaf size has been the recent focus of several reviews to which we refer the reader [135].

Comment [MM19]: Reference

MARGINAL PATTERNING IN SIMPLE AND COMPOUND LEAVES

Marginal patterning, which occurs during both primary and secondary morphogenesis, involves the formation of serrations, lobes and leaflets at the leaf margin, and flexibility in these patterning events further expands the variability in leaf form. The formation of marginal structures results from differential growth in adjacent regions and can be caused by a local restriction or promotion of growth [136, 137, 138]. As we discuss below, marginal patterning in simple and compound leaves involves partially overlapping mechanisms, many of which involve auxin signaling.

The interaction between auxin and *NO APICAL MERISTEM (NAM)/CUP-SHAPED COTYLEDON (CUC)* transcription factors is involved in marginal patterning in both simple and compound leaves. NAM/CUC transcription factors regulate many developmental processes, including boundary specification [36]. In simple *Arabidopsis* leaves, they promote leaf serrations [139], and in compound leaves they promote leaflet specification and separation [93]. The expression of *NAM/CUC* mRNA marks the boundary between the leaf margin and the future

leaflet in an array of species with compound leaves, and NAM/CUC silencing leads to leaf simplification [140, 141, 142]. A subset of *CUC* genes are negatively regulated by miR164. In tomato, the transgenic expression of a miR164-insensitive form of the NAM/CUC gene *GOBLET* (*GOB*) leads to ectopic initiation events in the leaflet margins, which later fuse to produce a final leaf form that is relatively simple and deeply lobed. Thus, both reduced and expanded expression domains of *GOB* lead to leaflet fusion [141], suggesting that distinct and sufficiently distant domains of *GOB* expression are essential for leaflet separation. NAM/CUC genes are therefore conserved modulators of the positioning and separation of marginal structures. In tomato, the *Potato-leaf* (*C*) gene, an ortholog of the *Arabidopsis* branching regulator *REGULATOR OF AXILLARY MERISTEMS1* (*RAX1*), also regulates leaf complexity; *c* mutants show reduced leaf complexity compared with the wild type, and smooth leaf blade margins. Interestingly, combining the *c* and the *gob* mutations results in the elimination of leaflet initiation, suggesting that they act partially redundantly in marginal patterning [143].

Auxin was also shown to be involved in leaf serration [139, 59] and in the initiation and separation of leaflets and lobes from the margin of compound leaf primordia, similar to its role in leaf initiation from the flanks of the SAM. In compound leaves, inhibition of auxin transport or activity resulted in the development of simplified leaves. Furthermore, PIN1 subcellular localization was found to converge at sites pre-marking leaflet initiation, leading to peaks in expression of the auxin-response sensor DR5, whereas external auxin application led to ectopic lamina growth and/or leaflet initiation [144, 145, 146, 147, 148, 149]. These observations indicate that discrete auxin maxima promote leaflet initiation and growth. Interestingly, in *M. truncatula*, leaves of the *MtPIN10/SLM1* (the *Medicago PIN1* ortholog) mutant exhibit increased complexity and decreased marginal patterning, suggesting a more complex effect of auxin on leaf patterning in *Medicago*. However, the increased complexity might result from fusion of several leaves [150, 151].

A role for auxin in margin patterning has also been implied based on studies of the tomato *ENTIRE* (*E*, *SIIAA9*) gene, which encodes a protein from the Aux/IAA family of auxin response repressors [141,152]. Leaves of the tomato mutant *e* are much simpler than wild-type leaves [153] and *e* leaf primordia initiate leaflets, but these fuse during the formation of the final *e* leaf form [147, 153, 149]. In *e* leaf primordia, the expression of the *PINI:PINI-GFP* reporter is upregulated and the expression of the auxin response sensor DR5 expands to the entire leaf margin [147, 149]. These observations suggest that *E* restricts lamina growth between developing leaflets by inhibiting auxin response. Together, these studies demonstrate that auxin promotes the formation and growth of diverse marginal structures.

How do NAM/CUC proteins and auxin interact in marginal patterning? Combining computational modeling and genetic approaches, it was proposed that, in *Arabidopsis*, *CUC2* promotes PIN1 localization, and auxin in turn represses *CUC2* expression, leading to regular patterns of leaf serrations [139]. Whereas in *Arabidopsis* auxin is thought to regulate *NAM/CUC* expression in both the SAM and the leaf [36, 139, 154, 112, 155], auxin in tomato affects *GOB* expression in apices but not in leaf primordia. Furthermore, the auxin response appears to act downstream of *GOB* in tomato leaf development, and it seems to be affected by both *GOB* and *E* [147]. Combining the *gob* and *e* mutations led to the complete elimination of leaflet initiation, suggesting that these factors also act via independent pathways [147]. These studies show that

Comment [MM20]: Is this a subtitle, might need to restructure sentence

the interaction between NAM/CUCs and auxin patterns margins in both simple *Arabidopsis* and compound tomato leaves, but the details of this interaction are tuned to pattern diverse leaf forms. The tomato LYRATE (LYR) gene, an ortholog of *JAG*, was shown to promote organ growth at the leaf margin, similar to the role of *JAG* in promoting growth of the main leaf lamina in *Arabidopsis*. Leaves of the *lyr* mutant have more leaflets in comparison to the wild type, and LYR overexpression leads to leaflet fusion [156, 157]. LYR possibly affects auxin response or distribution [157], and it will be interesting to see how it interacts with NAM/CUC genes in marginal patterning. Interestingly, CUC genes, *ASI* and auxin responsive genes were identified as targets of CIN-TCPs in *Arabidopsis* [158]. Combining downregulation of CIN-TCPs and upregulation of CUCs and *STIMPY/WOX9* genes led to substantially increased margin elaboration in *Arabidopsis*, giving rise to a leaf shape that resembles a compound leaf [159]. These studies show that common genes can affect both leaf maturation and marginal patterning.

Recent work has identified the REDUCED COMPLEXITY (RCO) homeodomain protein as necessary for leaflet development [138]. RCO is present in *C. hirsuta* and has evolved via duplication in the Brassicaceae family, but was lost in *Arabidopsis*, thus contributing to leaf simplification. RCO is thought to promote compound-leaf development by inhibiting growth between leaflets, but it does not affect auxin response distribution [138]. Another recent work compared the level of leaf dissection in various species of the genus *Capsella* and found that diversification in the RCO paralogs can account for naturally occurring leaf-shape variation in this Brassicaceae family. RCO expression can be temperature responsive in some cases, which is possibly involved in the plasticity of leaf shape under different temperatures [160]. In both *Capsella* and *C. hirsuta*, differential expression rather than protein function is thought to account for the evolution of the function in leaf complexity. It will be interesting to see how RCO interacts with other regulators of marginal patterning.

In addition to the genes and hormones discussed above, components of the trans-acting short interference RNA (tasiRNA) pathway are involved in leaf marginal patterning. Mutations in several genes in the tomato tasiRNA pathway, which are negative regulators of ARF2, 3 and 4, were shown to underlie the tomato 'wiry' syndrome of very narrow leaves with reduced complexity [161]. Interestingly, compromised tasiRNA pathway activity in *M. truncatula* led to a milder phenotype of increased leaf lobing with no effect on the number of leaflets [162], whereas leaf development in *Arabidopsis* was unaffected [163, 164]. Thus, whereas some mechanisms of marginal patterning are conserved among species, others differ substantially.

In summary, marginal patterning depends on the flexible positioning of regions in which lamina growth occurs and regions in which growth is inhibited. An indefinite number of leaf margin forms is achieved by tuning the interactions between plant hormones, transcription factors and growth regulators.

CONCLUSION

In conclusion, it is now clear that lateral root initiation comes about through the sequential activities of independent and / or overlapping auxin response modules. While various developmental systems have proved to be useful in understanding aspects of asymmetric cell division [5], lateral root initiation occurs deep within the primary root, hampering analysis of

what controls the polarity of the pericycle cells that will undergo this division, the coordination of the simultaneous nuclear migration, and the determination of the position of the division plane. However, with the improved imaging technology and the availability of new markers, elucidating these aspects of lateral root initiation and identifying the key players involved in these processes is possible. However, leaf development as a whole can be viewed as sequential developmental programs that are executed by different combinations of factors. Different developmental stages within a given program are often controlled by overlapping sets of factors or ‘tools’, thus comprising the ‘toolbox’ of leaf development. Particular examples of such tools that are involved in different stages of the same developmental program are discussed above. For instance, the involvement of YABBY family genes in several different stages and aspects of leaf development, together with their existence in seed plants only, has led to the notion that YABBY genes are integral to the ancestral specification of a leaf with determinate growth as opposed to a shoot from which a leaf is thought to have evolved [34]. Indeed, although for the purpose of clarity we have divided the analysis of leaf development into initiation, morphogenetic balance and marginal patterning, this division can be misleading, as many of the factors involved in fact affect several stages. For example, in addition to their role in leaflet initiation and separation, GOB, auxin and possibly ENTIRE/AUXIAA9 also affect the rate of leaf maturation [141].

Comment [MM21]: Review s more on leaves and less on the stem and roots

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

1. Dutta, A. C. (2007). Botany for Degree Students. (6th edition), *Oxford University Press, India*. pp. 196-197.
2. Kurup, S., Runions, J., Kohler, U., Laplaze, L., Hodge, S. and Haseloff, J. (2005). Marking cell lineages in living tissues. *Plant Journal* **42**: 444–453.
3. De Smet, I., Vanneste, S., Inze, D. and Beeckman T. (2006). Lateral root initiation or the birth of a new meristem. *Plant Molecular Biology* **60**: 871–887.
4. Peřet, B., De Rybel, B., Casimiro, I., Benkova, E., Swarup, R., Laplaze, L., Beeckman, T., and Bennett, M. (2009). *Arabidopsis* lateral root development: an emerging story. *Trends in Plant Science* **14**: 399–408.
5. De Smet, I. and Beeckman, T. (2011). Asymmetric cell division in land plants and algae: the driving force for differentiation. *Nature Reviews of Molecular Cell Biology* **12**: 177–188.
6. Schiefelbein, J. and Benfey, P. (1991). The development of plant roots: New approaches to underground problems. *Plant Cell*, **3**: 1147–1154.
7. Benfey, P. and Schiefelbein, J. (1994). Getting to the root of plant development: The genetics of *Arabidopsis* root formation. *Trends in Genetics*, **10**: 84–88.
8. Jiang, K. and Feldman L. (2005). Regulation of root apical meristem development. *Annual Review of Cell and Developmental Biology*, **21**: 485–509.
9. Gunning, B. (1982). *The root of the water fern Azolla: Cellular basis of development and multiple roles for cortical microtubules*. In *Developmental Order: Its Origin and Regulation*, S. Subtelny and P. 6. Green, eds. New York: Alan R. Liss, pp. 379-421.
10. Clowes, F. (1956). Localization of nucleic acid synthesis in root meristems. *Journal of Experimental Botany* **7**: 308-312.
11. Hashimoto, T., Hayashi, A., Amano, Y., Kohno, J., Iwanari, H., Usuda, S. and Yamada, Y. (1991). Hyoscyamine 6-p-hydroxylase, an enzyme involved in tropane alkaloid biosynthesis, is localized at the pericycle of the root. *Journal of Biological Chemistry*, **266**: 4648-4653.
12. Steeves, T. and Sussex, I. (1989). *Patterns in Plant Development*. (Cambridge, England: Cambridge University Press), p. 247.
13. Torrey, J. (1976). Root hormones and plant growth. *Annual Review of Plant Physiology*, **27**: 435-459.
14. Feldman, L. (1984). Regulation of root development. *Annual Review of Plant Physiology*, **35**: 223-242.
15. Klee, H., Horsch, R., Hinchee, M., Hein, M. and Offmann, N. (1987). The effects of overproduction of two *Agrobacterium tumefaciens* T-DNA auxin biosynthetic gene products in transgenic petunia plants. *Genes and Development*, **1**: 86-96.
16. Medford, J., Horgan, R., El-Sawi, Z. and Klee, H.J. (1989). Alterations of endogenous cytokinins in transgenic plants using a chimeric isopentenyl transferase gene. *Plant Cell*, **1**: 403-413.
17. Romano, C., Hein, M. and Klee, H. (1991). Inactivation of auxin in tobacco transformed with the indoleacetic acid-lysin synthetase gene of *Pseudomonas savastanoi*. *Genes and Development*, **5**: 438 - 446.
18. King, P. (1988). Plant hormone mutants. *Trends in Genetics*, **4**: 157-162.

19. Moore, R. and Evans, M. (1986). How roots perceive and respond to gravity. *American Journal of Botany*, **73**: 574-587.
20. Drew, M. and Saker, L. (1975). Nutrient supply and the growth of the seminal root system in barley. II. Localized compensatory increases in lateral root growth and rates of nitrate uptake when nitrate supply is restricted to only part of the root system. *Journal of Experimental Botany*, **26**: 79-90.
21. Fitter, A., Nichols, R. and Harvey, M. (1988). Root system architecture in relation to life history and nutrient supply. *Functional Ecology*, **2**: 345-351.
22. Coupland, R. and Johnson, R. (1965). Rooting characteristics of native grassland species in Saskatchewan. *Journal of Ecology*, **53**: 475-507.
23. Yu, P., Stolry, L. and Letey, J. (1969). Survival of plants under prolonged flooded conditions. *Agronomy Journal*, **61**: 844-847.
24. Lake, J. and Slack, G. (1961). Dependence on light of geotropism in plant roots. *Nature*, **191**: 300-302.
25. Wilkins, H. and Wain, R. (1974). The root cap and control of root elongation in *Zea mays* L. seedlings exposed to white light. *Planta*, **121**: 1-8.
26. Hart, J. and MacDonald, I. (1980). The influence of light on geotropism in cress roots. *Journal of Experimental Botany*, **31**: 903-911.
27. Pahlavanian, A. and Silk, W. (1988). Effect of temperature on spatial and temporal aspects of growth in the primary maize root. *Plant Physiology*, **87**: 529-532.
28. Fortin, M. and Poff, K. (1990). Thermotropism by primary roots of maize. *Plant Physiology*, **93**: 40 - 45.
29. Barley, K. and Greacen, E. (1967). Mechanical resistance as a soil factor influencing the growth of roots and underground shoots. *Advanced Agronomy*, **19**: 1-43.
30. Cannell, R. (1977). Soil aeration and compaction in relation to root growth and soil management. *Applied Biology*, **2**: 1-86.
31. Mahall, B. and Callaway, R. (1991). Root communication among desert shrubs. *Proceedings of National Academic Science, USA*, **88**: 874-876.
32. Floyd, S. and Bowman, J. (2010). Gene expression patterns in seed plant shoot meristems and leaves: homoplasy or homology? *Journal of Plant Research*, **123**: 43-55.
33. Kaplan, D. (2001). Fundamental concepts of leaf morphology and morphogenesis: a contribution to the interpretation of molecular genetic mutants. *International Journal of Plant Science*, **162**: 465-474.
34. Sarojam, R., Sappl, P., Goldshmidt, A., Efroni, I., Floyd, S., Eshed, Y. and Bowman, J. (2010). Differentiating Arabidopsis shoots from leaves by combined YABBY activities. *Plant Cell*, **22**: 2113-2130.
35. Zimmerman, W. (1952). Main results of the telome theory. *Paleobotanist*, **1**: 456-470.
36. Aida, M. and Tasaka, M. (2006). Genetic control of shoot organ boundaries. *Current Opinion on Plant Biology*, **9**: 72-77.
37. Žádníková, P. and Simon, R. (2014). How boundaries control plant development. *Current Opinion on Plant Biology*, **17**: 116-125.
38. Braybrook, S. and Kuhlemeier, C. (2010). *How a plant builds leaves*. *Plant Cell*, **22**: 1006-1018.
39. Snow, M. and Snow, R. (1932). Experiments on Phyllotaxis. I. The effect of isolating a primordium. *Philosophical Translation of Royal Society of London Biological Science*, **221**: 1-43.

40. Robinson, S., Burian, A., Couturier, E., Landrein, B., Louveaux, M., Neumann, E., Peaucelle, A., Weber, A. and Nakayama, N. (2013). Mechanical control of morphogenesis at the shoot apex. *Journal of Experimental Botany*, **64**: 4729-4744.
41. Green, P. (1980). Organogenesis-a biophysical view. *Annual Review of Plant Biology*, **31**: 51-82.
42. Milani, P., Gholamirad, M., Traas, J., Arnéodo, A., Boudaoud, A., Argoul, F. and Hamant, O. (2011). In vivo analysis of local wall stiffness at the shoot apical meristem in *Arabidopsis* using atomic force microscopy. *Plant Journal*, **67**: 1116-1123.
43. Kierzkowski, D., Nakayama, N., Routier-Kierzkowska, A., Weber, A., Bayer, E., Schorderet, M., Reinhardt, D., Kuhlemeier, C. and Smith, R. (2012). Elastic domains regulate growth and organogenesis in the plant shoot apical meristem. *Science*, **335**: 1096-1099.
44. Hamant, O., Heisler, M., Jonsson, H., Krupinski, P., Uyttewaal, M., Bokov, P., Corson, F., Sahlin, P., Boudaoud, A. and Meyerowitz, E. (2008). Developmental patterning by mechanical signals in *Arabidopsis*. *Science*, **322**: 1650-1655.
45. Uyttewaal, M., Burian, A., Alim, K., Landrein, B., Borowska-Wykręt, D., Dedieu, A., Peaucelle, A., Ludynia, M., Traas, J. and Boudaoud, A. (2012). Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in *Arabidopsis*. *Cell* **149**: 439-451.
46. Fleming, A., McQueen-Mason, S., Mandel, T. and Kuhlemeier, C. (1997). Induction of leaf primordia by the cell wall protein expansin. *Science*, **276**: 1415-1418.
47. Peaucelle, A., Braybrook, S., Le Guillou, L., Bron, E., Kuhlemeier, C. and Höfte, H. (2011). Pectin-induced changes in cell wall mechanics underlie organ initiation in *Arabidopsis*. *Current Opinion on Plant Biology*, **21**: 1720-1726.
48. Braybrook, S. and Peaucelle, A. (2013). Mechano-chemical aspects of organ formation in *Arabidopsis thaliana*: the relationship between auxin and pectin. *Plos One* **8**: e57813.
49. Heisler, M., Hamant, O., Krupinski, P., Uyttewaal, M., Ohno, C., Jönsson, H., Traas, J. and Meyerowitz, E. (2010). Alignment between PIN1 polarity and microtubule orientation in the shoot apical meristem reveals a tight coupling between morphogenesis and auxin transport. *Plos Biology*, **8**: e1000516.
50. Feraru, E., Feraru, M., Kleine-Vehn, J., Martinière, A., Mouille, G., Vanneste, S., Vernhettes, S., Runions, J. and Friml, J. (2011). PIN polarity maintenance by the cell wall in *Arabidopsis*. *Current Opinion on Plant Biology*, **21**: 338-343.
51. Nakayama, N., Smith, R., Mandel, T., Robinson, S., Kimura, S., Boudaoud, A. and Kuhlemeier, C. (2012). Mechanical regulation of auxin-mediated growth. *Current Opinion on Plant Biology*, **22**: 1468-1476.
52. Hay, A. and Tsiantis, M. (2010). KNOX genes: versatile regulators of plant development and diversity. *Development*, **137**: 3153-3165.
53. Barkoulas, M., Galinha, C., Grigg, S. and Tsiantis, M. (2007). From genes to shape: regulatory interactions in leaf development. *Current Opinion on Plant Biology*, **10**: 660-666.
54. Luo, M., Yu, C., Chen, F., Zhao, L., Tian, G., Liu, X., Cui, Y., Yang, J. and Wu, K. (2012). Histone deacetylase HDA6 is functionally associated with AS1 in repression of KNOX genes in *Arabidopsis*. *PLoS Genetics*, **8**: e1003114.

55. Lodha, M., Marco, C. and Timmermans, M. (2013). The Asymmetric Leaves complex maintains repression of KNOX homeobox genes via direct recruitment of Polycomb-repressive complex2. *Genes and Development*, **27**: 596-601.
56. Ha, C., Jun, J., Nam, H. and Fletcher, J. (2007). BLADE-ON-PETIOLE1 and 2 control Arabidopsis lateral organ fate through regulation of LOB domain and adaxial-abaxial polarity genes. *Plant Cell*, **19**: 1809-1825.
57. Ichihashi, Y., Aguilar-Martinez, J., Farhi, M., Chitwood, D., Kumar, R., Millon, L., Peng, J., Maloof, J. and Sinha, N. (2014). Evolutionary developmental transcriptomics reveals a gene network module regulating interspecific diversity in plant leaf shape. *Proceedings of National Academy of Science, USA*, **111**: E2616 – E2621.
58. Rast, M. and Simon, R. (2012). Arabidopsis Jagged Lateral Organs acts with Asymmetric Leaves2 to coordinate KNOX and PIN expression in shoot and root meristems. *Plant Cell* **24**: 2917-2933.
59. Hay, A., Barkoulas, M. and Tsiantis, M. (2006). ASYMMETRIC LEAVES1 and auxin activities converge to repress BREVIPEDICELLUS expression and promote leaf development in Arabidopsis. *Development* **133**: 3955-3961
60. Bolduc, N., Yilmaz, A., Mejia-Guerra, M., Morohashi, K., O'Connor, D., Grotewold, E. and Hake, S. (2012). Unraveling the KNOTTED1 regulatory network in maize meristems. *Genes and Development*, **26**, 1685-1690.
61. Scanlon, M., Henderson, D. and Bernstein, B. (2002). SEMAPHORE1 functions during the regulation of ancestrally duplicated knox genes and polar auxin transport in maize. *Development*, **129**: 2663-2673.
62. Tsiantis, M., Brown, M., Skibinski, G. and Langdale, J. (1999). Disruption of auxin transport is associated with aberrant leaf development in maize. *Plant. Physiology*, **121**: 1163-1168.
63. Hay, A., Kaur, H., Phillips, A., Hedden, P., Hake, S. and Tsiantis, M. (2002). The gibberellin pathway mediates KNOTTED1-type homeobox function in plants with different body plans. *Current Opinion on Plant Biology*, **12**: 1557-1565.
64. Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., Phillips, A., Hedden, P. and Tsiantis, M. (2005). KNOX action in Arabidopsis is mediated by coordinate regulation of cytokinin and gibberellin activities. *Current Opinion on Plant Biology*, **15**: 1560-1565.
65. Scofield, S., Dewitte, W., Nieuwland, J. and Murray, J. (2013). The Arabidopsis homeobox gene SHOOT MERISTEMLESS has cellular and meristem-organisational roles with differential requirements for cytokinin and CYCD3 activity. *Plant Journal*, **75**: 53-66.
66. Yanai, O., Shani, E., Dolezal, K., Tarkowski, P., Sablowski, R., Sandberg, G., Samach, A. and Ori, N. (2005). Arabidopsis KNOXI proteins activate cytokinin biosynthesis. *Current Opinion on Plant Biology*, **15**: 1566-1571.
67. Horstman, A., Willemsen, V., Boutilier, K. and Heidstra, R. (2014). Aintegumenta-Like proteins: hubs in a plethora of networks. *Trends in Plant Science*, **19**: 146-157.
68. Krizek, B. (2009). AINTEGUMENTA and AINTEGUMENTA-LIKE6 act redundantly to regulate arabidopsis floral growth and patterning. *Plant Physiology*, **150**: 1916 – 1929.
69. Yamaguchi, N., Wu, M., Winter, C., Berns, M., Nole-Wilson, S., Yamaguchi, A., Coupland, G., Krizek, B. and Wagner, D. (2013). A molecular framework for auxin-mediated initiation of flower primordia. *Development of Cell*, **24**: 271-282.

70. Pinon, V., Prasad, K., Grigg, S., Sanchez-Perez, G. and Scheres, B. (2013). Local auxin biosynthesis regulation by PLETHORA transcription factors controls phyllotaxis in *Arabidopsis*. *Proceedings of National Academy of Science USA*, **110**: 1107-1112.
71. Kumaran, M., Bowman, J. and Sundaresan, V. (2002). YABBY polarity genes mediate the repression of KNOX homeobox genes in *Arabidopsis*. *Plant Cell* **14**: 2761-2770.
72. Elliott, R., Betzner, A., Huttner, E., Oakes, M., Tucker, W., Gerentes, D., Perez, P. and Smyth, D. (1996). AINTEGUMENTA, an APETALA2-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *Plant Cell* **8**: 155-168.
73. Waites, R. and Hudson, A. (1995). phantastica: a gene required for dorsoventrality of leaves in *Antirrhinum majus*. *Development* **121**: 2143-2154.
74. Mizukami, Y. and Fischer, R. (2000). Plant organ size control: AINTEGUMENTA regulates growth and cell numbers during organogenesis. *Proceedings of National Academy of Science USA*, **97**: 942-947.
75. Juarez, M., Twigg, R. and Timmermans, M. (2004). Specification of adaxial cell fate during maize leaf development. *Development* **131**: 4533-4544.
76. Dai, M., Hu, Y., Zhao, Y., Liu, H. and Zhou, D. (2007). A Wuschel-Like HOMEBOX gene represses a YABBY gene expression required for rice leaf development. *Plant Physiology*, **144**: 380-390.
77. Ohno, C., Reddy, G., Heisler, M. and Meyerowitz, E. (2004). The *Arabidopsis* JAGGED gene encodes a zinc finger protein that promotes leaf tissue development. *Development* **131**: 1111-1122.
78. Dinneny, J., Weigel, D. and Yanofsky, M. (2006). NUBBIN and JAGGED define stamen and carpel shape in *Arabidopsis*. *Development* **133**: 1645-1655.
79. Schiessl, K., Muiño, J. and Sablowski, R. (2014). *Arabidopsis* JAGGED links floral organ patterning to tissue growth by repressing Kip-related cell cycle inhibitors. *Proceedings of National Academy of Science USA*, **111**: 2830 – 2835.
80. Ishiwata, A., Ozawa, M., Nagasaki, H., Kato, M., Noda, Y., Yamaguchi, T., Nosaka, M., Shimizu-Sato, S., Nagasaki, A. and Maekawa, M. (2013). Two WUSCHEL-related homeobox genes, narrow leaf2 and narrow leaf3, control leaf width in rice. *Plant Cell Physiology*, **54**: 779-792.
81. Lin, H., Niu, L., McHale, N., Ohme-Takagi, M., Mysore, K. and Tadege, M. (2013). Evolutionarily conserved repressive activity of WOX proteins mediates leaf blade outgrowth and floral organ development in plants. *Proceedings of National Academy of Science USA*, **110**: 366-371
82. Zhang, F., Wang, Y., Li, G., Tang, Y., Kramer, E. and Tadege, M. (2014). *Stenofolia* recruits TOPLESS to repress Asymmetric Leaves2 at the leaf margin and promote leaf blade outgrowth in *Medicago truncatula*. *Plant Cell*, **26**: 650-664.
83. Kawade, K., Horiguchi, G., Usami, T., Hirai, M. and Tsukaya, H. (2013). ANGUSTIFOLIA3 signaling coordinates proliferation between clonally distinct cells in leaves. *Current Opinion on Plant Biology*, **23**, 788-792.
84. Vercruyssen, L., Verkest, A., Gonzalez, N., Heyndrickx, K., Eeckhout, D., Han, S., Jegu, T., Archacki, R., Van Leene, J. and Andriankaja, M. (2014). ANGUSTIFOLIA3 binds to SWI/SNF chromatin remodeling complexes to regulate transcription during *Arabidopsis* leaf development. *Plant Cell*, **26**: 210-229.

85. Pinon, V., EtcHELLS, J., ROSSIGNOL, P., COLLIER, S., ARROYO, J., MARTIENSSSEN, R. and BYRNE, M. (2008). Three PIGGYBACK genes that specifically influence leaf patterning encode ribosomal proteins. *Development*, **135**: 1315-1324.
86. Szakonyi, D. and Byrne, M. (2011). Ribosomal protein L27a is required for growth and patterning in *Arabidopsis thaliana*. *Plant Journal*, **65**: 269-281.
87. Horiguchi, G., Van Lijsebettens, M., Candela, H., Micol, J. and Tsukaya, H. (2012). Ribosomes and translation in plant developmental control. *Plant Science*, **191**: 24-34.
88. Disch, S., Anastasiou, E., Sharma, V., Laux, T., Fletcher, J., and Lenhard, M. (2006). The E3 ubiquitin ligase BIG BROTHER controls *Arabidopsis* organ size in a dosage-dependent manner. *Current Opinion on Plant Biology*, **16**: 272-279.
89. Vi, S., Trost, G., Lange, P., Czesnick, H., Rao, N., Lieber, D., Laux, T., Gray, W., Manley, J. and Groth, D. (2013). Target specificity among canonical nuclear poly(A) polymerases in plants modulates organ growth and pathogen response. *Proceedings of National Academy of Science USA*, **110**: 13994-13999.
90. Kougioumoutzi, E., Cartolano, M., Canales, C., Dupre, M., Bramsiepe, J., Vlad, D., Rast, M., Dello Ioio, R., Tattersall, A. and Schnittger, A. (2013). SIMPLE LEAF3 encodes a ribosome-associated protein required for leaflet development in *Cardamine hirsuta*. *Plant Journal*, **73**: 533-545.
91. Tsukaya, H. (2014). Comparative leaf development in angiosperms. *Current Opinion on Plant Biology*, **17**: 103-109.
92. Hagemann, W. and Gleissberg, S. (1996). Organogenetic capacity of leaves: the significance of marginal blastozones in angiosperms. *Plant Syst. Evol.* **199**: 121-152.
93. Brand, A., Shirding, N., Shleizer, S. and Ori, N. (2007). Meristem maintenance and compound-leaf patterning utilize common genetic mechanisms in tomato. *Planta* **226**: 941-951.
94. Achard, P., Gusti, A., Cheminant, S., Alioua, M., Dhondt, S., Coppens, F., Beemster, G. and Genschik, P. (2009). Gibberellin signaling controls cell proliferation rate in *Arabidopsis*. *Current Opinion on Plant Biology*, **19**: 1188-1193.
95. Chandra-Shekhar, K. and Sawhney, V. (1991). Regulation of leaf shape in the *Solanifolia* mutant of tomato (*Lycopersicon esculentum*) by plant growth substances. *Annals of Botany*, **67**: 3-6.
96. Bassel, G., Mullen, R. and Bewley, J. (2008). Procera is a putative DELLA mutant in tomato (*Solanum lycopersicum*): effects on the seed and vegetative plant. *Journal Experimental Botany*, **59**: 585-593.
97. Fleishon, S., Shani, E., Ori, N. and Weiss, D. (2011). Negative reciprocal interactions between gibberellin and cytokinin in tomato. *New Phytology*, **190**: 609-617.
98. Jasinski, S., Tattersall, A., Piazza, P., Hay, A., Martinez-Garcia, J. F., Schmitz, G., Theres, K., McCormick, S. and Tsiantis, M. (2008). PROCERA encodes a DELLA protein that mediates control of dissected leaf form in tomato. *Plant Journal*, **56**: 603-612.
99. Robbins, W. (1957). Gibberellic acid and the reversal of adult hederata to a juvenile state. *American Journal of Botany*, **44**: 743-746.
100. Rogler, C. and Hackett, W. (1975). Phase change in *hedera helix*: induction of the mature to juvenile phase change by gibberellin A3. *Physiologia Plantarum* **34**: 141-147.
101. DeMason, D. and Chetty, V. (2011). Interactions between GA, auxin, and UNI expression controlling shoot ontogeny, leaf morphogenesis, and auxin response in *Pisum*

- sativum* (Fabaceae): or how the uni-tac mutant is rescued. *American Journal of Botany*, **98**: 775-791.
102. Sakamoto, T., Kamiya, N., Ueguchi-Tanaka, M., Iwahori, S. and Matsuoka, M. (2001). KNOX homeodomain protein directly suppresses the expression of a gibberellin biosynthetic gene in the tobacco shoot apical meristem. *Genes and Development*, **15**: 581-590.
103. Bolduc, N. and Hake, S. (2009). The maize transcription factor KNOTTED1 directly regulates the gibberellin catabolism gene *ga2ox1*. *Plant Cell* **21**: 1647-1658.
104. Yanai, O., Shani, E., Russ, D. and Ori, N. (2011). Gibberellin partly mediates Lanceolate activity in tomato. *Plant Journal*, **68**: 571-582.
105. Werner, T., Motyka, V., Strnad, M. and Schmulling, T. (2001). Regulation of plant growth by cytokinin. *Proceedings of National Academy of Science USA*, **98**: 10487-10492.
106. Holst, K., Schmülling, T. and Werner, T. (2011). Enhanced cytokinin degradation in leaf primordia of transgenic Arabidopsis plants reduces leaf size and shoot organ primordia formation. *Journal of Plant Physiol.* **168**: 1328-1334.
107. Frugis, G., Giannino, D., Mele, G., Nicolodi, C., Chiappetta, A., Bitonti, M., Innocenti, A., Dewitte, W., Van Onckelen, H. and Mariotti, D. (2001). Overexpression of KNAT1 in lettuce shifts leaf determinate growth to a shoot-like indeterminate growth associated with an accumulation of isopentenyl-type cytokinins. *Plant Physiology*, **126**: 1370-1380.
108. Efroni, I., Han, S.-K., Kim, H., Wu, M.-F., Steiner, E., Birnbaum, K., Hong, J., Eshed, Y. and Wagner, D. (2013). Regulation of leaf maturation by chromatin-mediated modulation of cytokinin responses. *Development of Cell*, **24**: 438-445.
109. Steiner, E., Efroni, I., Gopalraj, M., Saathoff, K., Tseng, T.-S., Kieffer, M., Eshed, Y., Olszewski, N. and Weiss, D. (2012). The Arabidopsis O-linked N-acetylglucosamine transferase SPINDLY interacts with class I TCPS to facilitate cytokinin responses in leaves and flowers. *Plant Cell* **24**: 96-108.
110. Shani, E., Ben-Gera, H., Shleizer-Burko, S., Burko, Y., Weiss, D. and Ori, N. (2010). Cytokinin regulates compound leaf development in tomato. *Plant Cell*, **22**: 3206-3217.
111. Benková, E., Michniewicz, M., Sauer, M., Teichmann, T., Seifertová, D., Jürgens, G. and Friml, J. (2003). Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell*, **115**: 591-602.
112. Heisler, M., Ohno, C., Das, P., Sieber, P., Reddy, G., Long, J. and Meyerowitz, E. (2005). Patterns of auxin transport and gene expression during primordium development revealed by live imaging of the Arabidopsis inflorescence meristem. *Current Opinion on Plant Biology*, **15**: 1899 – 1911.
113. Cheng, Y., Dai, X. and Zhao, Y. (2007). Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in Arabidopsis. *Plant Cell*, **19**: 2430-2439.
114. Reinhardt, D., Pesce, E., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M., Traas, J., Friml, J. and Kuhlemeier, C. (2003). Regulation of phyllotaxis by polar auxin transport. *Nature*, **426**: 255-260.

115. Hardtke, C. and Berleth, T. (1998). The Arabidopsis gene MONOPTEROS encodes a transcription factor mediating embryo axis formation and vascular development. *EMBO Journal*, **17**: 1405-1411.
116. Long, J. and Barton, M. (1998). The development of apical embryonic pattern in Arabidopsis. *Development* **125**: 3027-3035.
117. Guenot, B., Bayer, E., Kierzkowski, D., Smith, R., Mandel, T., Zadnikova, P., Benkova, E. and Kuhlemeier, C. (2012). Pin1-independent leaf initiation in Arabidopsis. *Plant Physiology*, **159**: 1501-1510.
118. Okada, K., Ueda, J., Komaki, M., Bell, C. and Shimura, Y. (1991). Requirement of the auxin polar transport system in early stages of Arabidopsis floral bud formation. *Plant Cell* **3**: 677-684.
119. Scarpella, E., Marcos, D., Friml, J. and Berleth, T. (2006). Control of leaf vascular patterning by polar auxin transport. *Genes and Development*, **20**: 1015-1027.
120. Friml, J., Yang, X., Michniewicz, M., Weijers, D., Quint, A., Tietz, O., Benjamins, R., Ouwerkerk, P. B. F., Ljung, K. and Sandberg, G. (2004). A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin efflux. *Science*, **306**: 862-865
121. Huang, F., Zago, M., Abas, L., van Marion, A., Galvan-Ampudia, C. and Offringa, R. (2010). Phosphorylation of conserved PIN motifs directs Arabidopsis PIN1 polarity and auxin transport. *Plant Cell*, **22**: 1129 – 1142..
122. Cheng, Y., Qin, G., Dai, X. and Zhao, Y. (2008). NPY genes and AGC kinases define two key steps in auxin-mediated organogenesis in Arabidopsis. *Proceedings of National Academy of Science USA*, **105**: 21017 – 21022.
123. Furutani, M., Nakano, Y. and Tasaka, M. (2014). MAB4-induced auxin sink generates local auxin gradients in Arabidopsis organ formation. *Proceedings of National Academy of Science USA*, **111**: 1198-1203.
124. O'Connor, D., Runions, A., Sluis, A., Bragg, J., Vogel, J., Prusinkiewicz, P. and Hake, S. (2014). A division in PIN-mediated auxin patterning during organ initiation in grasses. *PLoS Computational Biology*, **10**: 3447.
125. Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H. and Schmullig, T. (2003). Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *Plant Cell* **15**: 2532-2550.
126. Kurakawa, T., Ueda, N., Maekawa, M., Kobayashi, K., Kojima, M., Nagato, Y., Sakakibara, H. and Kyozuka, J. (2007). Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature*, **445**: 652-655.
127. Gordon, S., Chickarmane, V., Ohno, C. and Meyerowitz, E. (2009). Multiple feedback loops through cytokinin signaling control stem cell number within the Arabidopsis shoot meristem. *Proceedings of National Academy of Science USA*, **106**: 16529-16534.
128. Yoshida, S., Mandel, T. and Kuhlemeier, C. (2011). Stem cell activation by light guides plant organogenesis. *Genes and Development*, **25**: 1439-1450.
129. Lee, B., Yu, S. and Jackson, D. (2009). Control of plant architecture: the role of phyllotaxy and plastochron. *Journal of Plant Biology*, **52**: 277-282.
130. Giulini, A., Wang, J. and Jackson, D. (2004). Control of phyllotaxy by the cytokinin-inducible response regulator homologue ABPHYL1. *Nature*, **430**: 1031-1034.

131. Vidaurre, D., Ploense, S., Krogan, N. and Berleth, T. (2007). AMP1 and MP antagonistically regulate embryo and meristem development in *Arabidopsis*. *Development* **134**: 2561-2567.
132. Zhao, Z., Andersen, S., Ljung, K., Dolezal, K., Miotk, A., Schultheiss, S. and Lohmann, J. (2010). Hormonal control of the shoot stem-cell niche. *Nature*, **465**: 1089-1092.
133. Besnard, F., Refahi, Y., Morin, V., Marteaux, B., Brunoud, G., Chambrier, P., Rozier, F., Mirabet, V., Legrand, J. and Lainé, S. (2014). Cytokinin signalling inhibitory fields provide robustness to phyllotaxis. *Nature*, **505**: 417-421.
134. Efroni, I., Eshed, Y. and Lifschitz, E. (2010). Morphogenesis of simple and compound leaves: a critical review. *Plant Cell* **22**: 1019-1032.
135. Shleizer-Burko, S., Burko, Y., Ben-Herzel, O. and Ori, N. (2011). Dynamic growth program regulated by LANCEOLATE enables flexible leaf patterning. *Development*, **138**: 695-704.
136. Powell, A. and Lenhard, M. (2012). Control of organ size in plants. *Current Opinion on Plant Biology*, **22**: R360-R367.
137. Kawamura, E., Horiguchi, G. and Tsukaya, H. (2010). Mechanisms of leaf tooth formation in *Arabidopsis*. *Plant Journal*, **62**: 429-441.
138. Malinowski, R., Kasprzewska, A. and Fleming, A. (2011). Targeted manipulation of leaf form via local growth repression. *Plant Journal*, **66**: 941-952.
139. Vlad, D., Kierzkowski, D., Rast, M., Vuolo, F., Dello Ioio, R., Galinha, C., Gan, X., Hajheidari, M., Hay, A. and Smith, R. (2014). Leaf shape evolution through duplication, regulatory diversification, and loss of a homeobox gene. *Science*, **343**: 780-783.
140. Bilsborough, G., Runions, A., Barkoulas, M., Jenkins, H., Hasson, A., Galinha, C., Laufs, P., Hay, A., Prusinkiewicz, P. and Tsiantis, M. (2011). Model for the regulation of *Arabidopsis thaliana* leaf margin development. *Proceedings of National Academy of Science USA*, **108**, 3424-3429.
141. Blein, T., Pulido, A., Vialette-Guiraud, A., Nikovics, K., Morin, H., Hay, A., Johansen, I., Tsiantis, M. and Laufs, P. (2008). A conserved molecular framework for compound leaf development. *Science*, **322**: 1835-1839.
142. Berger, Y., Harpaz-Saad, S., Brand, A., Melnik, H., Sirding, N., Alvarez, J., Zinder, M., Samach, A., Eshed, Y. and Ori, N. (2009). The NAC-domain transcription factor GOBLET specifies leaflet boundaries in compound tomato leaves. *Development*, **136**: 823-832.
143. Cheng, X., Peng, J., Ma, J., Tang, Y., Chen, R., Mysore, K. and Wen, J. (2012). NO APICAL MERISTEM (MtNAM) regulates floral organ identity and lateral organ separation in *Medicago truncatula*. *New Phytology*, **195**: 71-84.
144. Busch, B., Schmitz, G., Rossmann, S., Piron, F., Ding, J., Bendahmane, A. and Theres, K. (2011). Shoot branching and leaf dissection in tomato are regulated by homologous gene modules. *Plant Cell*, **23**: 3595-3609.
145. Al-Hammadi, A., Sreelakshmi, Y., Negi, S., Siddiqi, I. and Sharma, R. (2003). The polycotyledon mutant of tomato shows enhanced polar auxin transport. *Plant Physiology*, **133**: 113-125.

146. Avasarala, S., Yang, J. and Caruso, J. (1996). Production of phenocopies of the lanceolate mutant in tomato using polar auxin transport inhibitors. *Journal of Experimental Botany*, **47**: 709-712.
147. Barkoulas, M., Hay, A., Kougioumoutzi, E. and Tsiantis, M. (2008). A developmental framework for dissected leaf formation in the Arabidopsis relative *Cardamine hirsuta*. *Natural Genetics*, **40**: 1136-1141.
148. Ben-Gera, H., Shwartz, I., Shao, M., Shani, E., Estelle, M. and Ori, N. (2012). ENTIRE and GOBLET promote leaflet development in tomato by modulating auxin response. *Plant Journal*, **70**, 903-915.
149. DeMason, D. and Polowick, P. (2009). Patterns of DR5::GUS expression in organs of pea (*Pisum sativum*). *International Journal of Plant Science*, **170**: 1-11.
150. Koenig, D., Bayer, E., Kang, J., Kuhlemeier, C. and Sinha, N. (2009). Auxin patterns *Solanum lycopersicum* leaf morphogenesis. *Development* **136**: 2997-3006.
151. Peng, J. and Chen, R. (2011). Auxin efflux transporter MtPIN10 regulates compound leaf and flower development in *Medicago truncatula*. *Plant Signal and Behavior*, **6**: 1537-1544.
152. Zhou, C., Han, L., Hou, C., Metelli, A., Qi, L., Tadege, M., Mysore, K. and Wang, Z. (2011). Developmental analysis of a *Medicago truncatula* smooth leaf margin1 mutant reveals context-dependent effects on compound leaf development. *Plant Cell*, **23**: 2106-2124.
153. Zhang, J., Chen, R., Xiao, J., Qian, C., Wang, T., Li, H., Ouyang, B. and Ye, Z. (2007). A single-base deletion mutation in SIIAA9 gene causes tomato (*Solanum lycopersicum*) entire mutant. *Journal of Plant Research*, **120**: 671-678.
154. Dengler, N. (1984). Comparison of leaf development in Normal (+/+), Entire (e/e), and Lanceolate (La/+) plants of tomato, *Lycopersicon esculentum* 'Ailsa Craig'. *Botanical Gazette*, **145**: 66-77.
155. Furutani, M., Vernoux, T., Traas, J., Kato, T., Tasaka, M. and Aida, M. (2004). PIN-FORMED1 and PINOID regulate boundary formation and cotyledon development in Arabidopsis embryogenesis. *Development* **131**: 5021 – 5030.
156. Vernoux, T., Kronenberger, J., Grandjean, O., Laufs, P. and Traas, J. (2000). PIN-FORMED 1 regulates cell fate at the periphery of the shoot apical meristem. *Development* **127**: 5157 – 5165.
157. Clayberg, C., Butler, L., Kerr, E., Rick, C. and Robinson, R. (1966). Third list of known genes in the tomato. *Journal of Hereditary*, **57**: 189-196.
158. David-Schwartz, R., Koenig, D. and Sinha, N. (2009). LYRATE is a key regulator of leaflet initiation and lamina outgrowth in tomato. *Plant Cell*, **21**: 3093-3104.
159. Koyama, T., Furutani, M., Tasaka, M. and Ohme-Takagi, M. (2007). TCP transcription factors control the morphology of shoot lateral organs via negative regulation of the expression of boundary-specific genes in *Arabidopsis*. *Plant Cell*, **19**: 473-484.
160. Blein, T., Pautot, V. and Laufs, P. (2013). Combinations of mutations sufficient to alter Arabidopsis leaf dissection. *Plants*, **2**: 230 – 247.
161. Sicard, A., Thamm, A., Marona, C., Lee, Y., Wahl, V., Stinchcombe, J., Wright, S., Kappel, C. and Lenhard, M. (2014). Repeated evolutionary changes of leaf morphology caused by mutations to a homeobox gene. *Current Opinion on Plant Biology*, **24**: 1880 – 1886.

162. Yifhar, T., Pekker, I., Peled, D., Friedlander, G., Pistunov, A., Sabban, M., Wachsman, G., Alvarez, J., Amsellem, Z. and Eshed, Y. (2012). Failure of the tomato trans-acting short interfering RNA program to regulate AUXIN RESPONSE FACTOR3 and ARF4 underlies the wiry leaf syndrome. *Plant Cell*, **24**: 3575 – 3589.
163. Zhou, C., Han, L., Fu, C., Wen, J., Cheng, X., Nakashima, J., Ma, J., Tang, Y., Tan, Y. and Tadege, M. (2013). The trans-acting short interfering RNA3 pathway and no apical meristem antagonistically regulate leaf margin development and lateral organ separation, as revealed by analysis of an argonaute7/lobed leaflet1 mutant in *Medicago truncatula*. *Plant Cell* **25**: 4845 – 4862.
164. Hunter, C., Willmann, M., Wu, G., Yoshikawa, M., de la Luz Gutiérrez-Nava, M. and Poethig, S. (2006). Trans-acting siRNA-mediated repression of ETTIN and ARF4 regulates heteroblasty in *Arabidopsis*. *Development*, **133**: 2973-2981.