

# Original Research Article

## Anti-oxidant and Anti-inflammatory activity of *Moringa oleifera* (Flowers)

Comment [O1]: Remove the brackets

### ABSTRACT

**Objectives:** The aim of the present study was isolate the new compound from ethyl acetate extract of *Moringa oleifera* flowers and identify the anti oxidant and anti inflammatory activities.

Comment [O2]: Should not be italis

Comment [O3]: Identify or determin???

**Methods:** Flavonoids, Biflavonoids are the major constituent of ethyl acetate extract of *Moringa Oleifera* flowers. The extraction and fractionation was carried out from solvents of ethanol, benzene, petroleum ether, diethyl ether and ethyl acetate. The Anti-inflammatory activity of the sample was determined by HRBC membrane stabilization and Albumin denaturation methods. Anti-oxidant activity of the sample was determined by DPPH assay and ABTS method.

Comment [O4]: What happened to the extracts from other solvents listed below?

Comment [O5]: Italics aand 'o' for oleifera should be low case

Comment [O6]: Was it only one sample studied???? If yes, which one????

**Results:** The results of the study, suggest that the sample isolated from the ethyl acetate fraction possesses anti-oxidant activity and anti-inflammatory activity. However, these effects need to be confirmed using in vivo models and clinical trials for its effective utilization as therapeutic agents.

Comment [O7]: Then why did you do other solvents as well?

**Keywords:** *Moringa oleifera*, Antioxidant activity, Anti-inflammatory activity, HRBC method, ABTS assay.

Comment [O8]: It would be good to add on the solvent used.

Comment [O9]: Add on the word flowers.

### 1. INTRODUCTION

The phytochemicals are extracted from all parts of the plant body, but the concentration of these components varies from part to part. Normally, parts known to contain the highest concentration of the principles are preferred to therapeutic purposes and it can either be the leaves, stems, barks, roots, bulks, rhizomes, woods, flowers, fruits or the seeds. The literature review revealed the antimicrobial, analgesic, antifertility and antibacterial, anti-inflammatory, anti-oxidant, purgative and hepatic protective activities of the plant body<sup>[1]</sup>. As oxidative stress plays a central role in liver pathologies and their progression, the use of antioxidants has been

Comment [O10]: Does the plant have a body with liver cells??

Comment [O11]: New paragraph

29 proposed as therapeutic agents, as well as drug coadjuvants, to counteract liver damage<sup>[2]</sup>. Pain  
30 and inflammatory responses in the peripheral and central nervous systems take part in key roles  
31 in the growth and persistence of many pathological pain states<sup>[3]</sup>. A variety of natural  
32 compounds are able to alleviate pain targeting inflammation by reducing the synthesis of  
33 inflammatory mediators, or modulating inflammatory and nociceptive pathways<sup>[4]</sup>.

34 *Moringa oleifera* Lam., a member of the Moringaceae family also known as Drumstick or  
35 Horseradish-tree, is home-grown to the sub- Himalayan regions of India, Pakistan, Bangladesh  
36 and Afghanistan. Due to the importance uses of *M. oleifera* in traditional medicine, many  
37 investigations have previously reported on pharmacological properties such as antifertility, anti-  
38 inflammatory, antispasmodic, and diuretic activities<sup>[5-7]</sup>.

**Comment [O12]:** You did not study the leaves but flowers!!!! How are flowers used from literature???????

**Comment [O13]:** Which part of the *Moringa oleifera* tree??

39

## 40 2. MATERIALS AND METHODS

### 41 2.1 Collection of Flowers

42 Fresh flowers of *Moringa oleifera* were collected from Karaikudi, Sivagangai (Dt), Tamil  
43 Nadu, India, during the month of April and identified by Dr.S.John Britto, Director, The rapinat  
44 Herbarium and Centre for Molecular Systematics (Authentication No. AR003 dated:  
45 05/04/2017). St.Joseph's College (Campus), Tiruchirappalli, Tamil Nadu, India.

46

### 47 2.2 Extraction and fractionation

48 Fresh flowers (3 kg) of *Moringa oleifera* collected were extracted with 90% ethanol. The  
49 combined alcoholic extract was concentrated in vacuo and the aqueous extract was successively  
50 fractionated with petroleum ether (60-80<sup>0</sup>C) (6x250ml), Peroxide free diethyl ether (4x250ml)  
51 and ethyl acetate (8x250ml). Petroleum ether fraction and diethyl ether fraction did not give in  
52 any isolable compounds. Ethyl acetate fraction on concentration yielded a dry powder. The dried  
53 compound was dissolved in DMSO and were used for further studies..

**Comment [O14]:** Which solvents were combined???

**Comment [O15]:** Meaning ????

**Comment [O16]:** How did you concentrate it ???

## 54 3. IN VITRO ANTIOXIDANT ACTIVITY

### 55 3.1 DPPH Assay Method

56 The DPPH free radical is reduced to a corresponding hydrazine, when it reacts with  
57 hydrogen donors. The DPPH radical is purple in colour and upon reaction with hydrogen donor

58 changes to yellow colour. It is a decoloration assay, which is evaluated by the addition of the  
59 antioxidant to a DPPH solution in ethanol or methanol and the decrease in absorbance was  
60 measured at 490nm<sup>[8]</sup>.

**Comment [O17]:** This is not clear. Please rewrite to bring out the meaning

#### 61 Reagents:

##### 62 A. Preparation of 2,2-Diphenyl 1-picryl hydrazyl solution (DPPH, 100µM):

63 22mg of DPPH was accurately weighed and dissolved in 100ml of methanol.  
64 From this stock solution, 18ml was taken and diluted to 100ml using methanol to obtain  
65 100µM DPPH solution.

##### 66 B. Preparation of test solutions:

67 21 mg of ethyl acetate fraction compound was dissolved in distilled DMSO to get  
68 a solution of 21mg/ml concentration. This solution was serially diluted to prepare lower  
69 concentrations.

**Comment [O18]:** What is fraction compound??

**Comment [O19]:** Does this exist????

**Comment [O20]:** Using what?????

70

##### 71 C. Preparation of standard solutions:

72 10mg of ascorbic acid was accurately weighed and dissolved in 1ml of Dimethyl  
73 sulfoxide (DMSO) to obtain 10mg/ml concentrations. These solutions were serially  
74 diluted with DMSO to prepare lower concentrations.

**Comment [O21]:** Find out the use of DMSO and its effects in large doses.

75

##### 76 D. Procedure:

77 The antioxidant activity was carried out in a 96 well micro titre plate. To 200µl  
78 of DPPH solution, 10µl of each of the test sample or the standard solution was added  
79 separately in wells of the micro titre plate. The final concentration of the test and  
80 standard solutions used were 1000, 500, 125 and 31.25 µg/ml. The plates were incubated  
81 at 37°C for 30 minutes and the absorbance of each solution was measured at 490 nm,  
82 using a micro plate reader.


**Comment [O22]:** What are these??

83

S. No	Concentration (µg/ml)	% CTC <sub>50</sub> Cytotoxicity (µg/ml)	IC <sub>50</sub> (µg/ml)
-------	-----------------------	--	--------------------------

84	1	1000	70.54	369.4
85	2	500	63.01	
86	3	125	46.57	
87	4	31.25	29.45	

88 **Table No. 1: Anti-oxidant activity of the compound isolated from the ethyl acetate fraction of flowers of**  
 89 **Moringa oleifera by DPPH assay**



91 How was the

92 cytotoxicity measured in your Experiment? On the graph, what are on the X and Y axes???

93 **Graph No.1: Graphical representation of anti-oxidant activity of the compound is olated from the ethyl acetate**  
 94 **fraction of flowers of Moringa oleifera by DPPH assay**

95 **3.2 ABTS radical scavenging activity:**


97 ABTS radical scavenging activity was performed as described by Re *et al.* (1999) with a  
 98 little modification. 7.0 mM ABTS in 14.7 mM ammonium peroxy-disulphate was prepared in 5.0  
 99 ml distilled water. The mixture was allowed to stand at room temperature for 24 hours. The  
 100 resulting blue green ABTS radical solution was further diluted such that its absorbance is  $0.70 \pm$   
 101  $0.020$  at 734 nm. Various concentrations of the sample solution (in ethanol) (20.0  $\mu$ L) were added  
 102 to 980.0  $\mu$ L of ABTS radical solution and the mixture was incubated in darkness for 10 minutes.  
 103 The decrease in absorbance was read at 734 nm. A test tube containing 20.0  $\mu$ L of ethanol  
 104 processed as described above was served as the control tube. Different concentrations of ascorbic  
 105 acid were used as reference compound.

**Comment [O23]:** What does this stand for???

**Comment [O24]:** Please use the word that will not change the meaning!!!

S. No	Concentration (µg/ml)	% CTC <sub>50</sub> Cytotoxicity(µg/ml)	IC <sub>50</sub> (µg/ml)
1	1000	76.34	211.9
2	500	68.19	
3	125	49.56	
4	31.25	36.78	

Table No. 2 : Anti-oxidant activity of the compound isolated from the ethyl acetate fraction of flowers of *Moringa oleifera* by ABTS assay.



Graph No.2: Graphical representation of radical scavenging activity of the compound isolated from the ethyl acetate fraction of flowers of *Moringa oleifera* by ABTS assay.

Comment [O25]: Same coments as above

#### 4. ANTI- INFLAMMATORY ACTIVITY

##### 4.1 The human red blood cell (HRBC) membrane stabilization method

The anti-inflammatory studies (Gopalkrishnan *et al.*, 2009; Sakat *et al.*, 2010) was adopted with few modifications. The blood was collected from human volunteer who had not intake any NSAIDS for 2 weeks prior to the experiment and mixed with equal volume of Alsever solution (2 % dextrose, 0.8 % sodium citrate, 0.5 % citric acid and 0.42 % NaCl) and centrifuged at 3,000 rpm. The packed cells were washed with isosaline and a 10 % suspension was prepared. Various concentrations of test drug were prepared in mg/ml using distilled water and to each concentration, 1 ml of phosphate buffer, 2 ml hypo saline and 0.5 ml of HRBC suspension were

Comment [O26]: What does it stand for????

122 added and incubated at 37°C for 30 min then centrifuged at 3,000 rpm for 20 minutes. The  
123 hemoglobin content of the supernatant solution was estimated spectrophotometrically at 560 nm.  
124 Diclofenac (100 Jg/ml) was act as a reference standard and its control was prepared by omitting  
125 the test drug. The experiments were performed in triplicates and mean values of the three were  
126 considered. The percentage (%) of HRBC membrane stabilization was calculated.<sup>[9,10]</sup>

Comment [O27]: ?????

$$\text{Percentage of Protection (\%)} = \frac{(100 - \text{OD of drug treated sample})}{\text{OD of Control}} \times 100$$

#### 129 4.2 Albumin denaturation method


130 The method as prescribed (Sakat *et al.*, 2010) was followed with some modifications. The  
131 reaction mixture was consisting of test sample and 1% solution of bovine albumin fraction. pH of  
132 the reaction mixture was adjusted using small amount of HCl. The mixtures were incubated at  
133 37°C for 20 minutes and then heated to 51°C for 20 minutes. After cooling the samples the  
134 turbidity was measured spectrophotometrically at 660 nm. Diclofenac sodium was used as a  
135 standard drug. The experiment was performed in triplicates and the mean value of the three was  
136 considered. Percentage inhibition of protein denaturation was calculated.<sup>[11,12]</sup>

Comment [O28]: To what level???

$$\text{Percentage of inhibition (\%)} = \frac{(\text{OD of Control} - \text{OD of Sample})}{\text{OD of Control}} \times 100$$

S. No	Concentration (µg/ml)	% of Inhibition
		Membrane Stabilization Mean ± S.E.M(S-I)
1	100	29.96 ± 0.41
2	200	39.48 ± 0.59
3	400	48.09 ± 0.61
4	600	58.93 ± 1.40
5	800	62.36 ± 1.86

142  
143 **Table 3: The human red blood cell (HRBC) membrane Stabilization activity of the compound isolated from the**  
144 **ethyl acetate fraction of flowers of *Moringa oleifera***  
145  
146  
147




148

149 *Graph 3: Graphical representation of human red blood cell (HRBC) membrane Stabilization activity of the*  
 150 *compound isolated from the ethyl acetate fraction of flowers of Moringa oleifera*

S. No	Concentration (µg/ml)	% of Inhibition
		Membrane Stabilization Mean ± S.E.M(S-I)
1	100	29.62 ± 0.58
2	200	39.18 ± 0.86
3	400	59.64 ± 0.94
4	600	68.17 ± 1.27
5	800	72.96 ± 1.49

151

152 *Table 4: The Inhibition of Albumin Denaturation activity of the compound isolated from the ethyl acetate*  
 153 *fraction of flowers of Moringa oleifera*



156

157

158

159 *Graph 4: Graphical representation of Inhibition of Albumin Denaturation activity of the compound isolated from*  
 160 *the ethyl acetate fraction of flowers of Moringa oleifera*

161

## 162 5. RESULTS AND DISCUSSION

### 163 5.1 Anti-oxidant activity:

164 The compound isolated from the ethyl acetate fractions of *Moringa oleifera* flowers  
165 exhibited significant anti-oxidant activity when compared with DPPH assay. It is evidenced from  
166 the data presented in Table-1. The result showed the percentage of cytotoxicity for 1000 µg/ml  
167 as 70.54, for 500 µg/ml as 63.01, for 125 µg/ml 46.57, and for 31.25 µg/ml 29.45. It is evident  
168 from the data presented in Table 2 that the sample possesses ABTS assay activity. The result  
169 showed the percentage of cytotoxicity for 1000 µg/ml as 76.34, for 500 µg/ml as 68.19, for 125  
170 µg/ml as 49.56, and for 31.25 µg/ml as 36.78.

**Comment [O29]:** Re-write this sentence

### 171 5.2 Anti-inflammatory activity:

172 The compound isolated from the ethyl acetate fractions of *Moringa oleifera* flowers  
173 exhibited significant anti-inflammatory activity of the human red blood cell (HRBC) membrane  
174 stabilization and the results are presented in Table 3. The result showed the percentage of  
175 inhibition in membrane stabilization for 100 µg/ml as  $29.96 \pm 0.41$ , for 200 µg/ml as  $39.48 \pm$   
176  $0.59$ , for 400 µg/ml as  $48.09 \pm 0.61$ , for 600 µg/ml as  $58.93 \pm 1.40$ , and for 800 µg/ml as  $62.36$   
177  $\pm 1.86$ . The inhibition of Albumin denaturation activity exhibited by the compound are given in  
178 Table 4. The results showed the percentage of inhibition in membrane stabilization for 100  
179 µg/ml as  $29.62 \pm 0.58$ , for 200 µg/ml  $39.18 \pm 0.86$ , for 400 µg/ml as  $59.64 \pm 0.94$ , for 600 µg/ml  
180 as  $68.17 \pm 1.27$ , and for 800 µg/ml as  $72.96 \pm 1.49$ . The anti-inflammatory effect of the  
181 compound isolated from ethyl acetate fraction (test sample) of *Moringa oleifera* may be due to  
182 presence of active constituent flavonoids. The results strongly suggest anti-inflammatory effects  
183 and anti-oxidant effects of the test sample by percentage of inhibitions, which are given in the  
184 Table 1,2,3,4.

**Comment [O30]:** Re-write the whole paragraph so that the reaser can appreciate your results. Are there no other researchers that studied M.oleifera flowers for its anti –oxidant properties???

**Comment [O31]:** Also re-write this sentence.

185

186

187



188 **6. CONCLUSION**

189 In the present study, both DPPH assay and ABTS have showed a highest potential  
190 antioxidant activity and also the human red blood cell (HRBC) membrane stabilization activity  
191 of the test sample. It could be concluded that the compound isolated from the ethyl acetate  
192 fraction of flowers of *Moringa oleifera* is of phytopharmaceutical importance.

**Comment [O32]:** Please re-write and say exactly what you mean.

194 **COMPETING INTERESTS DISCLAIMER:**

195 Authors have declared that no competing interests exist. The products used for this research are  
196 commonly and predominantly use products in our area of research and country. There is absolutely no  
197 conflict of interest between the authors and producers of the products because we do not intend to use  
198 these products as an avenue for any litigation but for the advancement of knowledge. Also, the research  
199 was not funded by the producing company rather it was funded by personal efforts of the authors.

200

201

202 **7. REFERENCES**

**Comment [O33]:** Look up newer references which are in plenty. They will help you to improve your style of writing and communication scientific research.

- 203 1. Solomon S, Muruganatham N and Senthamilselvi M M. Anti-oxidant and  
204 anti-inflammatory activity of *Thespesia populnea* (flowers) Pharmacophore. 2015; 6(1):  
205 53-59
- 206 2. Mukazayire MJ, Allaey V, Duez P, Calderon PB, Stevigny C, and Bigendako, MJ.  
207 “Evaluation of the hepatotoxic and hepatoprotective effect of Rwandese herbal drugs on  
208 *in vivo* (guinea pigs barbiturate-induced sleeping time) and *in vitro* (rat precision-cut liver  
209 slices, PCLS) models,” Experimental and Toxicologic Pathology. 2010; 62(3):289–299.
- 210 3. Watkins LR, Milligan ED, Maier SF, Glial proinflammatory cytokines mediate  
211 exaggerated pain states: implications for clinical pain, Advances in Experimental  
212 Medicine and Biology. 2003; 521: 1–21.
- 213 4. Smejkal K, Selected natural phenolic compounds - potential treatment for peripheral  
214 neuropathy? Ceska Slovenska Farmacie. 2014; 63:55–70.
- 215 5. Fahey JW. *Moringa oleifera*: a review of the medical evidence for its nutritional,  
216 therapeutic, and prophylactic properties. Part 1. Tree for Life Journal. 2005; 1:5.

- 217 6. Shukla S, Mathur R, Prakash A O. Antifertility profile of the aqueous extract of  
218 Moringa oleifera roots. *Journal of Ethnopharmacology*. 1988; 22:51–62.
- 219 7. Caceres A, Saravia A, Rizzo S, Zabala L, De Leon E, Nave F. Pharmacologic properties  
220 of Moringa oleifera. 2: Screening for antispasmodic, anti-inflammatory and diuretic  
221 activity. *Journal of Ethnopharmacology*. 1992; 36:233–237.
- 222 8. Ilavarsan R and Mallika M et al. (2005), “*African Journal of Traditional, Complementary  
223 and Alternative medicines*”. 2005; 2(1):70-85.
- 224 9. Mizushima Y and Kobayashi M. Interaction of anti-inflammatory drugs with serum  
225 proteins especially with some biologically active proteins. *Journal of Pharmacy  
226 and Pharmacology*. 1968; 20:169-173.
- 227 10. Sakat S, Juvekar AR, Gambhire MN. In vitro antioxidant and inflammatory activity of  
228 methanol extract of *Oxalis corniculata* Linn. *International Journal of Pharma and  
229 Pharmacological Sciences*. 2010; 146-155.
- 230 11. Brown JH and Mackey HK. Inhibition of heat-induced denaturation of serum proteins by  
231 mixtures of nonsteroidal anti-inflammatory agents and aminoacids. *Proceedings of the  
232 Society for Experimental Biology and Medicine*, 1968; 128:225-8.
- 233 12. Grant NH, Alburn HE and Kryzanaukas C. Stabilization of serum albumin by  
234 anti-inflammatory drugs. *Biochemical Pharmacology*. 1970; 19:715-22.

235