

Original Research Article

Impact of zinc fertilization on yield, yield attributes and quality parameters of finger millet varieties under rainfed *alfisols* of southern zone, Andhra Pradesh

ABSTRACT

Aims: To study the effect of zinc fertilization through soil and foliar at different stages of finger millet on yield, yield attributes and quality in two major finger millet varieties under rainfed *alfisols* of southern zone, Andhra Pradesh

Study design: Split-plot design

Place and Duration of Study: Wetland farm, S.V Agricultural College, Tirupati and during *kharif* season of 2019 and 2020 (Two seasons)

Methodology: Zinc fertilization to two major finger millet varieties *viz.*, Vakula and Tirumala through soil and foliar application at different crop stages with following treatments *viz.*, Control (No fertilizers and manures); RDF (60 -30-20 kg N-P-K + FYM @ 4 t ha⁻¹); RDF + soil application of ZnSO₄ @ 25 kg ha⁻¹ as basal; RDF+Soil application of chelated-ZnSO₄ @ 5 kg ha⁻¹; RDF+foliar application of 0.2% ZnSO₄ at ear head emergence stage; RDF+foliar application of 0.2% ZnSO₄ at grain filling stage; and RDF+foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stage. The yield, yield attributes and quality parameters *viz.*, protein, zinc and iron content in grains were determined by adopting standard protocols.

Results: The application of zinc significantly improved the yield and quality parameters over control. The foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stage was significantly ($p \leq 0.05$) improved the yield and yield attributes of finger millet over RDF. The grain yield, straw yield, no. of productive tillers per plant, no. of fingers per plant were increased to 57.0%, 83.2%, 44.6% and 51.7%, respectively over RDF *i.e.*, 60-30-20 kg N-P-K + FYM @ 4 t ha⁻¹. The quality parameters namely protein, grain zinc and iron also increased up to 40.7%, 69.5% and 43.2%, respectively over RDF.

Conclusion: Application of zinc sulphate at ear head emergence and grain filling stages enhanced the yield, yield parameters and quality parameters compared to other treatments of tirumala variety under rainfed *alfisols* of southern zone of Andhra Pradesh.

Keywords: Zn fertilization; fortification; grain yield; protein, Recommended fertilizer dose

1. INTRODUCTION

Finger millet (*Eleusine coracana*) is commonly known as "Nutritious millet" owing to its nutritional superiority over many cereal crops (rice, maize and sorghum) in terms of

proteins, minerals, iron, calcium and vitamins. The grains contains about 5 to 8 per cent protein, 65 to 75 per cent carbohydrates, 15 to 20 per cent dietary fiber and 2.5 to 3.5 per cent minerals and it has 30 times more calcium than rice (344 mg/100 g). Finger millet well recognized with their health beneficial effects viz., anti-diabetic, anti-tumorigenic, atherosclerogenic, antioxidant and antimicrobial properties (1).

Finger millet extensively cultivated in the tropical and sub-tropical regions, which accounts for about 85% of total millet production in India. It is an important small millet crop ranked third in cultivated area, production and productivity of 1.19 mha, 1.98 mt and 1661 kg ha⁻¹, respectively (2) and it has the pride of place in having the highest productivity among the millets after sorghum and pearl millet (3). In Andhra Pradesh it covers an area of 31.63 thousand ha with a production and productivity of 35,000 tonnes and 1087 kg ha⁻¹, respectively. The grains have long storability even under ambient conditions and have made them "famine reserves". This aspect is at most important as Indian agriculture suffers from vagaries of monsoon (4). Under increased probability of occurrence of drought and soil fertility degradation, many farmers opted to raise this crop, hence the cultivated area was allocated for this crop has significantly increased over the last decade (5).

Zinc (Zn) is considered the major limiting micronutrient in most of the areas limiting the crop yields. Zinc has role in diverse physiological functions in biological systems. Zinc is typically the second most abundant transition metal in organisms after iron and the only metal represented in all six enzyme classes viz., oxidoreductase, transferases, hydrolases, lyases, isomerases and ligases (6). Further around 200 enzymes responsible for growth, development, immune function and resistance to infections are regulated by zinc in plant system (7). Hence, Zinc insufficiency in soils may cause lower yield or sometimes crop failure and leads to poor accumulation of zinc into grains causing zinc malnutrition in humans (8) (9). Zinc deficiency is 5th leading cause of deaths in the developing world and about 0.8 million people die annually of which 0.45 million are children under the age of five as per WHO reports. The extent of zinc deficiency was 49% in Indian soils (10). The soils of Andhra Pradesh are also deficient in Zn, hence there is a dietary need to increase grain Zn content of finger millet, improving the remobilization of absorbed and accumulated Zn to grain is a research priority. Zn application also reported to increase the grain iron (Fe) concentration in pearl millet (11).

The agronomic bio-fortification is a easier and faster approach to increase grain Zn concentration in finger millet. Several studies revealed that Zn fertilization increased Zn concentration in rice grain from 35 to 141 percent (12), increased from 24 to 48 percent in wheat (13) and up to 72 percent in maize (14). But limited research has done on impact of zinc fertilization on yield, quality and bio-fortification in finger millet. Hence present study was initiated to study the effect of Zn fertilization on yield, yield attributes and grain fortification in finger millet under rainfed condition.

2. MATERIAL AND METHODS

2.1 Description of experimental site

2.1.1 Climate

. Field experiment was carried out at wetland farm, S.V.Agricultural College, Tirupati Andhra Pradesh, India during *kharif* season in the year 2019 and 2020. Geographically located between 13.5° N and 79.5° E with an altitude of 182.9 m above MSL. The region has a semi-arid type climate. During *kharif* 2019, crop received 769.0 mm of rainfall in 43 rainy days with standard week wise mean maximum and minimum temperature ranged from 28.2 to 36.1 °C and 21.3 to 27.6 °C, respectively. The mean sunshine hours and mean evaporation (USWB Class-A Open Pan evaporimeter) ranged from 2.0 to 8.8 hours day⁻¹ and 1.9 to 6.0 mm per day with an average of 4.7 hr day⁻¹ and 4.2 mm per day, respectively.

During *kharif* 2020, crop received 723.8.0 mm of rainfall in 31 rainy days with standard week wise mean maximum and minimum temperature ranged from 28.4 to 36.0°C and 20.8 to 24.7°C, respectively. The mean sunshine hours and mean evaporation (USWB Class-A Open Pan evaporimeter) ranged from 0.2 to 8.2 hours day⁻¹ and 2.1 to 5.0 mm per day with an average of 3.8 hr day⁻¹ and 3.4 mm per day, respectively.

2.1.2 Initial Soil characteristics

Composite soil sample at 0-15 cm depth was collected, processed and analyzed for different physical, chemical properties by following the standard procedures and The soil was sandy clay loam in texture (18.3% clay, 5.5% silt and 76.2% sand), slightly alkaline (7.87) in reaction, non-saline (0.423 dS m⁻¹) in nature. The oxidizable organic carbon was medium (6.5 g kg⁻¹). The available nitrogen was low (213 kg ha⁻¹) and available phosphorus and potassium was in high category (189 and 564 kg ha⁻¹ respectively) whereas, DTPA extractable zinc was sufficient (1.33 g kg⁻¹).

Commented [ma1]: use unit in meters, , for example 0.00 - 0.15 m

2.2 Treatments and Experimental Design

The experiment was laid out in split plot design with two finger millet varieties as main treatments viz., Vakula and Tirumala released by Agricultural Research Station, Perumallapalli and zinc fertilization at different methods and crop stages as sub treatments viz., T₁: Control (No fertilizers and manures), T₂: 60 -30-20 kg N-P-K + FYM @ 4 t ha⁻¹, T₃: T₂ + Soil application of ZnSO₄ @ 25 kg ha⁻¹ as basal, T₄: T₂ + Soil application of chelated zinc sulphate @ 5 kg ha⁻¹, T₅:T₂+Foliar application of 0.2% ZnSO₄ at ear head emergence stage, T₆: T₂ + Foliar application of 0.2% ZnSO₄ at grain filling stage and T₇:T₂+Foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stage. The treatments randomized in split plot design with three replications. The recommended dose of 60 kg N, 30 kg P₂O₅ and 20 kg K₂O ha⁻¹ applied through urea, SSP and MOP, respectively. Fertilizer nitrogen was applied in two equal splits as first half dose at the time of transplanting and second half at 30 DAT and the full dose of FYM @ 4 t ha⁻¹, phosphorus and potassium applied at the time of transplanting.

Commented [ma2]: How many plants represented the sample unit? if possible inform

2.3 Soil and Plant analysis

Post harvest soil samples were collected from each treatment before after harvesting of both the seasons during 2019 and 2020 at 0 to 15 cm. The samples were air-dried at room temperature, pulverized, sieved through a 2-mm sieve. The available zinc DTPA method (15). Weighed 10 g of soil into a 150 ml conical flask, added 20 ml of DTPA extractant (0.005 M DTPA, 0.01 M CaCl₂ and 0.1 M Triethanol amine (TEA) with pH 7.3) and shake the contents on a horizontal shaker for 2 hrs and Filtered the suspension through whatman No.42 filter paper and zinc content was determined in the extractant using Atomic absorption spectrophotometer (model No: Spectra A varian 220).

2.4 Yield and yield attributes

The grain obtained from the net plot area including the grain of the sampled plants was thoroughly sundried to a safe moisture level of 14 per cent, weighed and expressed in kg ha⁻¹. Similarly straw was sun dried to a constant weight and expressed in kg ha⁻¹.

2.5 Estimation of quality parameters

2.5.1 Grain zinc and iron content (mg kg⁻¹)

Di-acid digestion was carried out using a mixture of HNO₃:HClO₄ (9:4) by taking one gram of powdered grain sample in 150 ml conical flask and added 10 ml of di-acid mixture and mixed

by swirling. The contents were placed on hot plate in a digestion chamber. The contents were further evaporated until the volume was reduced to 3 to 5 ml but not to dryness. The completion of digestion was confirmed by white fumes and kept for cooling. Added double distilled water and filter the contents into a 100 ml volumetric flask by using whatman No. 42 filter paper and made upto 100 ml. The filtrate was used for estimating zinc by AAS. Grain zinc (ppm) = [AAS reading x volume made (100 ml)]/ wt of the plant sample (g).

2.5.2 Grain protein content (%)

Estimation of total protein content in seeds of finger millet was done as per the method developed by(16). Weighed 0.5 g grain, grounded with pestle and mortar by adding 10 ml of phosphate buffer. The contents were centrifuged at 3500 rpm for 15 min. The supernatant was used for protein estimation. Aliquot of 0.2 ml of sample extract was pipette out in test tube and made up to 1.0 ml volume. A test tube with 1 ml volume of water was used as a blank. 5 ml of reagent-C was added to all the test tubes including the blank. The contents were mixed well and allowed to stand for 10 min. added 0.5 ml of reagent-D and mixed well, incubated for 30 min at room temperature in dark. The colour intensity was read at 660 nm using spectrophotometer (Model: Genesys 10S UV-VIS). From the standard curve, concentration of protein in different samples was determined and expressed in percentage.

2.6 Statistical Analysis

The experimental data were analyzed statistically by following standard procedure outlined by (17). Significance was tested by comparing 'F' value at 5 per cent level of probability. Treatmental differences that were non-significant were denoted as NS and the data analysed by OPSTAT.

3. RESULTS AND DISCUSSION

3.1 Effect of zinc on yield attributes

The yield attributes among the two varieties (main plots) and the interaction effect were found non-significant. Among the zinc application treatments (sub plots) there is a significant difference. Two years (2019 & 2020) data was presented. The response was almost similar among main, sub plots and interactions in two years of experiments and hence only pooled date are used to highlight the results.

3.1.2 Productive tillers per plant

Number of productive tillers per plant has been tabulated in table 1. From the pooled data, it was noticed that the productive tiller number per plant was significantly influenced by the application of zinc. The treatment RDF + FYM +foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stages (T₇) recorded significantly more number of productive tillers per plant (1.88) which was on par with the treatments RDF + FYM + foliar application of 0.2% ZnSO₄ at ear head emergence stage (T₅) (1.68), RDF + FYM +foliar application of 0.2% ZnSO₄ at grain filling stage (T₆) (1.62) and RDF + FYM +soil application ZnSO₄ @ 25 kg ha⁻¹ (T₃) (1.53) regardless of main plot treatments. The lowest was expressed in (T₁) control (1.17). The main plot treatment ranged from 1.43 to 1.61 and there was no significant difference between the varieties. The interaction effect was noticed as non-significant and ranged from 1.07 to 1.80. The results are in coincidence with (18) and (19). Increased number of productive tillers per plant due to optimum supply of zinc which increases the availability of other nutrients (macro & micro) results in the enhancement of metabolic activities of plant and finally increased the yield.

3.1.2 No. of finger number per plant

The data pertaining to finger number per plant was presented in table 1. From the pooled data, The treatment RDF + FYM +foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stages (T₇) recorded significantly more number of finger number per plant (21.58) which was on par with the treatments RDF + FYM + foliar application of 0.2% ZnSO₄

at ear head emergence stage (T_5) (19.40) and RDF + FYM + foliar application of 0.2% $ZnSO_4$ at grain filling stage (T_6) (18.17) and the lowest was noticed in (T_1) control (12.82) regardless of main plot treatments. There was no significant difference among main plot treatments and interaction effect. The values ranged between 15.82 to 18.35 and 22.17 to 11.00 respectively. Matching results were expressed in the year 2019 and 2020 experiments conducted by (20) and (21) respectively.

3.1.3 Test weight (1000 grains)

The data pertaining to finger number per plant was presented in table 1. From the pooled data, The treatment RDF + FYM + foliar application of 0.2% $ZnSO_4$ at ear head emergence and grain filling stages (T_7) recorded significantly highest test weight (3.05 g) which was on par with RDF + FYM + foliar application of 0.2% $ZnSO_4$ at grain filling stage (T_6) (2.98 g) and RDF + FYM + foliar application of 0.2% $ZnSO_4$ at ear head emergence stage (T_5) (2.95 g). Lowest test weight was registered by absolute control (T_1) (2.72 g). Among the main plot and interaction, the effect was non-significant. The values ranged from 2.86 g to 2.94 g and 2.66 g to 3.06 g respectively. The highest test weight (g) in T_7 treatment was due to foliar spray which helps to rapid absorption of zinc through leaf resulted in better grain filling ability of the crop. The supremacy of yield components is due to combined application of RDF + FYM + $ZnSO_4$ which enhanced the photosynthesis and better translocation of photosynthates from source to sink. The above results of yield components are in coincidence with the experimental results conducted by (22) and (23).

3.1.4 Grain Yield (kg ha⁻¹)

Grain yield is the combination of yield attributing characters viz., number of productive tillers per plant, no. of fingers per plant and test weight (g). From the pooled data, it was revealed that significantly grain yield (2298.27 kg ha⁻¹) was observed in (V2) tirumala variety when compared to vakula variety of finger millet. Among sub plots, the highest grain yield was received by the treatment applied with RDF + FYM + Foliar application of 0.2% $ZnSO_4$ at both ear head emergence and grain filling stages (T_7) (3150.55 kg ha⁻¹). The lowest grain yield of 1452.9 kg ha⁻¹ was registered in absolute control (T_1) compared to other treatments. Regardless of application of Zinc sulphate, incorporation of FYM + RDF recorded 38% higher when compared to absolute control. The increase in grain yield in T_7 was 57% compared to T_2 (RDF + FYM). The interaction effect also shows significant difference and highest was noticed by Tirumala variety (V_2) (3048 kg ha⁻¹) by the foliar spray of 0.2% $ZnSO_4$ at both ear head emergence and grain filling stages (T_7) i.e., (V_2T_7) and the lowest (1434 kg ha⁻¹) was noticed by Vakula variety (V_1) with absolute control (T_1) i.e., V_1T_1 . Similar results were reported by (24) and (25)

3.1.4 Straw yield (kg ha⁻¹)

Application of RDF + FYM + foliar application of 0.2% $ZnSO_4$ at ear head emergence and grain filling stages (T_7) improved the straw yield (7364.58 kg ha⁻¹) of finger millet by 83.2 per cent compared to absolute control (T_1) and 51.5 % compared to T_2 . The main plot and interactions were non-significant and the values are in the range of 5531 to 5423 and 7404 to 3866. kg ha⁻¹ respectively. These findings are in matching with (24) and (19). The highest straw yield (kg ha⁻¹) in T_7 treatment was due to increase in cell division and cell elongation. The lowest straw yield of 3566 kg ha⁻¹ was expressed by T_1 (absolute control). The grain yield was the ultimate end product of many yield contributing attributes, physiological and morphological processes that took place in plants during its life cycle. The Integrated application of organic and inorganic sources showed beneficial effect on physiological process of plant metabolism and growth, thereby resulting in higher grain and straw yield.

3.2 Quality parameters

3.2.1 Protein content in grain (%)

There was significant difference among the zinc application treatments (sub-plots) and the highest protein content (9.63%) in grain was recorded in the treatment receiving RDF + FYM + foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stages (T₇). Lowest grain yield was registered with absolute control (4.49%). Table. 3. The protein content of two varieties (main plots) showed non-significant and the values ranged between 7.90 to 7.56%. The interaction between varieties and zinc application treatments was non-significant (table. 3) on protein content of grain. The values ranged between 9.93% (V₂T₇) to 4.54% (V₁T₁). The results were coincided with the results obtained by (22) and (26). The increase in protein content with zinc application is in agreement with the hypothesis that zinc plays an important role in protein synthesis.

3.3.2 Zinc content in grain (mg kg⁻¹)

Among the two varieties, Tirumala (V₂) recorded significantly highest zinc content (28.68 mg kg⁻¹) in grain. Among the treatments (zinc application) of sub-plot, (T₇) RDF + FYM + foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stages recorded significantly highest zinc content (39.06 mg kg⁻¹) in grain. The lowest was registered by absolute control (T₁) (12.02 mg kg⁻¹). The interaction effect shows non-significant results and values ranged between 11.17 to 40.82 mg kg⁻¹. The results are in coincidence with the experiments conducted during the year 2019 and 2020 by (27) and (28) respectively. Enhancement of zinc content in grain of finger millet by 69.5% compared to RDF + FYM (T₂) was due to foliar application of 0.2% ZnSO₄ at both ear head emergence and grain filling stages. Whereas, the increase of protein content of T₇ by (69.5%) compared to T₂

3.3.3 Iron content in grain (mg kg⁻¹)

The iron content (mg kg⁻¹) in grain was significantly influenced by sub plots (zinc application) and main plot (varieties) and interaction between varieties and zinc application showed non-significant. Highest iron content (151.81 mg kg⁻¹) was registered significantly with Tirumala variety (V₂). Significantly highest iron content (178.92 mg kg⁻¹) in grain was recorded by T₇ (RDF + FYM + foliar application of 0.2% ZnSO₄ at ear head emergence and grain filling stages) which was on par with T₆ (168.91 mg kg⁻¹) (RDF + FYM + foliar application of 0.2% ZnSO₄ at grain filling stage). Both the treatments were significantly superior over other treatments which were tried in this experiment. The results are similar with the field experiments conducted by (29) and (30). The Lowest iron content (114.28 mg kg⁻¹) in grain of finger millet was recorded with absolute control (T₁). The interaction effect ranged from 182.07 to 113.80 mg kg⁻¹ in grains of finger millet. The iron content was increased by (43.2%) compared to RDF + FYM (T₂) whereas the increase in iron content by 55.2% compared to absolute control (T₁).

The foliar application of ZnSO₄ at both ear head emergence and grain filling stages enhanced all grain quality parameters.

Table 1. Yield and yield attributes as influenced by zinc application of finger millet genotypes

Treatments	No. of productive tillers per plant			Finger number per plant			Test weight (1000 grain weight)		
	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
Varieties (V)									
(V ₁) : Vakula	1.97	0.86	1.43	16.81	15.17	15.98	2.95	2.76	2.86
(V ₂) : Tirumala	2.26	0.95	1.61	18.80	17.89	18.35	2.98	2.90	2.94
S.E (m)	0.125	0.044	0.062	1.061	1.015	1.048	0.040	0.014	0.025
C.D (p=0.05)	NS	NS	NS	NS	NS	NS	NS	0.095	NS
Method and stage of Zn application									
Control (no fertilizers and manures)	1.66	0.68	1.17	13.08	12.55	12.82	2.76	2.68	2.72
RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)	1.85	0.75	1.30	14.98	13.45	14.22	2.93	2.77	2.86
RDF+ Soil application of ZnSO ₄ @ 25 kg/ha as basal	2.18	0.87	1.53	17.93	16.73	17.33	2.93	2.85	2.89
RDF+ Soil application of chelated Zinc sulphate @ 5 kg/ha	2.07	0.83	1.47	17.33	15.95	16.63	2.90	2.80	2.85
RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence stage	2.33	0.98	1.68	19.98	18.78	19.40	3.02	2.87	2.95
RDF+ foliar application of 0.2% ZnSO ₄ at grain filling stage	2.51	1.25	1.88	22.40	20.78	21.58	3.14	2.95	3.05
RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence & grain filling stage	2.22	0.97	1.62	18.90	17.45	18.17	3.06	2.89	2.98

S.E (m)	0.200	0.137	0.125	0.876	0.980	0.915	0.050	0.061	0.037
C.D (p=0.05)	NS	NS	0.366	2.571	2.879	2.686	0.148	NS	0.109
Interaction (main x sub)	NS	NS	NS	NS	NS	NS	NS	NS	NS

UNDER PEER REVIEW

Table 2. Grain yield, straw yield as influenced by zinc application in finger millet varieties

Treatments	Grain yield (kg ha ⁻¹)			Straw yield (kg ha ⁻¹)		
	2019	2020	Pooled	2019	2020	Pooled
Varieties (V)						
(V ₁) : Vakula	2481	1980	2230	5922	4923	5423
(V ₂) : Tirumala	2585	2010	2298	6195	4867	5531
S.E (m)	6.1	6.7	6.03	142.7	188.2	146.4
C.D (p=0.05)	40.0	NS	40.2	NS	NS	NS
Method and stage of Zn application						
Control (no fertilizers and manures)	1666	1239	1452	4641	3395	4018
RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)	2183	1823	2003	5262	4454	4858
RDF+ Soil application of ZnSO ₄ @ 25 kg/ha as basal	2413	1990	2201	5516	4662	5089
RDF+ Soil application of chelated Zinc sulphate @ 5 kg/ha	2353	1933	2143	5366	4591	4979
RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence stage	2521	2073	2297	6691	4866	5779
RDF+ foliar application of 0.2% ZnSO ₄ at grain filling stage	3036	2169	2603	6908	5595	6252
RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence & grain filling stage	3560	2740	3150	8025	6704	7364
S.E (m)	76.8	25.7	40.5	335.7	236.8	190.8
C.D (p=0.05)	22.5	75.4	119.0	985.8	695.4	560.2
Interaction (main x sub)	NS	111.8	171.1	NS	NS	NS

Table 3. Grain quality parameters of finger millet as influenced by zinc application

Treatments	Protein (%)			Zinc (mg kg ⁻¹)			Iron (mg kg ⁻¹)			Calcium (%)		
	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled	2019	2020	Pooled
Varieties (V)												
(V ₁) : Vakula	7.70	7.41	7.56	25.10	24.06	24.58	149.38	147.5	148.4	0.27	0.68	0.48
(V ₂) : Tirumala	8.10	7.69	7.90	29.39	27.97	28.68	152.93	150.6	151.8	0.29	0.74	0.51
S.E (m)	0.09	0.053	0.07	0.96	0.72	0.25	2.81	4.6	1.92	0.007	0.17	0.09
C.D (p=0.05)	NS	NS	NS	NS	NS	1.682	NS	NS	NS	NS	NS	NS
Method and stage of Zn application												
Control (no fertilizers and manures)	4.61	4.36	4.49	12.69	11.36	12.02	116.2	114.2	115.2	0.30	0.52	0.41
RDF (60 -30-20 kg N-P-K & FYM @ 4 t/ha)	6.89	6.78	6.84	23.66	22.44	23.05	123.0	126.7	124.8	0.27	0.57	0.42
RDF+ Soil application of ZnSO ₄ @ 25 kg/ha as basal	8.62	8.35	8.48	27.47	25.88	26.68	156.3	154.2	155.3	0.28	0.97	0.63
RDF+ Soil application of chelated Zinc sulphate @ 5 kg/ha	8.34	8.07	8.21	25.13	24.29	24.71	149.1	144.2	146.6	0.30	0.87	0.59
RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence stage	8.16	7.64	7.90	28.36	27.36	27.86	162.7	159.3	161.0	0.27	0.40	0.33
RDF+ foliar application of 0.2% ZnSO ₄ at grain filling stage	8.69	8.37	8.53	33.54	32.55	33.04	169.9	167.9	168.9	0.26	0.78	0.52

RDF+ foliar application of 0.2% ZnSO ₄ at ear head emergence & grain filling stage	9.97	9.30	9.63	39.87	38.25	39.06	180.8	177.0	178.9	0.29	0.85	0.57
S.E (m)	0.334	0.244	0.277	1.617	1.459	1.084	7.859	7.207	5.567	0.020	0.210	0.107
C.D (p=0.05)	0.981	0.715	0.813	4.747	4.284	3.184	23.0	21.16	16.34	NS	NS	NS
Interaction(main x sub)	NS	NS	NS	NS	NS	NS	NS	NS	NS	0.089	NS	NS

UNDER PEER REVIEW

4. CONCLUSION

The results of the present experiment confess that significantly yield attributes, yield and quality parameters of finger millet were recorded highest with the application of RDF+ foliar spray of 0.2% ZnSO₄ at both ear head emergence and grain filling stages. It might be acknowledged that combination of inorganic and organic nutrient sources and addition of external application of zinc through foliar spray proved to be superior over other treatments. So, we can informed that the application of zinc sulphate at both stages enhanced the yield, yield parameters and quality parameters compared to other treatments of tirumala variety under rainfed alfisols of southern zone of Andhra Pradesh.

REFERENCES

1. Chethan S. Nagappa Malleshi G. Finger Millet Polyphenols: Characterization and their Nutraceutical Potential. *American Journal of Food Technology*. 2007;2(7): 2167 – 2179.
2. Ministry of Agriculture, Government of India Indiastat. 2017-18. www.indiaagristat.2017-18.
3. Saravanpandian P, Subramanian S, Paramasivam P, Kumaraswamy K. Organic farming in sustaining soil health. *Agricultural Reviews*. 2005; 26(2): 141 – 147.
4. Shanmugan A. Stanly Joseph Michaelraj PA. Study on millets cultivation in Karur district of Tamilnadu. *International Journal of Management Research & Review*. 2013;3(1):2167-2179.
5. Erenso D, Asfaw A, Taye T, Tesfaye T. Genetic resources, breeding and production of millets in Ethiopia. *Proceedings of an International Conference*, 19-21 September, Bern, Switzerland. 2009;43-56.
6. Auld DS. Zinc coordination sphere in biochemical zinc sites, biometals *Australian soils*. *Australian Journal of Soil Research*. 2001;30:45-53
7. Fischer WC, Black RE. Zinc and the risk for infectious disease. *Annu Rev Nutr*. 2004;24: 255–75.
8. De Valença AW, Bake A, Brouwer ID, Giller KE. Agronomic biofortification of crops to fight hidden hunger in sub-Saharan Africa. *Global Food Secur*. 2017;12:8–14.
9. Liu J, Yang M, Li H, Li D, Shi X, Zhang Y. Genetic processes of iron and zinc accumulation in edible portion of crops and their agro-biofortification: a review. *Am J Agric For*. 2017;5(3):65–72.
10. Singh MV. Micro and Secondary Nutrients and Pollutant Elements Research in India. *Progress Report*, IISS, Bhopal, 2006;1-110.
11. Ramegowda Y, Venkategowda R, Govind G, Hagadur GJ, Udayakumar M, Shankar AG. Effect of Zn application on its uptake, distribution and concentration of Fe and Cu in finger millet (*Eleusine coracana* (L.) Gaertn). *Journal of Plant Nutrition*. 2016; 39(4): 569-580
12. Shivay YS, Kumar D, Prasad R, Ahlawat IPS. Relative yield and zinc uptake by rice from zinc sulphate and zinc oxide coatings onto urea. *Nutr cyc agroecosyst*. 2008;80:181-188.
13. Yang XW, Tian XH, Gale WJ, Cao YM, Lu XC, Zhao AQ. Effect of soil and foliar application on zinc concentration and bioavailability in wheat grain grown on potentially zinc deficient soils. *Cereal Res Commun*. 2011;39: 553-543.
14. Hossain MA, Jahiruddin M, Islam MR, Mian MH. The requirement of zinc for improvement of crop yield and mineral nutrition in maize-mungbean-rice system. *Plant soil*. 2008;306: 13-22.
15. Lindsay WL, Norvell WA. Developments of DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of American Journal*. 1978;42: 421-428

16. Lowry OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. *Journal of Biochemistry*. 1951;193: 265-375.
17. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*, 1985.ICAR, New Delhi
18. Patel NI, Patel BR, Patel CK and Patel FB. Effect of foliar spray of nutrients at different growth stages on pearl millet under Dryland condition. *International Journal of Chemical Studies*. 2019;7(5): 05-09.
19. Logesh Kumar S, Sankaran, VM, Hemalatha M and Suresh S. Effect of different level of fertilizers and foliar nutrition on yield attributes and yield of Pearl millet under rainfed condition. *International Journal of Chemical Studies*. 2020; 8(6): 979-982.
20. Kamal G. Comparative response of finger millet [*Eleusine coracana* (L.) Gaertn] and pearl millet [*Pennisetum glaucum* (L.) R.Br. emend Stuntz] to applied nutrients under semi arid conditions of Rajasthan. M.Sc(Ag.)Thesis submitted to Sri Karan Narendra Agriculture University, Jobner, India. 2019.
21. Vijaya kumar M, Sivakumar M and Tamilselvan N. Effect of Zinc and Iron application on yield attributes, available nutrients status and nutrient uptake of finger millet under rainfed condition. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(5): 3237 – 3246.
22. Akshay G. Effect of zinc biofortification on soil properties, yield, nutrient uptake and quality of *rabi* sorghum. M.Sc(Ag.)Thesis submitted to Vasantnaik Marathwada Krishi Vidyapeeth, Parbhani, India. 2019.
23. Prashantha GM, Prakash SS, Umesh S, Chikkaramappa T, Subbarayappa CT and Ramanamurthy V. Direct and residual effect of zinc and boron on yield and yield attributes of finger millet – groundnut cropping system. *International Journal of Pure and Applied Bioscience*. 2019;7(1): 124 – 134.
24. Ashoka P, Rajakumar GR and Priya P. Response of soil and foliar application of nutrients on yield and nutrient uptake of foxtail millet (*Setaria italica* L.). *International Journal of Chemical Studies*. 2020; 8(4): 2282-2285.
25. Chowdary KA and Patra BC. Effect of Micronutrient application with different sources of NPK on growth and yield of finger millet crop in red laterite zone. *Journal of Agricultural Science and Technology*. 2019;9: 403-416.
26. Sharad KM, Suresh K, Madhavi A and Shanti M. Agronomic biofortification and nutrient uptake of sorghum (*Sorghum bicolor* L.) as influenced by fertilization strategies. *International Journal of Chemical Studies*. 2020; 8(3): 2107-2110.
27. Sharanappa HS, Latha BK, Desai B, Koppalkar G and Ravi MV. Effect of Agronomic Biofortification with Zinc and Iron on Yield and Quality of Pearlmillet [*Pennisetum glaucum* (L.)] Genotypes. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(9): 1312-1321.
28. Ajay Kumar E, Surekha K, Bhanu K, Rekha and Harish Kumar Sharma S. Effect of Various Sources of Zinc and Iron on Grain Yield, Nutrient Uptake and Quality Parameters of Finger Millet (*Eleusine coracana* L.). *International Research Journal of Pure & Applied Chemistry*. 2020; 21(2): 46-55.
29. Amar KJ, Shrivastava S and Vinay A. Response of organic manure, zinc and iron on soil properties, yield and nutrient uptake by Pearl millet crop grown in inceptisol. *International Journal of Pure and Applied Bioscience*. 2018; 6 (1): 426-435.
30. Sravani K. Zinc fortification for grain quality and yield of finger millet (*Eleusine Coracana* (L.) Gaertn). M.Sc(Ag.)Thesis submitted to Acharya N G Ranga Agricultural University, Hyderabad, India. 2018.