

## **Original Research Article**

# **AVAILABLE NUTRIENT STATUS OF SOIL AS INFLUENCED BY COMBINED APPLICATION OF HUMIC ACID AND INORGANIC NITROGEN**

---

### **ABSTRACT**

A field experiment was carried out to study the influence of different levels of humic acid (10, 20, 30 kg ha<sup>-1</sup>) and inorganic N fertilizer viz., 100 % of recommended dose and 75 % of recommended dose on chemical properties of soil under direct sown rice at Agricultural college farm, Bapatla during 2019. The experiment was laid out in RBD with ten treatments replicated thrice with BPT-5204 variety of rice as test crop. Soil samples collected at tillering, panicle initiation and harvest stages of crop were analyzed for chemical properties like available N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, Sulphur and cationic micro-nutrients (Fe, Mn, Zn, Cu). Results indicated that increased availability of N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, Sulphur and cationic micro-nutrients (Fe, Mn, Zn, Cu) were observed with the treatment T<sub>6</sub> which received 100% RDN and HA @ 30 kg ha<sup>-1</sup>.

*Keywords: Humic acid, Direct sown rice, Soil chemical properties*

### **1. INTRODUCTION**

Due to intensive cultivation, tropical soils are low in organic matter and also due to increased use of chemical fertilizers enhanced the degradation of soil organic matter by micro-organisms which resulted in depletion of soil organic carbon. The sustainable soil productivity is the key to crop production in any crop or cropping system. For this, maintenance of soil organic matter at satisfactory levels is necessary under Indian conditions. Humic Substances are considered as the most important constituents of soils. They form the largest fraction of soil organic matter and play the most dominant role in soil conditioning and plant growth. Humic substances improve the crop productivity by improving physical, chemical and biological properties of soil. Physically, humic substances improve soil structure and water holding capacity. Chemically, they serve as adsorption and retention complexes for plant nutrients. Complexed elements remain more available to plant roots (Randhawa and Broadbent, 1965) because complexing shields them from immobilization. Biologically, they improve growth of useful soil micro-organisms by acting as source of nutrients and energy to soil biota. The present study was carried out in a view to observe the influence of humic acid and inorganic nitrogen on chemical properties of soil under direct sown rice.

## 2. MATERIAL AND METHODS

A field experiment was conducted at the Agricultural College Farm, Bapatla of Acharya N.G. Ranga Agricultural University during 2019.

put an image for a better understanding of where in the world the experiment was carried out

put description of the place, climate, soil type, geographic coordinates, etc...

The experiment was laid out in a randomized block design with three replications and ten treatments *viz.*, T<sub>1</sub>- (control for N), T<sub>2</sub> - 100% Recommended Dose of Nitrogen (RDN), T<sub>3</sub> - 100 % RDN + FYM @ 10 t ha<sup>-1</sup>, T<sub>4</sub> - 100 % RDN + Soil application of humic acid @ 10 kg ha<sup>-1</sup>, T<sub>5</sub> - 100 % RDN + Soil application of humic acid @ 20 kg ha<sup>-1</sup>, T<sub>6</sub> -100 % RDN + Soil application of humic acid @ 30 kg ha<sup>-1</sup>, T<sub>7</sub> - 75 % RDN + FYM @ 10 t ha<sup>-1</sup>, T<sub>8</sub> -75 % RDN + Soil application of humic acid @ 10 kg ha<sup>-1</sup>, T<sub>9</sub> -75 % RDN + Soil application of humic acid @ 20 kg ha<sup>-1</sup>, T<sub>10</sub> -75 % RDN + Soil application of humic acid @ 30 kg ha<sup>-1</sup> (NOTE: RDF: 120:60:40 N-P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O (kg ha<sup>-1</sup>) through Urea, SSP and MOP. Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>). The experimental soil was low in available nitrogen (226 kg ha<sup>-1</sup>), medium in available phosphorus (51.88 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>), high in available potassium content (601 kg K<sub>2</sub>O ha<sup>-1</sup>) and sufficient in available sulphur (15 mg kg<sup>-1</sup>) and available cationic micro-nutrients *viz.*, Fe (42.08 mg kg<sup>-1</sup>), Mn (5.65 mg kg<sup>-1</sup>), Zn (2.15 mg kg<sup>-1</sup>), Cu (2.03 mg kg<sup>-1</sup>). Well decomposed farmyard manure @ 10 t ha<sup>-1</sup> was applied to the field as per recommended treatments one week before sowing. Nitrogen @120 kg ha<sup>-1</sup> (100 % RDN) and @ 90 kg ha<sup>-1</sup> (75 % RDN) was applied as per the treatments in the form of urea in three equal splits *i.e.*, 1/3<sup>rd</sup> as basal, 1/3<sup>rd</sup> at active tillering stage and 1/3<sup>rd</sup> at panicle initiation stage. A common dose of phosphorus @ 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in the form of single super phosphate was applied as basal just before sowing. A common dose of 40 kg K<sub>2</sub>O ha<sup>-1</sup> was applied as muriate of potash, in two equal splits as half at basal and half at panicle initiation stage by taking the plot size into consideration. Humic acid @ 10, 20, 30 kg ha<sup>-1</sup> was applied to the soil directly in granular form as basal just before sowing. Soil samples collected at different growth stages of rice were analysed for chemical properties *viz.*, available N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, Sulphur and cationic micro-nutrients (Fe, Mn, Zn, Cu) at initial, tillering, panicle initiation and harvest stages of direct sown rice using standard procedures.

describe the methods used to determine nutrient content

describe how the statistical analysis was performed

## 3. RESULTS AND DISCUSSION

### 3.1 Nitrogen

Data pertaining to available nitrogen content in soil at different growth stages were furnished in Table 1. Perusal of data revealed that application of HA showed a significant increase in available nitrogen content at all the stages of crop growth.

The maximum available nitrogen content (297, 275, 261 kg ha<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>), while the lowest available nitrogen content (234, 227, 226 kg ha<sup>-1</sup>) was recorded with T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively. The treatments T<sub>5</sub>, T<sub>3</sub> and T<sub>10</sub> were on par with the treatment T<sub>6</sub>.

Among the humic acid treatments, increase in available nitrogen content was upto a dose of 30 kg ha<sup>-1</sup>. However, the significant increase was observed only upto 20 kg ha<sup>-1</sup> when it is combined with 100% RDN. Similar significant increase in available nitrogen was observed with increase in N dose in the presence of humic acid or FYM.

The application of humic acid increases CEC of soil and hence ammonia loss becomes reduced. The negative sites due to ionization of carboxylic (COOH) and phenolic (OH) might have improved  $\text{NH}_4^+$  retention, hence reduction in N loss (Stevenson, 1994) and thereby increased the availability of N in soils. Increase in available nitrogen content may also be due to the newly formed quinines from humic acid that inhibited nitrification process and consequently decreased the leaching losses (Flaig, 1964). Usually, urea gets rapidly hydrolysed. On application of humic acid combined with urea, it allows the controlled release of urea and provides a continuity in supply of nitrogen (Dong and Yuan 2009). The increase in availability of nitrogen might be due to N contributed by native nitrogen by the increased microbial activities induced by HA. Independent application of urea could also suffer severe volatilization losses as ammonia not engaged by plants is quickly oxidized. This chemolithoautotrophic oxidation of ammonia to  $\text{NO}_2$  is restricted by  $\text{NH}_3$  availability (Laanbroek and Woldendorp, 1995). The increased N availability might be due to decreased no. of nitrifying micro-organisms (Quraishi and Cornfield, 1973).

**Table 1. Effect of different levels of Humic acid and inorganic nitrogen on Available Nitrogen ( $\text{kg ha}^{-1}$ ) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	234	227	226
T2 : 100% RDN	251	233	229
T3 : 100% RDN + FYM @ $10 \text{ t ha}^{-1}$	281	261	249
T4 : 100% RDN + HA @ $10 \text{ kg ha}^{-1}$	267	246	236
T5 : 100% RDN + HA @ $20 \text{ kg ha}^{-1}$	294	262	257
T6 : 100% RDN + HA @ $30 \text{ kg ha}^{-1}$	297	275	261
T7 : 75% RDN + FYM @ $10 \text{ t ha}^{-1}$	255	231	230
T8 : 75% RDN + HA @ $10 \text{ kg ha}^{-1}$	247	232	224
T9 : 75% RDN + HA @ $20 \text{ kg ha}^{-1}$	258	236	235
T10:75% RDN + HA @ $30 \text{ kg ha}^{-1}$	278	259	240
SEm ( $\pm$ )	8.13	9.10	8.23
CD (P=0.05 %)	24	27	24
CV (%)	5.3	6.4	6.0

(NOTE: Recommended dose of P & K was applied from  $T_1$  to  $T_{10}$ )

### 3.2 Phosphorus

The data pertaining to available Phosphorus content in soil at different growth stages of rice were furnished in Table 2. Perusal of data indicated that available phosphorus was not

significantly influenced by different treatments at any growth stage of rice. However, an increase in available phosphorus content was observed in humic acid applied plots at all three stages of crop growth. The available Phosphorus content ranged from 67.33 to 82.78, 64.64 to 80.09 and 63.40 to 78.24 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively.

Among the treatments, available phosphorus content (105.96, 98.63, 96.78 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) was recorded highest in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>) and the lowest available phosphorus content (90.51, 83.18, 80.09 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively.

**Table 2. Effect of different levels of Humic acid and inorganic nitrogen on Available Phosphorus (kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	67.33	64.64	63.40
T2 : 100% RDN	70.42	67.17	65.87
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	75.52	75.15	75.15
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	71.97	72.30	72.05
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	76.60	78.24	77.00
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	82.78	80.09	78.24
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	71.97	67.73	67.73
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	68.88	65.87	64.02
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	71.97	71.44	70.82
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	73.51	73.29	73.91
SEm (±)	4.30	4.08	3.47
CD (P=0.05 %)	NS	NS	NS
CV (%)	10.2	9.9	8.5

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

Humic acid has the ability to reduce P fixation and solubilize insoluble P compounds and hence increases the P concentration in soil (Sibanda and Young, 1986). Application of humic acid increases the phosphatase activity in soil. Phosphatase enzyme which is responsible for hydrolysis of phosphate esters into inorganic phosphorus might have contributed to increased P availability in the soil (Malcolm and Vaughan, 1979). Humic acid causes slow

and continuous dissolution of phosphate minerals in soil which might have accounted for its increased availability throughout the crop growth (Wang *et al.*, 1995).

### 3.3 Potassium

Perusal of data presented in the Table 3 indicated that soil available potassium did not differ significantly among the treatments at any stage of crop growth. However, available potassium content in soil increased on application of humic acid along with inorganic nitrogen at all the stages of crop growth. The available Potassium content in soil ranged from 700 to 763, 637 to 754 and 610 to 709 kg ha<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively.

Maximum available potassium content (763, 754, 709 kg K<sub>2</sub>O ha<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>), while the lowest available potassium content (700, 637, 610 kg K<sub>2</sub>O ha<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively.

**Table 3. Effect of different levels of Humic acid and inorganic nitrogen on Available Potassium (kg K<sub>2</sub>O ha<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	700	637	610
T2 : 100% RDN	709	655	628
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	745	718	691
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	718	691	655
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	754	745	700
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	763	754	709
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	709	682	637
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	709	646	628
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	718	691	637
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	745	700	682
SEm (±)	24.81	40.26	40.70
CD (P=0.05 %)	NS	NS	NS
CV (%)	5.9	10.1	10.7

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

The addition of compost-HA is suspected that HA-NH<sub>4</sub><sup>+</sup> and -K<sup>+</sup> exchanged to soil -Na<sup>+</sup> and soil-Ca<sup>+2</sup> caused reduced soil salinity. This reduction of Na was because of K<sup>+</sup> replaced

them on the surface of adsorption colloid so the proportion of K was increased. Because the three ions have a similar valence, their exchangeability is determined by the affinity of the cations (Tan, 2003). Increase in soil available potassium may be contributed due to the reduced K fixation as well as release of fixed K by humic acid. The humic compounds are capable of penetrating the inter-micellar spaces of clays and reaches the specific sorption sites for K<sup>+</sup> ions where they might compete for sites with K and increase in K availability in soil (Bharath *et al.*, 2015).

### 3.4 Sulphur

Data presented in the Table 4, revealed that available sulphur in soil was not significantly influenced by different treatments at any stage of crop growth. The available Sulphur content in soil ranged from 15.10 to 16.39, 14.96 to 16.11 and 14.16 to 15.37 mg kg<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively. Application of humic acid combined with inorganic nitrogen increased available sulphur content in soil at all the stages of the crop growth.

**Table 4. Effect of different levels of Humic acid and inorganic nitrogen on Available Sulphur (mg kg<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	15.10	14.96	14.16
T2 : 100% RDN	15.10	15.03	14.22
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	15.91	15.37	14.76
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	16.11	15.84	15.03
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	16.38	16.04	15.10
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	16.39	16.11	15.37
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	15.44	15.30	14.49
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	15.23	15.17	14.43
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	15.50	15.30	14.49
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	16.24	15.97	15.03
SEm (±)	0.54	0.67	0.50
CD (P=0.05 %)	NS	NS	NS
CV (%)	5.9	7.4	5.8

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

Highest available sulphur content (16.39, 16.11, 15.37 mg kg<sup>-1</sup> at maximum tillering, panicle initiation and harvest stages respectively) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>)

<sup>1</sup>) and the lowest available sulphur content (15.10, 14.96, 14.16 mg kg<sup>-1</sup> at maximum tillering, panicle initiation and harvest stages respectively) was recorded in T<sub>1</sub> (control). The increase in available sulphur might be due to mineralization of sulphur containing amino acids during decomposition of organics.

### 3.5 Iron

Data pertaining to the effect of humic acid and inorganic nitrogen on available iron content were presented in the Table 5. It was observed that application of HA did not significantly influence available iron content in soil at any growth stage of rice. The available iron content in soil ranged from 42.45 to 48.68, 41.20 to 46.41 and 35.68 to 43.57 mg kg<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively. Application of humic acid in combination with either 100% or 75% N (inorganic) increased the available iron content in soil over control and 100% RDN alone.

**Table 5. Effect of different levels of Humic acid and inorganic nitrogen on Available Iron (mg kg<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	42.45	41.20	35.68
T2 : 100% RDN	42.89	41.64	38.54
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	43.89	43.13	39.35
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	44.79	43.39	39.62
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	45.79	44.37	41.32
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	48.68	46.41	43.57
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	43.46	42.54	38.70
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	43.21	42.08	38.62
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	44.19	42.77	39.20
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	45.39	43.87	40.71
SEm (±)	2.06	1.80	1.61
CD (P=0.05 %)	NS	NS	NS
CV (%)	8.0	7.2	7.1

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

Among various treatments, highest available iron content (48.68, 46.41, 43.57 mg kg<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>) and the lowest available iron content (42.45, 41.20, 35.68 mg kg<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively.

### 3.6 Manganese

Data pertaining to the effect of humic acid and inorganic nitrogen on available manganese content were presented in the Table 6. It is evident that there is no significant effect of treatments on available manganese content in soil at any of the three stages of crop growth. However, an increase in available manganese content in soil was observed on application of humic acid combined with inorganic nitrogen. The available manganese content in soil ranged from 5.96 to 6.72, 5.80 to 6.64 and 5.54 to 6.35 mg kg<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively.

Highest available manganese content (6.72, 6.64, 6.35 mg kg<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>) and the lowest available manganese content (5.96, 5.80, 5.54 mg kg<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively.

**Table 6. Effect of different levels of Humic acid and inorganic nitrogen on Available Manganese (mg kg<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	5.96	5.80	5.54
T2 : 100% RDN	6.14	5.95	5.86
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	6.65	6.44	6.22
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	6.67	6.53	6.26
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	6.72	6.63	6.28
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	6.72	6.64	6.35
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	6.24	6.16	6.09
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	6.16	6.08	6.03
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	6.27	6.19	6.10
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	6.70	6.59	6.28
SEm (±)	0.20	0.28	0.20
CD (P=0.05 %)	NS	NS	NS
CV (%)	5.5	7.7	5.5

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

### 3.7 Zinc

Data pertaining to the effect of humic acid and inorganic nitrogen on available zinc content were presented in the Table 7. It was observed that available zinc in soil did not differ

significantly among the treatments at any stage of crop growth. It was observed that there is a non-significant increase in available zinc on application of humic acid combined with inorganic N. The available zinc content in soil ranged from 4.30 to 5.81, 3.76 to 5.23 and 3.34 to 4.21 mg kg<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively.

Highest available zinc content (5.81, 5.23, 4.21 mg kg<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>) and the lowest available zinc content (4.30, 3.76, 3.34 mg kg<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively. Increased zinc availability might be attributed due to prevention of formation of immobile and insoluble hydroxides of zinc by humic acid (Stumm and Morgan, 1970).

**Table 7. Effect of different levels of Humic acid and inorganic nitrogen on Available Zinc (mg kg<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
<b>T1 : Control (0N-P-K)</b>	4.30	3.76	3.34
<b>T2 : 100% RDN</b>	4.70	3.92	3.43
<b>T3 : 100% RDN + FYM @ 10 t ha<sup>-1</sup></b>	5.74	4.99	4.05
<b>T4 : 100% RDN + HA @ 10 kg ha<sup>-1</sup></b>	5.75	5.04	4.06
<b>T5 : 100% RDN + HA @ 20 kg ha<sup>-1</sup></b>	5.81	5.21	4.17
<b>T6 : 100% RDN + HA @ 30 kg ha<sup>-1</sup></b>	5.81	5.23	4.21
<b>T7 : 75% RDN + FYM @ 10 t ha<sup>-1</sup></b>	5.20	4.91	4.04
<b>T8 : 75% RDN + HA @ 10 kg ha<sup>-1</sup></b>	4.82	4.47	3.75
<b>T9 : 75% RDN + HA @ 20 kg ha<sup>-1</sup></b>	5.50	4.94	4.04
<b>T10:75% RDN + HA @ 30 kg ha<sup>-1</sup></b>	5.78	5.11	4.17
SEm (±)	0.36	0.35	0.21
CD (P=0.05 %)	NS	NS	NS
CV (%)	11.5	12.7	9.4

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

### 3.8 Copper

On perusal of data furnished in the Table 8, it is evident that application of HA had no significant effect on available copper content in soil at any of the three stages of crop growth. However, application of humic acid combined with inorganic nitrogen increased the copper content in soil over control (T<sub>1</sub>) and 100% RDF alone (T<sub>2</sub>). The available copper content in soil ranged from 2.69 to 3.04, 2.60 to 2.96 and 2.44 to 2.82 mg kg<sup>-1</sup> at tillering, panicle initiation and harvest stages of the crop respectively.

Among different treatments imposed, highest available copper content (3.04, 2.96, 2.82 mg kg<sup>-1</sup>) was recorded in T<sub>6</sub> (100% RDN + HA @ 30 kg ha<sup>-1</sup>) and the lowest available copper content (2.69, 2.60, 2.44 mg kg<sup>-1</sup>) was recorded in T<sub>1</sub> (control) at maximum tillering, panicle initiation and harvest stages respectively.

Humic acid attracts positive ions and forms chelate with micronutrients serving as a reservoir of essential plant nutrients. Humic acid being polyvalent molecule (Schnizter and Khan, 1972; Spostio, 1989) attracts micronutrient cations and release them to the plants. Humic acid might have enhanced the nutrient availability by breakdown of minerals in two ways. Humic acids might have attacked minerals and decomposed them thereby releasing them from molecular state to adsorbed state or the HA formed stable organo-mineral complexes of ions such as HA-Fe<sup>+2</sup>, HA-Mn<sup>+2</sup>, HA-Zn<sup>+2</sup>, HA-Cu<sup>+2</sup> which become easily available to the plant (Brady, 1996).

**Table 8. Effect of different levels of Humic acid and inorganic nitrogen on Available Copper (mg kg<sup>-1</sup>) in soil at different growth stages of direct sown rice**

Treatment	Maximum Tillering stage	Panicle Initiation stage	Harvest stage
T1 : Control (0N-P-K)	2.69	2.60	2.44
T2 : 100% RDN	2.73	2.66	2.50
T3 : 100% RDN + FYM @ 10 t ha <sup>-1</sup>	2.97	2.87	2.68
T4 : 100% RDN + HA @ 10 kg ha <sup>-1</sup>	2.97	2.88	2.70
T5 : 100% RDN + HA @ 20 kg ha <sup>-1</sup>	3.02	2.94	2.81
T6 : 100% RDN + HA @ 30 kg ha <sup>-1</sup>	3.04	2.96	2.82
T7 : 75% RDN + FYM @ 10 t ha <sup>-1</sup>	2.84	2.71	2.61
T8 : 75% RDN + HA @ 10 kg ha <sup>-1</sup>	2.78	2.68	2.55
T9 : 75% RDN + HA @ 20 kg ha <sup>-1</sup>	2.89	2.81	2.63
T10:75% RDN + HA @ 30 kg ha <sup>-1</sup>	3.00	2.89	2.73
SEm (±)	0.11	0.17	0.18
CD (P=0.05 %)	NS	NS	NS
CV (%)	6.8	10.7	12.1

(NOTE: Recommended dose of P & K was applied from T<sub>1</sub> to T<sub>10</sub>)

#### 4. CONCLUSION

The treatment involving application of humic acid @ 30 kg ha<sup>-1</sup> combined with 100% RDN (T<sub>6</sub>) recorded significantly highest available nitrogen content and the treatments T<sub>5</sub>, T<sub>3</sub> and T<sub>10</sub> were on par with T<sub>6</sub>, whereas, the available P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, Sulphur and cationic micro-nutrients (Fe, Mn, Zn, Cu) were not influenced at any stage of crop growth with the imposed treatments.

it is not possible to do this information since for all nutrients the statistic presented is not significant

#### REFERENCES

- Bharath, T. and Madhavi, K. 2015. Nutrient uptake of sorghum crop influenced by top dressing and foliar application of organic nutrient sources. *Indian Journal of Applied Research*. 94: 386-396.
- Brady, N. C and Weil, R. R. 1996. *The nature and properties of soil*. Upper saddle river, 11th Edition, Prentice hall, Inc.
- Dong, L and Yuan, H. 2009. Nitrogen incorporation into lignite humic acids during microbial degradation. *Geomicrobiology Journal*. 26: 484–490.
- Flaig, W. 1964. Effects of microorganisms on the transformation of lignin to humic substances. *Geochem Cosmochimica Acta*. 28: 1523.
- Laanbroek, H.J and Woldendorp, J.W. 1995. Activity of chemolithotrophic nitrifying bacteria under stress in natural soils. In: J. G. Jones (ed.). *Advances in Microbial Ecology*. Plenum Press, New York, USA. pp. 275–304.
- Malcolm, R.E and Vaughan, D. 1979. Humic substances and phosphatase activities in plant tissues. *Soil Biology and Biochemistry*. 11: 253- 259.
- Quraishi, M. S. I and Cornfield, A. H. 1973. Incubation study of nitrogen mineralisation and nitrification in relation to soil pH and level of copper (II) addition. *Environmental Pollution*. 4: 159.
- Schnitzer, M and Khan, S.U. 1972. *Humic substances in the soil environment*. Marcel Dekker, New York.
- Sibanda, H. M and Young, S. D. 1986. Competitive adsorption of humus acids and phosphate on goethite, gibbsite, and two tropical soils. *Journal of Soil Science*. 37: 197–204.
- Sposito, G. 1989. *The chemistry of soils*. Oxford Univ. press, Inc, Oxford, Newyork.
- Stevenson, F.J. 1994. *Humus Chemistry: Genesis, Composition, Reaction*. 2<sup>nd</sup> Ed. John Wiley & Sons, Inc. New York. p. 36.
- Stumm, W and Morgan, J. J. 1970. *Aquatic chemistry. An introduction emphasising chemical equilibrium in natural waters*. Wiley, Interscience, London.
- Tan, K. H. 2003. *Humic Matter in Soil and the Environment. Principles and Controversies*. University of Georgia. Athens, Georgia, U.S.A.

Wang, X.J., Wang, Z.Q and Li, S.G. 1995. The effect of humic acids on the availability of phosphorus fertilizers in alkaline soils. *Soil Use and Management*. 11: 99-12.

UNDER PEER REVIEW