

## PREPARATION OF *BOERHAAVIA DIFFUSA* MEDIATED SELENIUM BASED MOUTHWASH-A COMPARATIVE MICROBIAL AND CYTOTOXIC EFFECT

**Running title:** A comparative study on microbial and cytotoxic effect of *boerhaaviadiffusa* based mouthwash

### ABSTRACT

**Introduction:** The extract of *Boerhaaviadiffusa* root, a plant used in Indian traditional medicine, has significant immunomodulatory **Potential**. Selenium, a nutrient element that has a massive function in biological systems, is one of the **the** interesting compounds to integrate with antibacterial agents. Recently several studies have pointed out the ability of selenium nanoparticles to exhibit anticancer, antioxidant, antibacterial and antibiofilm properties.

**Aim:** To analyze the antimicrobial and cytotoxic activity of *Boerhaaviadiffusa* mediated selenium nanoparticles based mouthwash.

**Materials and method:** 1g of *Boerhaaviadiffusa* was added in 100 ml of distilled water. It was boiled; the plant extract was filtered using Whatman's no.1 filter paper. In 250 ml conical flask, 60 ml of 20 millimolar sodium selenite was prepared and 40 ml of the filtered plant extract was mixed. This flask was kept in a magnetic stirrer. The nanoparticle solution was centrifuged at 8000rpm to prepare nanoparticle pellets. The nanoparticle pellet was dried in a hot air oven at 80 degree celsius. The dried powder was sent for **Characterization**. A mouthwash is prepared.

The prepared mouthwash was tested for cytotoxic activity by brine shrimp lethality assay and antimicrobial activity evaluated **the** zone of inhibition of agar well diffusion method.

**Results and Discussion:** It was proved that cytotoxic activity of selenium bound mouthwash was less compared to the commercial mouthwash. The antibacterial activity of the selenium

Comment [VA1]: potential

Comment [VA2]: delete

Comment [VA3]: characterization

Comment [VA4]: by the

bound mouthwash against *C. albicans* and *S. mutans* was significant when compared to the standard antimicrobial agent.

Comment [VA5]: Write full form

**Conclusion:** Based on the results, this mouthwash has the required qualities to be commercially used. Therefore, further studies can be done to prove that this mouthwash can be used commercially.

**Keywords:** Mouthwash, antimicrobial activity, selenium-based, *B. diffusa*, nanoparticles

Comment [VA6]: Full form

## INTRODUCTION

Plant extracts have been widely evaluated for possible immunomodulatory properties. The ethanolic extract of *Boerhaavia diffusa* root, a plant used in Indian traditional medicine, has significant immunomodulatory potential. (1) High incidence of bacterial infections and antibiotic resistance as a growing problem has urged the need for novel antibacterial agents. (2) *Boerhaavia diffusa* roots (Punarnavamool) are in use since the beginning of the Ayurveda era for various therapeutic benefits. Root extract of this plant induces strong systemic resistance in susceptible host plant. (3) *Boerhaavia diffusa* belonging to family Nyctaginaceae has a wide distribution, occurring on major parts of the globe. It is used as traditional medicine by indigenous people of many countries in the world for its protective role against inflammation, prostatic hyperplasia, diabetes, cancer, gastrointestinal problems, arthritis etc. The whole plant contains numerous bioactive compounds which are responsible for its pharmacological activities. Experiments are being done to evaluate full potential of the plant. (4) *Boerhaavia diffusa* contains a large number of phytoconstituents namely, flavonoids, alkaloids, steroids, triterpenoids, lipids, lignins, carbohydrates, proteins, glycoproteins, punarnavine, boeravinone A–F, hypoxanthine 9- $\beta$ -arabinofuranoside, ursolic acid, punarnavoside, punarnavoside and liri dendrin. (5)

Cytotoxic agents are known as all the elements that are toxic to the cells, which include the factors that prevent their growth and sometimes cause death, and are also used to treat certain disorders. Chemical and biological substances or physical agents can cause cytotoxicity by affecting the cells in varying degrees. Antimicrobial activity refers to the process of killing or inhibiting the disease causing microbes. Various antimicrobial agents are used for this purpose. Antimicrobial may be anti-bacterial, anti-fungal or antiviral. In this study the antibacterial and antifungal properties have been analysed.

In previous research to find the microbial activity of selenium particles, resistance of *Escherichia coli* and *Staphylococcus aureus* was compared in the presence of individual Nano and Bio counterparts as well as the nanohybrid system. Upon interaction of SeNPs with Lysozyme, the nanohybrid system efficiently enhanced the antibacterial activity compared to the protein. Therefore, it was concluded that SeNPs play an important role in inhibition of bacterial growth at very low concentrations of protein. (6). Selenium, a nutrient element that has a massive function in biological systems, is one of the interesting compounds to integrate with antibacterial agents. Recently several studies have pointed out the ability of selenium nanoparticles to exhibit anticancer, antioxidant, antibacterial and antibiofilm properties. Oral mucositis (OM) is a complication of high-dose chemotherapy (HDC) followed by hematopoietic SCT (HSCT) with few effective treatments. Selenium has a cytoprotective role via the glutathione peroxidase (Glu.Px) enzyme and prevents chemotherapy-induced toxicities. It is involved in several key metabolic activities through selenoproteins, which are essential for protection against oxidative damage. In other words, selenium is a cofactor for glutathione peroxidase (Glu.Px), an endogenous enzyme system, which is able to scavenge free radicals. Some animal studies have shown that adequate supplementation of selenium could produce cytoprotective effects and anti-ulcer activity (7). Mouthwashes (MWs) have been particularly well accepted by individuals due to their ease of use. It is an effective method for delivery of antimicrobial agents thus preventing bacterial adhesion, colonization, and metabolism. However emergence of bacterial resistance to such agents has become a common phenomenon, which is represents a major problem. This has encouraged the development of alternative strategies to tackle drug-resistance problems. (8) Our team has extensive knowledge

Comment [VA7]: italic

Comment [VA8]: Italic

Comment [VA9]: delete

Comment [VA10]: Hematopoietic stem-cell transplantation

Comment [VA11]: delete

and research experience that has translate into high quality publications(9–20),(21–25).(26)(27)(28)

In the present study the cytotoxic activity and microbial activity of the *B.diffusa* mediated selenium based mouthwash was analysed by comparing with the commercial mouthwash and also the standard antimicrobial agents.This study could have also analyzed various other characters such as antioxidant, anti inflammatory activities etc.

Comment [VA12]: analyzed

## MATERIALS AND METHOD

### Extract preparation

1g of stem of *Boerhaviadiffusa* was added in 100ml of distilled water. It was boiled for 10 - 15 minutes at 70 degree celsius. After boiling, the plant extract was filtered using Whatman's no.1 filter paper. In 250 ml conical flask, 60ml of 20millimolar sodium selenite was prepared and 40ml of the filtered plant extract was mixed. This flask was kept in a magnetic stirrer. The synthesised nanoparticles were preliminarily analysed by using UV visible spectrophotometer. The nanoparticle solution was centrifuged at 8000rpm to prepare nanoparticle pellets. The nanoparticle pellet was dried in a hot air oven at 80 degree celsius. The dried powder was sent for Characterization.A mouthwash is prepared.(Figure 1)

Comment [VA13]: analyzed

### MOUTHWASH PREPARATION

0.3gsucrose,0.001g sodium benzoate,0.01g of sodium lauryl sulphate Dissolved in 10ml distilled water. To that nanoparticle sample 600 microlitre was added.Andflavouring agent peppermint oil was added- 50microlitres

Comment [VA14]: dissolved

### BRINE SHRIMP LETHALITY ASSAY:Cytotoxicity test

Salt water preparation :

2g of iodine free salt was weighed and dissolved in 200ml of distilled water. 6 well ELISA plates were taken and 10-12 ml of saline water was filled. To that 10 nauplii were slowly added to each well (20µL,40 µL,60 µL,80 µL,control). Then the nanoparticles were added according to the concentration level. The plates were incubated for 24 hours.

After 24 hours, the ELISA plates were observed and noted for number of live nauplii present and calculated by using following formula,

$$\text{number of dead nauplii} / (\text{number of dead nauplii} + \text{number of live nauplii}) \times 100$$

### ANTIMICROBIAL TEST

Antimicrobial activity of respective nanoparticles against the strain *Staphylococcus aureus*, *E. faecalis*, *S. mutans* and *C. albicans* MHA agar was utilized for this activity to determine the zone of inhibition. Muller hinton agar was prepared and sterilized for 45 minutes at 120 lbs. Media poured into the sterilized plates and let them stabilize for solidification. The wells were cut using the well cutter and the test organisms were swabbed. The nanoparticles with different concentrations were loaded and the plates were incubated for 24 hours at 37 ° C. After the incubation time the zone of inhibition was measured.



Figure 1: extract preparation and mouthwash preparation of *B. diffusa* mediated selenium nanoparticles based mouthwash.

## RESULTS

The cytotoxic activity of the selenium based mouthwash was tabulated. The viability of the Nauplii was analysed for various selenium nanoparticles concentrations synthesised from mouthwash aided by *B. diffusa* (figure 2). The nauplii in the well was noticed after 24 hours that 80% of Nauplii were alive in a minimum concentration of 5 $\mu$ L and 10 $\mu$ L. 90% of the Nauplii were alive at 20 $\mu$ L and 40  $\mu$ L concentration. At 80 $\mu$ L of concentrations 80% of the nauplii were alive. Whereas 100% of the nauplii were alive as shown by the control. Thus, according to other research, the increase in concentration enhanced cytotoxicity. In order to study the exact mechanism of action, more precise studies of cytotoxicity need to be performed. In the present analysis the cytotoxicity percentage was higher at 20 $\mu$ L and 40 $\mu$ L concentrations. (Table 1)(graph 1) For comparing the selenium based mouthwash, commercial mouthwash was also analysed. It was noticed after 24 hours that 90% of the Nauplii were alive in the minimum concentration of 5 $\mu$ L and 70 % of the Nauplii were alive at 10 $\mu$ L ,20 $\mu$ L and 40 $\mu$ L. Only 50% of the nauplii were alive at 80 $\mu$ L concentration of the commercial mouthwash. Also at control 100% of the nauplii were alive. (table 2)(graph 2) Therefore as the concentration increases the number of nauplii decreases proving it to be highly cytotoxic whereas the selenium bound was less cytotoxic comparatively. The results were statistically analysed and was insignificant. p value was 0.226. (p value < 0.05)

Comment [VA15]: analyzed


### Cytotoxic activity

concentration	5 $\mu$ L	10 $\mu$ L	20 $\mu$ L	40 $\mu$ L	80 $\mu$ L	control
No of nauplii	8	8	9	9	8	10


Table 1: Brine shrimp lethality assay for different concentrations of *B. diffusa* mediated selenium based mouthwash.

concentration	5 $\mu$ L	10 $\mu$ L	20 $\mu$ L	40 $\mu$ L	80 $\mu$ L	control
No of nauplii	9	7	7	7	5	10

Table 2: Brine shrimp lethality assay for different concentrations of commercial mouthwash observed after 24 hours



Graph 1: The bar graphs represents the cytotoxic activity of the commercial mouthwash. X axis depicts the number of live nauplii in percentage. Y axis represents the concentration of commercial mouthwash added. Blue represents day 1 and orange represents day 2.



Graph 2 :The bar graphs represent the cytotoxic activity of the selenium based mouthwash.




Figure 2:Brine shrimp lethality assay comparison ELISA plate well with different concentrations of selenium-based mouthwash and commercial mouthwash Observed for presence of live nauplii after 24 hours.

#### Antimicrobial activity




Table 3 demonstrates inhibition of bacterial growth at varying concentrations of biosynthesized selenium nanoparticles based mouthwash. The antimicrobial activity was evaluated based on their zone of inhibitions and the results were compared with the standard antibacterial agent. The antibacterial activity against *S. aureus* showed a zone of inhibition of 12mm at the concentration of 25µl. The antibacterial activity against *S. aureus* showed a zone of inhibition of 18mm at the concentration of 50µL. Antimicrobial activity against *S. aureus* showed a zone of inhibition of 20mm at the concentration of 100µL. The standard antibacterial agent had 20mm of zone of inhibition. Therefore comparatively the antimicrobial activity of selenium based mouthwash is moderate against *S. aureus*. The antimicrobial activity of selenium nanoparticles based mouthwash against *S. mutans* showed a zone of inhibition 25mm at concentrations of 25µL, 20mm at 50µl and 22mm at concentration of 100µl. The standard antimicrobial agent has zone of inhibition of 18mm. Therefore this mouthwash when compared to the standard shows high antimicrobial activity. The antimicrobial activity of selenium bases mouthwash against *E. faecalis* showed a zone of inhibition of 10mm at the concentration of 5µl. The antimicrobial activity of selenium nanoparticles based mouthwash against *E. faecalis* showed a zone of inhibition of 12mm at the concentration of 50µl. The antimicrobial activity against *E. faecalis* showed a zone of inhibition of 15mm at the concentration of 100µl. When compared to the standard antimicrobial agent it is observed that this mouthwash has low level of antimicrobial activity against *E. faecalis*. The antimicrobial activity of selenium bases mouthwash against *C. albicans* showed a zone of inhibition of 15mm at the concentration of 5µl. The antimicrobial activity of selenium nanoparticles based mouthwash against *C. albicans* showed a zone of inhibition of 18mm at the concentration of 50µl. The antimicrobial activity against *C. albicans* showed a zone of inhibition of 21mm at the concentration of 100µl. (figure 3) When compared to the standard antimicrobial agent that is 13mm it is observed that this mouthwash has high level of antimicrobial activity against *C. albicans*. (table3) Therefore the antimicrobial activity of the *B. diffusa* mediated selenium nanoparticles based mouthwash was high against *C. albicans* and *S. mutans*. (graph 3).

*B diffusa* mediated selenium particles -microbial effect

	25µL	50µL	100µL	AB
--	------	------	-------	----


C albicans	15	18	21	13
E.faecalis	10	12	15	32
S aureus	12	18	20	20
S mutans	25	20	22	18

Table 3:Shows zone of inhibition of various oral pathogens at different concentrations.



Graph 3: The graph represents the antimicrobial activity of the selenium nanoparticles based mouthwash against C.albicans, E. Faecalis, S.aureus and S.mutans in different concentrations and is compared with the standard antimicrobial agent. X axis represents the concentration of the selenium nanoparticles based mouthwash. Y axis represents the percentage of inhibition.

C.albicans S.mutans



E. faecalis S. aureus

Figure 3: Antimicrobial activity of B. diffusa mediated selenium nanoparticles based mouthwash.

## DISCUSSION

In the analysis of toxicity, the brine shrimp lethality assay is an important test that provides us knowledge on the cytotoxic effect of a bioactive compound on cells (29)(30)(31)(32).The viability of the Nauplii was analysed for various selenium nanoparticles concentrations synthesised from mouthwash aided by *B diffusa*(figure 2).The antimicrobial activity of selenium bound mouthwash that were synthesized with *Boerhaaviadiffusa*,showed various zones of inhibition at different concentrations.(33)(34)(35)(36)(37)(38).

Comment [VA16]: analyzed

In previous articles, it has been observed that the cytotoxic activity of selenium particles was high which is not similar to the present study. Also The results of the previous study indicate that Selenium nanoparticles can be potentially used as antimicrobial and antioxidant agents similar to the present study(39)Many studies were done to analyze the cytotoxic activity and microbial activity of selenium particles but not many articles were published on selenium based mouthwash.(40).Brine Shrimp Lethality was done to check the cytotoxicity of nanoparticles and in previous articles it was found that there was increase of cytotoxic level with increased concentration of the administered nanoparticles which is unlike the present study(41).In a study done by vyas,the selenium nanoparticles were synthesised using garlic and he concluded that its antioxidant potential was high.(42) The present study only concentrates on cytotoxic activity and microbial activity which is a limitation of this study.Clove and cinnamon mediated selenium nanoparticles had high antimicrobial activity against *C.albicans* which shows similar results as that of present study.(43)Unlike the present study, the zone of inhibition was found to be high mainly for *S. aureus and E. faecalis*.(44)In this study mainly for *S.mutans* and *C.albicans* the zone of inhibition were found to high.Similar to this study ,The results of a article done previously showed that there was a lower toxicity rate of SeNPs.(45)

Only antimicrobial and cytotoxic activity of the mouthwash have been analyzed.Further more studies must be done to gain knowledge about the different characteristics of the *Boerhaaviadiffusamediated* selenium particles based mouthwash.Errors may occur while counting the live nauplii as well as in measuring the zone of inhibition of various microbes.

## CONCLUSION

Within the limitations of the study we can conclude that *B.diffusamediated* selenium nanoparticles based mouthwash had low cytotoxic potential and high antimicrobial

activity. Therefore it is best to be used commercially. Further studies can be done to prove various other characteristics of this mouthwash to use this mouthwash commercially.

**NOTE:**

The study highlights the efficacy of "Ayurveda" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

**REFERENCES**

1. Hussain AG, Prof DrNormah, Hussin K. Nature's Medicine: A collection of Medicinal Plants from Malaysia's Rainforest. Landskap Malaysia; 323 p.
2. Ghorbanpour M, Bhargava P, Varma A, Choudhary DK. Biogenic Nano-Particles and their Use in Agro-ecosystems. Springer Nature; 2020. 606 p.
3. Dubey NK. Plants as a Source of Natural Antioxidants. CABI; 2014. 317 p.
4. Kaur H. Boerhaavia Diffusa: Bioactive Compounds and Pharmacological Activities. Biomedical and Pharmacology Journal. 2019 Dec 28;12(4):1675–82.
5. Kaur M, Goel RK. Anti-Convulsant Activity of Boerhaaviadiffusa: Plausible Role of Calcium Channel Antagonism. Evid Based Complement Alternat Med. 2011 Jan 9;2011:310420.
6. Vahdati M, T TM. Synthesis and Characterization of Selenium Nanoparticles-Lysozyme Nanohybrid System with Synergistic Antibacterial Properties. Sci Rep [Internet]. 2020 Jan 16 [cited 2021 Feb 12];10(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/31949299/>
7. Jahangard-Rafsanjani Z, Gholami K, Hadjibabaie M, Shamshiri AR, Alimoghadam K, Sarayani A, et al. The efficacy of selenium in prevention of oral mucositis in patients undergoing hematopoietic SCT: a randomized clinical trial [Internet]. Vol. 48, Bone Marrow Transplantation. 2013. p. 832–6. Available from: <http://dx.doi.org/10.1038/bmt.2012.250>
8. Elgamily H, Mosallam O, El-Sayed H, Mosallam R. Antibacterial effectiveness of probiotic-based experimental mouthwash against cariogenic pathogen: An in vitro study

[Internet]. Vol. 12, European Journal of Dentistry. 2018. p. 007–14. Available from: [http://dx.doi.org/10.4103/ejd.ejd\\_253\\_17](http://dx.doi.org/10.4103/ejd.ejd_253_17)

9. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species [Internet]. Vol. 94, Archives of Oral Biology. 2018. p. 93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>
10. VijayashreePriyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol*. 2019 Dec;90(12):1441–8.
11. Paramasivam A, VijayashreePriyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res*. 2020 Feb;43(2):153–4.
12. VijayashreePriyadharsini J, SmilineGirija AS, Paramasivam A. An insight into the emergence of *Acinetobacterbaumannii* as an oro-dental pathogen and its drug resistance gene profile - An in silico approach. *Heliyon*. 2018 Dec;4(12):e01051.
13. Paramasivam A, VijayashreePriyadharsini J. Novel insights into m6A modification in circular RNA and implications for immunity. *Cell Mollimmunol*. 2020 Jun;17(6):668–9.
14. Paramasivam A, Priyadharsini JV, Raghunandhakumar S. Implications of m6A modification in autoimmune disorders. *Cell Mollimmunol*. 2020 May;17(5):550–1.
15. Girija ASS, Shankar EM, Larsson M. Could SARS-CoV-2-Induced Hyperinflammation Magnify the Severity of Coronavirus Disease (CoViD-19) Leading to Acute Respiratory Distress Syndrome? *Front Immunol*. 2020 May 27;11:1206.
16. Jayaseelan VP, Arumugam P. Exosomal microRNAs as a promising theragnostic tool for essential hypertension. *Hypertens Res*. 2020 Jan;43(1):74–5.
17. Ushanthika T, SmilineGirija AS, Paramasivam A, Priyadharsini JV. An in silico approach towards identification of virulence factors in red complex pathogens targeted by reserpine. *Nat Prod Res*. 2021 Jun;35(11):1893–8.
18. Ramalingam AK, Selvi SGA, Jayaseelan VP. Targeting prolyltripectidyl peptidase from *Porphyromonasgingivalis* with the bioactive compounds from *Rosmarinusofficinalis*. *Asian Biomed*. 2019 Oct 1;13(5):197–203.
19. Kumar SP, Girija ASS, Priyadharsini JV. Targeting NM23-H1-mediated inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *Ganoderma lucidum*: A computational study. *pharmaceutical-sciences* [Internet]. 2020;82(2). Available from: <https://www.ijpsonline.com/articles/targeting-nm23h1mediated-inhibition-of-tumour-metastasis-in-viral-hepatitis-with-bioactive-compounds-from-ganoderma-lucidum-a-comp-3883.html>

20. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting Epstein-Barr virus nuclear antigen 1 (EBNA-1) with Murrayakoengii bio-compounds: An in-silico approach. *Acta Virol.* 2020;64(1):93–9.
21. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent.* 2021 May;31(3):436–41.
22. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent.* 2021 Mar;31(2):285–6.
23. Barma MD, Muthupandiyani I, Samuel SR, Amaechi BT. Inhibition of Streptococcus mutans, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol.* 2021 Jun;126:105132.
24. Teja KV, Ramesh S. Is a filled lateral canal - A sign of superiority? *J Dent Sci.* 2020 Dec;15(4):562–3.
25. Reddy P, Krithikadatta J, Srinivasan V, Raghu S, Velumurugan N. Dental Caries Profile and Associated Risk Factors Among Adolescent School Children in an Urban South-Indian City. *Oral Health Prev Dent.* 2020 Apr 1;18(1):379–86.
26. Jayaseelan VP, Paramasivam A. Emerging role of NET inhibitors in cardiovascular diseases. *Hypertens Res.* 2020 Dec;43(12):1459–61.
27. IswaryaJaisankar A, SmilineGirija AS, Gunasekaran S, VijayashreePriyadharsini J. Molecular characterisation of csgA gene among ESBL strains of A. baumannii and targeting with essential oil compounds from Azadirachtaindica. *Journal of King Saud University - Science.* 2020 Dec 1;32(8):3380–7.
28. Girija AS. Fox3 (+) CD25 (+) CD4 (+) T-regulatory cells may transform the nCoV's final destiny to CNS! COMMENT. WILEY 111 RIVER ST, HOBOKEN 07030-5774, NJ USA; 2021.
29. Nasim I, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, India. Cytotoxicity and anti-microbial analysis of silver and graphene oxide bio nanoparticles [Internet]. Vol. 16, *Bioinformation.* 2020. p. 831–6. Available from: <http://dx.doi.org/10.6026/97320630016831>
30. Rajeshkumar S, Sherif MH, Malarkodi C, Ponnaniakajamideen M, Arasu MV, Al-Dhabi NA, et al. Cytotoxicity behaviour of response surface model optimized gold nanoparticles by utilizing fucoidan extracted from padina tetrastrumatica [Internet]. Vol. 1228, *Journal of Molecular Structure.* 2021. p. 129440. Available from: <http://dx.doi.org/10.1016/j.molstruc.2020.129440>
31. Nandhini JT, Ezhilarasan D, Rajeshkumar S. An ecofriendly synthesized gold nanoparticles induces cytotoxicity via apoptosis in HepG2 cells. *Environ Toxicol [Internet].* 2020 Aug 14; Available from: <http://dx.doi.org/10.1002/tox.23007>

32. Mohapatra S, Leelavathi L, Rajeshkumar S, D. SS, P. J. Assessment of Cytotoxicity, Anti-Inflammatory and Antioxidant Activity of Zinc Oxide Nanoparticles Synthesized Using Clove and Cinnamon Formulation - An In-Vitro Study [Internet]. Vol. 9, Journal of Evolution of Medical and Dental Sciences. 2020. p. 1859–64. Available from: <http://dx.doi.org/10.14260/jemds/2020/405>
33. Kumar SA, Aravind Kumar S, Department of Orthodontics, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Science (SIMATS), Saveetha University, et al. Antimicrobial activity of silymarin mediated zinc oxide and hydroxy apatite nanoparticles against oral pathogens [Internet]. Vol. 16, Bioinformation. 2020. p. 863–8. Available from: <http://dx.doi.org/10.6026/97320630016863>
34. Wu S, Rajeshkumar S, Madasamy M, Mahendran V. Green synthesis of copper nanoparticles using *Cissus vitiginea* and its antioxidant and antibacterial activity against urinary tract infection pathogens [Internet]. Vol. 48, Artificial Cells, Nanomedicine, and Biotechnology. 2020. p. 1153–8. Available from: <http://dx.doi.org/10.1080/21691401.2020.1817053>
35. Synthesis of Silver Nanoparticles Using *Ocimum Sanctum* L (Tulsi) Leaf Extract & Study of Antimicrobial Activity: A Green Chemistry Approach [Internet]. Vol. 8, International Journal of Green and Herbal Chemistry. 2019. Available from: <http://dx.doi.org/10.24214/ijghc/gc/8/3/60109>
36. Barma MD. Synthesis of Triphala Incorporated Zinc Oxide Nanoparticles and Assessment of its Antimicrobial Activity Against Oral Pathogens : An In-Vitro Study [Internet]. Vol. 13, Bioscience Biotechnology Research Communications. 2020. p. 74–8. Available from: <http://dx.doi.org/10.21786/bbrc/13.7/14>
37. Vikneshan M, Saravanakumar R, Mangaiyarkarasi R, Rajeshkumar S, Samuel SR, Suganya M, et al. Algal biomass as a source for novel oral nano-antimicrobial agent. *Saudi J Biol Sci.* 2020 Dec;27(12):3753–8.
38. Chellapa LR, Shanmugam R, Indiran MA, Samuel SR. Biogenic nanoselenium synthesis, its antimicrobial, antioxidant activity and toxicity [Internet]. Vol. 9, Bioinspired, Biomimetic and Nanobiomaterials. 2020. p. 184–9. Available from: <http://dx.doi.org/10.1680/jbibn.19.00054>
39. Boroumand S, Safari M, Shaabani E, Shirzad M, Faridi-Majidi R. Selenium nanoparticles: synthesis, characterization and study of their cytotoxicity, antioxidant and antibacterial activity [Internet]. Vol. 6, Materials Research Express. 2019. p. 0850d8. Available from: <http://dx.doi.org/10.1088/2053-1591/ab2558>
40. Forootanfar H, Adeli-Sardou M, Nikkhoo M, Mehrabani M, Amir-Heidari B, Shahverdi AR, et al. Antioxidant and cytotoxic effect of biologically synthesized selenium nanoparticles in comparison to selenium dioxide. *J Trace Elem Med Biol.* 2014 Jan;28(1):75–9.
41. S. RJ, Roy A, Shanmugam R, E. DW. Preparation and Characterization of Cinnamon Oil



Mediated Gold Nanoparticles and Evaluation of Its Cytotoxicity Using Brine Shrimp Lethality Assay [Internet]. Vol. 9, Journal of Evolution of Medical and Dental Sciences. 2020. p. 2894–7. Available from: <http://dx.doi.org/10.14260/jemds/2020/633>

42. Vyas J, Rana S. Antioxidant activity and green synthesis of selenium nanoparticles using allium sativum extract [Internet]. Vol. 9, International Journal of Phytomedicine. 2017. p. 634. Available from: <http://dx.doi.org/10.5138/09750185.2185>
43. Kumar A, Arvina R, Rajeshkumar S. ANTIFUNGAL ACTIVITY OF CLOVE AND CINNAMON MEDIATED SELENIUM NANOPARTICLES: AN In vitro STUDY. PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY. 2020 Aug 24;18–23.
44. Barma MD, Kannan SD, Indiran MA, Rajeshkumar S, Pradeep Kumar R. Antibacterial Activity of Mouthwash Incorporated with Silica Nanoparticles against *S. aureus*, *S. mutans*, *E. faecalis*: An in-vitro Study [Internet]. Journal of Pharmaceutical Research International. 2020. p. 25–33. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1630646>
45. Francis T, Rajeshkumar S, Roy A, Lakshmi T. Anti-inflammatory and Cytotoxic Effect of Arrow Root Mediated Selenium Nanoparticles [Internet]. Vol. 12, Pharmacognosy Journal. 2020. p. 1363–7. Available from: <http://dx.doi.org/10.5530/pj.2020.12.188>