

## **Review Article**

### **Overview on Causes and Updated Management of Favism**

#### **Abstract:**

Favism is most common in those who have G6PD deficiency from the Mediterranean region. As ~~h~~Hemolytic anaemia is the most common complication of G6PD deficiency, and it can be life-threatening in certain people. Infection, hyperglycemia, certain meals, and certain drugs can all cause hemolysis ~~therefore~~, ~~The~~~~the~~ most prevalent enzymopathy is glucose-6-phosphate dehydrogenase (G6PD) deficiency, which affects an estimated 400 million individuals, globally. ~~and~~ ~~e~~Exposure to some medicines might cause hemolytic anaemia. The most important management technique is to avoid oxidative stresses ~~by avoiding in the first place to avoid~~ a hemolytic crisis. ~~A~~ ~~And~~ ~~also~~, avoidance of exposure to food and med~~i~~ecines that causes hemolytic anaemia episodes

This review looks at etiology, ~~epidemiology~~, ~~pathophysiology~~, ~~evaluation~~ and management of the disease.

## Introduction:

Favism characterises a subgroup of patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency's susceptibility to, and clinical presentation of, acute haemolytic crises as a result of eating broad beans. The most prevalent enzymopathy is glucose-6-phosphate dehydrogenase (G6PD) deficiency, which affects an estimated 400 million individuals globally.[1]

Hemolytic anaemia is the most common complication of G6PD deficiency, and it can be life-threatening in certain people. Infection, hyperglycemia, certain meals, and certain drugs can all cause hemolysis. Though G6PD insufficiency is found in limited areas throughout Africa, the Middle East, and Southeast Asia, it affects around one in every ten African-American men in the United States. As global travel to the United States expands and immigrants gain access to the American health-care system, the disease may become more relevant in the future, resulting in a higher-than-expected prevalence.[1]

The enzyme glucose-6-phosphate dehydrogenase (G6PD) is found in the cytoplasm of all cells in the human body. It's a maintenance enzyme that protects cells from reactive oxygen species damage (ROS). This is accomplished by supplying substrates that prevent oxidative damage.[2]

Because of their involvement in oxygen transport and their incapacity to replace cellular proteins as mature cells, erythrocytes are particularly sensitive to ROS. During times of heightened reactive oxygen species production, inherited deficits in glucose-6-phosphate dehydrogenase can cause severe hemolytic anaemia.[2]

Patients who depend on [red blood cell \(RBC\)](#) transfusions may be at a higher risk of receiving blood from G6PD-deficient donors. [SCD](#) patients, for example, frequently require transfusions of blood group antigen-negative RBCs to avoid or respond to alloantibody formation; thus, based on blood group antigen frequencies, they are more likely to receive RBCs from African-American donors, who ~~may are also more likely to be~~ G6PD deficient.[5,4]

If these individuals ~~have got~~ an infection or were given a drug that caused oxidative stress ~~simultaneously at the same time~~, they would be at [a](#) greater risk (e.g. a pregnant SCD patient treated with nitrofurantoin for a urinary tract infection) .[5]

The most prevalent life-threatening manifestation of G6PD deficiency is hemolytic anaemia. Patients, typically report with symptoms such as jaundice, lethargy, back pain, tachypnea, and tachycardia. Reduced haemoglobin and red

Write out the word in **:[1]MS]Comment**  
fullb before abbreviating

blood cell counts, reticulocytosis, elevated lactate dehydrogenase, and increased unconjugated bilirubin are all relevant diagnostic results.[1]

### Etiology:

The G6PD enzyme is represented by the Gd gene. Because this gene is found on the long arm of the X chromosome, ~~and it~~ is inherited in an X-linked manner. Mutations that affect the protein structure and so limit its activity or the amount of enzyme generated, can cause G6PD deficiency. There are now 186 known human G6PD mutations, the majority of which are single-nucleotide variants. ~~Because~~ Complete inactivation of G6PD would be harmful to a growing embryo. ~~N,~~ none of the mutation patterns found in humans produce it. [2,3]

The G6PD gene is found on the long arm of the X chromosome and is made up of 13 exons that cover about 18 kb. The gene's coding sequence is 1,545 base pairs (bp) long, ~~and it~~ codes for 515 amino acids from the G6PD main sequence. G6PD insufficiency is the most prevalent human enzyme abnormality, affecting over 400 million people worldwide. A deficient hemizygote is a guy who inherits the damaged X chromosome from his mother and carries the deficient gene. Each parent's X chromosome is faulty in female homozygotes. Only one faulty X chromosome is passed down to female heterozygotes.[6]

Exposure to some medicines might cause hemolytic anaemia episodes: acetanilid, cotrimoxazole, dapsone, doxorubicin, furazolidone, methylene blue, moxifloxacin, nalidixic acid, naphthalene, niridazole, nitrofuratoin, norfloxacin, pamaquine, pentaquine, phenazopyridine, phenylhydrazine, primaquine, ras The degree to which a person is susceptible to a substance varies from person to person.[7]

### Epidemiology:

G6PD is the most prevalent human enzyme deficiency, affecting an estimated 400 million people globally. Due to X-linked inheritance, men are more typically impacted when compared to than women. Tropical and subtropical regions are the most affected. Surprisingly, there is evidence that G6PD deficiency protects against uncomplicated malaria, however, ~~but~~ not against severe malaria. G6PD deficiency and malaria have a protective mechanism that is still being researched. G6PD insufficiency is more common in people of African, Mediterranean, or Asian cultures, possibly because to its malaria-protective effect.[2]

G6PD is the rate-limiting first step of the pentose phosphate pathway, which converts nicotinamide adenine dinucleotide phosphate (NADP) to its reduced form, NADPH, using glucose-6-phosphate. NADPH is essential in red blood

Repeated .Remove :[2]MS]Comment information

Rewrite sentence for :[3]MS]Comment clarity

Repeated .Remove :[4]MS]Comment information

Individuals from the :[5]MS]Comment ?... torpicaland regions are

cells for preventing cellular components from being damaged by oxygen-free radicals. It accomplishes this by acting as a substrate for the glutathione reductase enzyme. Reduced glutathione can be utilised to convert hydrogen peroxide to water and protect cellular components, especially the cell walls of mature red blood cells (RBCs), which have limited repair capacity. [2]

Remove :[6]MS]Comment

### Pathophysiology:

The first step in the pentose phosphate pathway is catalysed by glucose-6-phosphate dehydrogenase (G6PD). Converting glucose to ribose-5-phosphate, a precursor of RNA, DNA, ATP, CoA, NAD, and FAD, is part of the pentose phosphate pathway (PPP). The route also involves the production of NADPH, which provides the cell with reducing energy by ensuring that reduced glutathione remains within the cell. Reduced glutathione protects cells from oxidative damage by acting as an antioxidant.[2]

Most cells have a backup set of metabolic pathways that can generate the necessary intracellular NADPH. Red blood cells, on the other hand, lack the "other NADPH makers." As a result, G6PD deficiency is particularly dangerous in red blood cells, where any oxidative stress leads to hemolytic anaemia. Consumption of fava beans, certain medicines, infections, and certain metabolic disorders such as diabetic ketoacidosis can all cause oxidative stress.[7]

Puddling of haemoglobin occurs as a result of oxidative denaturation, and bite or hemiblisters cells form in the peripheral smear. Denatured haemoglobin precipitates, and can be seen as often known as "Heinz bodies," can be seen. A normal peripheral blood smear, which shows red blood cells of uniform size and shape, can be distinguished from this.[7]

"State exactly what is :[7]MS]Comment  
" this

### Clinical implications:

When managing with a G6PD-deficient patient, every health care professional should ~~must~~ exercise caution. Hemolysis can be caused by fava beans, certain medicines, infections, and metabolic disorders. Inadequate care for G6PD-deficient people, who develop acute hemolytic anaemia, can result in severe brain impairment or death. As a result, the practitioner should advise the high-risk patient and his or her caregiver to check for signs and symptoms of a hemolytic crisis which includes (cyanosis, headache, dyspnea, exhaustion, substernal pain, jaundice, scleral icterus, and dark urine). Prior to the follow-up appointment, a simple postoperative phone call to check on the patient, could be critical to his or her health. Hemoglobin concentrations are thought to begin to recover after removal of the offending hemolytic agent, after 8 to 10 days;

Remove :[8]MS]Comment

Remove :[9]MS]Comment

consequently, acute hemolysis seldom leads to severe anaemia which needs a blood transfusion (especially in infants) [7]

Post - operatively, the G6PD-deficient patient may exhibit clinical symptoms that necessitate further support or therapy. Hemolysis usually appears 1 to 3 days after contact with the triggering factors. Acute hemolysis is self-limiting, however, but it can be severe enough to necessitate a blood transfusion in rare cases. Cyanosis, fatigue, tachycardia, dyspnea, lethargy, lumbar/substernal pain, abdominal pain, splenomegaly, hemoglobinuria, and/or scleral icterus are all possible symptoms. Furthermore, haemoglobin breakdown products accumulate in the blood, causing jaundice. The breakdown products, and they can be excreted in the urine, resulting in a dark brown discoloration. [7]

Avoidance of oxidative stresses is the major treatment for G6PD deficiency. Anemia can be severe enough to necessitate a blood transfusion in rare cases. [9-11]

Again repeat :[10]MS]Comment  
?information

Hemolysis manifests itself clinically after 24 to 72 hours of medication administration, and anaemia develops until about day 7. This makes it difficult for doctors to detect a hemolytic crisis in patients undergoing outpatient or short-stay hospital operations (less than 24). Therefore, so the most important management technique is to avoid oxidative stresses thereby in the first place to avoid a hemolytic crisis. Acute hemolysis in G6PD-deficient adults, on the other hand, is usually short-lived and does not require therapy. The offending agent should be eliminated in the event of a hemolytic crisis, and the patient should be continuously watched. A daily complete blood count should be performed at the very least to monitor the requirement for a blood transfusion.[7]

### **Evaluation :**

- Neonatal evaluation:-

In newborn infants, check for jaundice by looking for a yellowish color on the skin in a well-lit room. In infants, total serum bilirubin (TSB) or transcutaneous bilirubin (TcB) readings are more objective. A risk-stratified bilirubin nomogram can be used to help select the best treatment for newborns with elevated bilirubin levels.[2]

Although G6PD deficiency screening tests are available, they are not commonly used in the United States. However, screening should be explored in infants with severe jaundice that is resistant to phototherapy. Screening tests should also, be available to patients or who with have a family history or ethnicity that suggests G6PD deficiency. To determine the production of NADPH from

NADP, the most frequent screening approach is a fast fluorescent spot test. Quantitative spectrophotometric analysis can also be used for screening.[2]

- Children and adult evaluation:-

The evaluation of older individuals with G6PD deficient problems begins with a thorough medical history, including new drugs, and a search for a family history of related symptoms. In patients with G6PD deficiency, the stress of infection might induce a hemolytic event, so it's also important to check for infection.

CBC, bilirubin levels, reticulocyte count, serum aminotransferases, and lactate dehydrogenase levels are among the tests performed in the laboratory. Schistocytes may be seen on a peripheral blood smear, indicating hemolysis. [2]

### **Management :**

- Neonates

The infant patient's treatment focuses on preventing kernicterus and minimizing jaundice. This includes phototherapy that follows stated standards. An exchange transfusion may be required in severe cases.[2,15,16]

- Adults and Children

**The treatment of older individuals** is mostly determined by the overall clinical picture. Supportive therapy, as well as the withdrawal and avoidance of the offending medications, can be used to manage less severe symptoms. Infections should be treated as suggested by the history and examination. Transfusions may be required in more severe situations. [2]

Avoidance of oxidative stresses is the major treatment for G6PD deficiency. Anemia can be severe enough to necessitate a blood transfusion in rare cases. Splenectomy is generally not advised. Although G6PD deficiency is frequently asymptomatic and the resulting hemolysis is usually short-lived, folic acid and iron may be beneficial in hemolysis. Antioxidants like vitamin E and selenium haven't been shown to help with G6PD deficiency.[9-11]

Favism is most common in those who have G6PD deficiency from the Mediterranean region[8].

To manage hemolysis in individuals with G6PD impairment, it is important to identify and eliminate the precipitating factor. Anemia should be treated appropriately, with the understanding that hemolysis is self-limiting and usually resolves in 8 to 14 days. Transfusions are only used in exceptional circumstances. In most cases, splenectomy is ineffective.[12-14]

The majority of people who have glucose-6-phosphate dehydrogenase (G6PD) deficiency don't require treatment. They should, however, be educated to stay away from medications and chemicals that can induce oxidative stress. Broad beans should also be avoided by patients (ie, fava beans).[12-14]

Foods, medicines, and chemicals that can cause hemolysis should be avoided by people with G6PD deficiency. The degree of enzyme insufficiency and consequently the risk posed by such chemicals is determined, in part, by the person's G6PD variation.[12-14]

Fava beans are the most well-known food for causing hemolysis. ~~in fact, Favism~~, or symptomatic attacks of hemolytic anaemia caused by eating fava beans, has been known since antiquity. Other foods to avoid for people with G6PD deficiency include the following:

- a glass of red wine
- Legumes in general
- Blueberries
- Products made from soya
- Water with a tonic effect [13,14]

#### Complications:

- Acute severe haemolytic anaemia causes death (relatively rare).
- Intraocular intravascular haemolysis causes ophthalmological damage. [17]

**Patients with an acute onset of favism** had severe ophthalmological problems, ~~however, but~~ not all of ~~patients~~ suffered from generalised haemorrhagic symptoms at the same time.[18]

- Kidney injury that ~~was~~ severe.
- Infection susceptibility.[17]

#### Conclusion:

Fava beans are the most well-known food for causing hemolysis. ~~in fact, Favism~~, or symptomatic attacks of hemolytic anaemia ~~are~~ caused by eating fava beans. The most important management technique is to avoid oxidative stresses ~~and to in the first place to~~ avoid a hemolytic crisis. Acute hemolysis in

G6PD-deficient adults, on the other hand, is usually short-lived and does not require therapy.

When managing with a G6PD-deficient patient, every health care professional must exercise caution.

Hemolysis can be caused by fava beans, certain medicines, infections, and metabolic disorders. Inadequate care for G6PD-deficient people who develop acute hemolytic anaemia can result in severe brain impairment or death.

## References

1. Bulp J, Jen M, Matuszewski K. Caring for Glucose-6-Phosphate Dehydrogenase (G6PD)-Deficient Patients: Implications for Pharmacy. *P T*. 2015 Sep;40(9):572-4. PMID: 26417175; PMCID: PMC4571844.
2. Richardson SR, O'Malley GF. Glucose 6 Phosphate Dehydrogenase Deficiency. [Updated 2021 Jul 26]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan .-
3. Lauden SM, Chongwain S, Achidi A, Helm E, Cusick SE, Krug A, Slusher TM, Lund TC. Prevalence of glucose-6-phosphate dehydrogenase deficiency in Cameroonian blood donors. *BMC Res Notes*. 2019 Apr 02;12(1):195.
4. Francis R, Jhang J, Hendrickson J, Zimring J, Hod E, Spitalnik S. Frequency of glucose-6-phosphate dehydrogenase-deficient red blood cell units in a metropolitan transfusion service. *Transfusion*. 2012.
5. Francis RO, Jhang JS, Pham HP, Hod EA, Zimring JC, Spitalnik SL. Glucose-6-phosphate dehydrogenase deficiency in transfusion medicine: the unknown risks. *Vox Sang*. 2013 Nov;105(4):271-82. doi: 10.1111/vox.12068. Epub 2013 Jul 2. PMID: 23815264; PMCID: PMC3797222.
6. Arese P, Gallo V, Pantaleo A, Turrini F. Life and Death of Glucose-6-Phosphate Dehydrogenase (G6PD) Deficient Erythrocytes - Role of Redox Stress and Band 3 Modifications. *Transfus Med Hemother*. 2012 Oct;39(5):328-34. doi: 10.1159/000343123. Epub 2012 Sep 17. PMID: 23801924; PMCID: PMC3678266.
7. Elyassi AR, Rowshan HH. Perioperative management of the glucose-6-phosphate dehydrogenase deficient patient: a review of literature. *Anesth Prog*. 2009 Autumn;56(3):86-91. doi: 10.2344/0003-3006-56.3.86. PMID: 19769422; PMCID: PMC2749581.
8. Oliveira S, Pinheiro S, Gomes P, Horta AB, Castro AS. Favismo [Favism]. *Acta Med Port*. 2000 Jul-Aug;13(4):237-40. Portuguese.



9. JENNIFER E. FRANK, Diagnosis and Management of G6PD Deficiency, 2005 Oct 1; 72(7):1277-1282.  
<https://www.aafp.org/afp/2005/1001/p1277.html#afp20051001p1277-b6>
10. Beutler E. G6PD deficiency. Blood. 1994 Dec 1. 84(11):3613-36.
11. Glucose-6-phosphate dehydrogenase deficiency. Accessed online July 20, 2005,  
[at:http://www.thedoctorslounge.net/clinlounge/diseases/hematology/g6pd.htm](http://www.thedoctorslounge.net/clinlounge/diseases/hematology/g6pd.htm).
12. Srikanth Nagalla, Francisco Talavera, et al. Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency Medication, Updated: Jul 19, 2021 Medscape, <https://emedicine.medscape.com/article/200390-treatment#showall>.
13. Samanta S, Kumar P, Kishore SS, Garewal G, Narang A. Donor blood glucose 6-phosphate dehydrogenase deficiency reduces the efficacy of exchange transfusion in neonatal hyperbilirubinemia. Pediatrics. 2009 Jan. 123(1):e96-e100.
14. Drugs that should be avoided - Official List. G6PD Deficiency Association. Available at [https://www.g6pd.org/en/G6PDDeficiency/SafeUnsafe/DaEvitare\\_ISS-it](https://www.g6pd.org/en/G6PDDeficiency/SafeUnsafe/DaEvitare_ISS-it). Accessed: July 19, 2021.
15. Mohamed GS, Lemine SM, Cheibetta S, Mohamed A. [Neonatal screening for glucose-6-phosphate dehydrogenase (G6PD) deficiency in Mauritania]. Pan Afr Med J. 2018;30:224.
16. Tripathi P, Kumar R, Agarwal S. Spectrum and hematological profile of hereditary anemia in North Indians: SGPGI experience. Intractable Rare Dis Res. 2018 Nov;7(4):258-263.
17. Dr Louise Newson, et al [Favism], Last edited 21 Sep 2015. <https://patient.info/doctor/favism>.
18. CHOREMIS C, JOANNIDES T, KYRIAKIDES B. Severe ophthalmologic complications following favism. Br J Ophthalmol. 1960 Jun;44(6):353-6. doi: 10.1136/bjo.44.6.353. PMID: 13810004; PMCID: PMC509945.

Check : [11]MS]Comment

check : [12]MS]Comment

Use Surname and : [13]MS]Comment  
 ?Initial of the author thereafter the

not consistent with : [14]MS]Comment  
 ?the other references

Not consistent : [15]MS]Comment