

**Nutritional composition, functional properties and food applications of Millet grains**

**Abstract**

Millet is an important nutritious ancient minor cereal food crop. This work reviews the composition, functional properties and food application of millet grains. The review shows that this cereal grains is a good source of carbohydrate and starch, with minute proteins, fat, vitamins and other nutrients. The functional properties of the cereal grains (Millet) was also evaluated and the findings gotten from various authors shows that the grains has a good functional properties in terms of their bulk density, oil absorption capacity, water absorption capacity, least gelatinization temperature and host of others. The pasting properties were also researched on and various authors attested to the potentiality of the grains in terms of the pasting properties. The food applications of the grains was not left out since the basic essence of this review is to see to the betterment of the livelihood of human, as such the various foods that can be produced from these grains were also looked into foods like millet ball “Fura”, tuwo, gruel, alcoholic beverages (like pito, burukutu) and non-alcoholic beverages (like “kunu zaki”) where all examined and conclusively the grains were rich sources of meals.

**Keywords:** Millet, Flour, Kunu, cereals, grains.

**1.0 Introduction**

22 The word 'millet' gets its origin from the French word "*mille*" which means thousand, with a  
23 handful of millet containing up to 1000 grains [1]. Millet belongs to the group of small-seeded  
24 species of cereal crops or grains which are annual plants [2]. The grain belongs to the family  
25 *Poaceae* which originated in Ethiopia and the sub-family *Chloridodeae* [3]. Different cultivars of  
26 millet grains exist: brown, light brown and white [4, 5], with grain colour used as the distinct  
27 means of cultivar differentiation. The white cultivars have been developed mainly for the baking  
28 industry, the brown and light brown types used for porridge while the brown cultivar is utilized  
29 for brewing traditional opaque beer in Southern Africa [6].

30 The grain millet is a semi-arid region crop cultivated in dry areas with limited rainfall and can  
31 adapt to various agro-climatic conditions [7]. Period of cultivation of the grain ranges between  
32 February and August with harvest period set in June or January. Millet grains are cultivated in  
33 Nepal [8], Sri Lanka, Bhutan and the Himalayan regions of India. The grain is also cultivated in  
34 Taiwan, China, Japan (to a limited extent), as well as in South Carolina in the United States.  
35 About 55-60% of globally produced millet is cultivated in Africa [9] mainly in Ethiopia, Kenya,  
36 Nigeria, Malawi, Tanzania, Uganda, Zambia and Zimbabwe. The grain is widely cultivated in  
37 Africa using different names. The total annual production of all millets worldwide is  
38 approximately 4.5-5 million tons [10], with India alone producing about 2.5 million tons and  
39 some countries in Africa accounting for about 2 million tons of the grains. India is thus reported  
40 to be the largest producer of millet [2], contributing a total of 60% of the global production [6].

41 Millet grains are gluten-free, non-acid-forming [9], easy to digest with low glycemic index [11].  
42 Its low glycemic index food property is reported to be a good choice for people with celiac  
43 disease (disease caused by gluten-containing cereal protein ingestion) and diabetes as  
44 consumption of the grain assist in the regulation of blood glucose level [12]. The grains consist

45 of dietary fiber, carbohydrates, iron and calcium in high concentration when compared to other  
46 cereal grains. Millet grains also contain high amount of magnesium and phosphorus [13].  
47 Krishnan *et al.* [14] reported that millet grains contain polyphenols and phytates which are  
48 known to influence the bioavailability of minerals. In addition to their nutritive value, several  
49 potential health benefits such as preventing cancer and cardiovascular diseases, reducing tumor  
50 incidence, lowering blood pressure, risk of heart disease, cholesterol and rate of fat absorption,  
51 delaying gastric emptying, and supplying gastrointestinal bulk have been reported for millet [13,  
52 15, 16].

53 Millets are small-seeded with different varieties such as pearl millet (*Pennisetum glaucum*),  
54 finger millet (*Eleusine coracana*), kodo millet (*Paspalum setaceum*), proso millet (*Penicum*  
55 *miliaceum*), foxtail millet (*Setaria italic*), little millet (*Panicum sumatrense*), and barnyard millet  
56 (*Echinochloa utilis*). They are known as coarse cereals beside maize (*Zea mays*), sorghum  
57 (*Sorghum bicolor*), oats (*Avena sativa*), and barley (*Hordeum vulgare*) [17]. Millet is known as  
58 *ragi* and *mandia* in the Bastar region of Chhattisgarh and offers both nutritional and livelihood  
59 security for human beings and also feed security for diverse livestock populations in dry land  
60 regions of rural India [18].

61 Millets are not placed as a single important commodity in the North American and European  
62 food basket at the present time, but their importance as an ingredient in multigrain and gluten-  
63 free cereal products has been highlighted. However, in many African and Asian areas, millets  
64 serve as a major food component and various traditional foods and beverages, such as bread  
65 (fermented or unfermented), porridges, and snack foods are made of millet, specifically among  
66 the non-affluent segments in their respective societies [19]. Millet grains, before consumption  
67 and for preparing of food, are usually processed by commonly used traditional processing

68 techniques include decorticating, malting, fermentation, roasting, flaking, and grinding to  
69 improve their edible, nutritional, and sensory properties. Processing of millet grains comprises  
70 both the traditional and modern methods. The modern methods of processing can be employed in  
71 the manufacture of value-added products such as soaked, cooked, malted, fermented, popped or  
72 puffed, extruded and multi-grain flour. Traditional methods of millet processing also include a  
73 method of spreading and drying the grains in the sun for a period of one week. Upon drying, the  
74 grains are stored in a bag and later used for the processing and manufacture of different food  
75 products [20]. Dried millet grains can be stored for more than 5-10 years, but a major hurdle is  
76 that the grains are very tiny and not easy to handle. The grains are resistant to diseases and  
77 insects but are easily invaded by fungal disease [21]. Despite its usefulness and health beneficial  
78 properties, there is little research and innovation on millet grains/flours as compared to  
79 conventional cereal grains such as maize, sorghum, rice and wheat.

## 80 **2.0 Nutritional Composition of Millet**

81 Nutritional quality of food is a key element in maintaining human overall physical well-being  
82 because nutritional well-being is a sustainable force for health and development and  
83 maximization of human genetic potential. Therefore, for solving the problem of deep-rooted food  
84 insecurity and malnutrition, dietary quality should be taken into consideration [22].

85 In addition to the cultivating advantages of millet, they were found to have high nutritive value  
86 and comparable to that of major cereals such as wheat and rice [23]. Millet proteins are good  
87 sources of essential amino acids except lysine and threonine but are relatively high in  
88 methionine. Millets are also rich sources of phytochemicals and micronutrients [24, 25]. For  
89 example, pearl millet was found significantly rich in resistant starch, soluble and insoluble  
90 dietary fibers, minerals, and antioxidants [26]. It contains about 92.5% dry matter, 2.1% ash,

91 2.8% crude fiber, 7.8% crude fat, 13.6% crude protein, and 63.2% starch [27]. Also, foxtail  
92 millet protein characterization showed that its protein concentrate is a potential functional food  
93 ingredient and the essential amino acid pattern suggests possible use as a supplementary protein  
94 source to most cereals because it is rich in lysine [28]. Finger millet also is known to have  
95 several potential health benefits and some of the health benefits are attributed to its polyphenol  
96 contents [29]. It has a carbohydrate content of 81.5%, protein 9.8%, crude fiber 4.3%, and  
97 mineral 2.7% that is comparable to other cereals and millets [30]. Its crude fiber and mineral  
98 contents are markedly higher than those of wheat (1.2% fiber, 1.5% minerals) and rice (0.2%  
99 fiber, 0.6% minerals); its protein is relatively better balanced; it contains more lysine, threonine,  
100 and valine than other millets [31]. In addition, black finger millet contains 8.71 mg/g dry weight  
101 fatty acid and 8.47 g/g dry weight protein [32]. Kodo millet and little millet were also reported to  
102 have 37% to 38% of dietary fiber, which is the highest among the cereals; and the fat has higher  
103 polyunsaturated fatty acids [33]. The protein content of proso millet (11.6% of dry matter) was  
104 found to be comparable with that of wheat with significant higher content of leucine, isoleucine,  
105 and methionine [34]. Thus, the presence of all the required nutrients in millets makes them  
106 suitable for large-scale utilization in the manufacture of food products such as baby foods, snack  
107 foods, and dietary food and, increasingly, more millet products have entered into the daily lives  
108 of people, including millet porridge, millet wine, and millet nutrition powder from both grain and  
109 flour form [35, 36].

### 110 **3.0 Functional Properties of Millet**

111 Cereal grains contain 60 to 70% starch and are excellent energy rich food for human. Doctors  
112 recommended cereals as the first food to be added to infant diets and a healthy diet for adults  
113 should have most of its calories in the form of complex carbohydrates such as cereals grain

114 starch. Cereals and millets form the staple food of diets in about 75% of the countries of the  
115 world [37].

116 Cereals are an excellent source of vitamin and minerals including fat soluble vitamin E, which is  
117 an essential antioxidant. The cereal grains are an easy protein source as required by  
118 Recommended Daily Allowance (RDA) but unfortunately they lack the essential amino acid  
119 lysine and therefore they must not be used as the sole source of dietary protein [38]. Cereal  
120 grains contain about 58 to 72% carbohydrates, 8 to 13% protein, 2 to 5% fat, and 2 to 11%  
121 indigestible fibre. They also contain 300 to 350 kcal/100 g of the grain. Carbohydrates are  
122 present in the form of digestible starches and sugars. The operations of milling generally remove  
123 much of the indigestible fibre and fat from the grains when they are to be consumed for human  
124 food.

125 Functional properties are the fundamental physico-chemical properties that reflect the complex  
126 interaction between the composition, structure, molecular conformation and physico-chemical  
127 properties of food components together with the nature of environment in which these are  
128 associated and measured [39, 40]. Functional characteristics are required to evaluate and possibly  
129 help to predict how new proteins, fat, fibre and carbohydrates may behave in specific systems as  
130 well as demonstrate whether or not such protein can be used to stimulate or replace conventional  
131 protein [39, 40].

132 The food property is characterized of the structure, quality, nutritional value and /or acceptability  
133 of a food product. A functional property of food is determined by its physical, chemical, and/or  
134 organoleptic properties. Example of such functional properties may include solubility,  
135 absorption, and water retention, frothing ability, elasticity and absorptive capacity for fat and

136 foreign particulars, pasting properties, emulsification, hydration (water binding), viscosity,  
137 foaming, solubility, gelation, cohesion and adhesion.

### 138 **3.1 Water Absorption Capacity (WAC)**

139 Water absorption capacity is the amount of water taken up by flour to achieve the desired  
140 consistency and create a quality end-product. Flour with high water absorption may have more  
141 hydrophilic constituents such as polysaccharides. Protein has both hydrophilic and hydrophobic  
142 nature and therefore they can interact with water in foods. Increase in the WAC has always been  
143 associated with increase in the amylose leaching and solubility, and loss of starch crystalline  
144 structure. Thilagavathi *et al.* [41], compared WAC of various types of millet with wheat and  
145 soybeans flour and found out that it ranged from 74.08 to 76.83 ml/100g, 74.08 to 78.83 ml/100g  
146 and (58.17- 60.02 ml/100g) for millet, wheat and soybean flour respectively. The observed  
147 variation in different flours may be due to different protein concentration, their degree of  
148 interaction with water and conformational characteristics [42].

### 149 **3.2 Oil Absorption Capacity (OAC)**

150 OAC has been attributed to the physical entrapment of oil; this is important since oil acts as  
151 flavor retainer and increases the consumers' taste of food [43]. OAC of millet flour has been a  
152 wide research that has been conducted by various researchers at varying conditions and different  
153 results obtained. The oil absorption capacity according to Amir *et al.* [44], on finger millet flour  
154 was found to be 1.93g/g and that of pearl millet flour is 1.60g/g. There is an advantage for best  
155 organoleptic characteristics of meal that high water and oil absorption capacity of the flour can  
156 positively influence the flavor, moisture and fat content in food [45].

157

158

### 159 3.3 Bulk Density

160 Bulk density is a measure of heaviness of flour and is generally affected by the particle size and  
161 the density of the flour. It is very important in determining the packaging requirement, material  
162 handling and application in wet processing in the food industry. According to Amir *et al.* [44],  
163 the bulk density of pearl millet and finger millet flours ranged from 0.67g/ml to 0.54g/ml. The  
164 differences in the values of bulk density between these flours are likely due to varietal  
165 differences. Krishnan *et al.* [46] found a bulk density of 0.5g/ml, 0.50g/ml and 0.6g/ml in native,  
166 malted and hydrothermally treated finger millet seed coat.

### 167 3.4 Foaming Capacity and Stability

168 The Forming capacity (FC) and foaming stability (FS) are determined by a loss of liquid  
169 resulting from destabilization that is measured as a volume decrease. Foaming formation is  
170 governed by three factors: transportation, penetration, and reorganization of the molecule at the  
171 air–water interface. Therefore, for good foaming, the protein should be capable of migrating at  
172 the air–water interface, unfolding and rearranging at the interface [47].

173 Yagoub and Abdalla, [48] presented results of FC which varied from 116.55% to 151.58%. They  
174 were in agreement with those of cowpea and millet flour as reported by Akubor [49] and  
175 Jayathilake *et al.* [50] respectively. An increase in FC might be initiated by a decrease in surface  
176 tension of the air and water interface, which consequently caused absorption of soluble protein  
177 molecules for hydrophobic interactions. The FC of a food materials depended on the surface  
178 active properties of its protein [51].

### 179 3.5 Gelatinization Temperature

180 Gelatinization temperature of all the flour samples investigated by Iwe *et al.* [52], ranged from  
181 29.00 to 74.00°C and it fell within the range (<75°C) reported by ARSO [53]. There is a



182 significant variation between the flour varieties in their gelatinization temperatures.  
183 Gelatinization temperature is the temperature at which starch molecules in a food substance lose  
184 their structure and leach out from the granules as swollen amylose and it affects the time required  
185 for the cooking of food substances [54]. According to Chandra and Samsher, [55], a flour which  
186 has a higher starch content takes lower temperature for gelatinization and those with lower starch  
187 content takes higher temperature to gelatinize.

### 188 **3.7 Pasting Properties**

189 Pasting properties are important in predicting the behavior of flours during and after cooking.  
190 The difference in Peak viscosity observed in the samples is an indication of various degrees of  
191 starch gelatinization and difference in amylose content of the blends. Sanni *et al.* [56] noted that  
192 high peak viscosity is closely associated with the high starch damage which in turn enhances  
193 viscosity.

194 Bhupender *et al.* [57] did a research on the pasting properties of starches from different pearl  
195 millet cultivars measured using RVA. Starches from different cultivars displayed a significant  
196 variation in all their pasting parameters. The starch suspensions showed gradual increase in  
197 viscosity with increase in temperature. The increase in viscosity with temperature may be  
198 attributed to the removal of water from the exuded amylose by the granules as they swell. Peak  
199 Viscosity is an indicator of water binding capacity and ease with which the starch granules are  
200 disintegrated and often correlated with final product quality [58]. Peak viscosity (PV) of different  
201 starch samples was observed to be in the range from 1665 to 1998 cP., Breakdown viscosity  
202 (BV) of starch from different pearl millet cultivars differed ranged from 414 to 769 cP. The  
203 breakdown is caused by disintegration of gelatinized starch granules structure during continued  
204 stirring and heating, thus, indicating the shear thinning property of starch [59]. A low breakdown

205 value suggests the stability of starches under hot conditions. Amylose content is believed to have  
206 a marked influence on the breakdown viscosity (measure of susceptibility of cooked starch  
207 granule to disintegration) and the setback viscosity (measure of recrystallization of gelatinized  
208 starch during cooling) [60]. Lower level of amylose to reinforce the molecular network within  
209 the granules resulted in greater breakdown viscosity. High amylose content has also been  
210 suggested as the major factor contributing to the non-existence of a peak, a high stability during  
211 heating, and a high setback during cooling [61, 62]. Setback viscosity of pearl millet starches  
212 ranged from 627 to 1064 cP. Final viscosity indicates the ability of the starch to form a viscous  
213 paste. Final viscosity of pearl millet starches ranged from 1931 to 2476 cP. A higher final  
214 viscosity relates to the high resistance to shear. Increase in final viscosity might be due to the  
215 aggregation of the amylose molecules. Pasting temperature of starch suspensions ranged from  
216 88.1 to 90.2°C. Stability ratio explains the resistance of a starch paste to viscosity breakdown as  
217 shear is applied.

#### 218 **4.0 Food Application of Millet**

##### 219 **4.1 Tuwo Production**

220 Tuwo is a local delicacy of the Northern part of Nigeria; it is made from millet, maize or  
221 sorghum as the case may be and the choice of the producer. Tuwo is a solid food which is made  
222 in forms of balls or swallows; it is made from the flour produced from any of the above  
223 mentioned grains [63]. According to Odusola *et al.* [64], tuwo is produced by getting millet, the  
224 grain is then sorted, dehulled, winnowed and mill into smooth flour and then sieve appropriately,  
225 water will be heated in the pot to boil, little portion of the flour is used to make a slurry in a cold  
226 water and its then transferred in to the boiling water and is allowed to boil together properly.  
227 After which the sieved flour of the millet is pour gradually into the pot containing the boiled

228 water and the boiled slurry and stir until a desired thickness is obtained, the food is allowed to  
229 heat for additional ten to fifteen minutes and it's then stirred and it's ready to be served with any  
230 desired soup.

#### 231 **4.2 Millet Ball Production “Fura”**

232 Fura is a staple food for the Fulanis and Hausas. The single most important cereal grain for fura  
233 production is millet or its twin grain sorghum [65]. In tropical Africa, millet grains are milled  
234 and used to produce thick porridges which are known by various names in different parts of the  
235 continent. In west Africa particularly in Nigeria, Ghana and Burkina Faso, one such thick  
236 porridge is called ‘fura’ - a semi- solid dumpling cereal meal [66]. Fura is produced mainly from  
237 moist millet flour, blended with spices, compressed into balls and boiled for 30 minutes. While  
238 still hot, the cooked dough is worked in the mortar with the pestle (with addition of hot water)  
239 until a smooth, slightly elastic, cohesive lump (fura) is formed. The fura dough is rolled into 25–  
240 30 g balls by hand and dusted with flour. The fura is made into porridge by crumbling the fura  
241 balls into fermented whole milk (kindrimo) or 24 fermented skim milk (nono) [66]. Sugar may  
242 be added to taste, the mixture is called ‘fura da nono’ in Nigeria. It is a popular mid-day meal.  
243 Fura is produced at home both for family and commercial consumption. The producers of fura  
244 still use the modern method to dehull the grain and to reduce the dehulled grain into flour on like  
245 the formal traditional methods of mortar and pestle thing. Fura is typically distributed with  
246 minimum packaging. Processors and retailers of fura are primarily concerned with reducing  
247 waste and having a container for their food. Fura has a limited shelf-life of one day at ambient  
248 temperature [66]. Usually, a day after production, fura shows visible mold growth on the surface.  
249 The short shelf-life has always been a major deterrent to large-scale production. Thus, improving

250 the processing, packaging and storage life of fura are of interest before food manufacturers can  
251 think of large scale production.

### 252 **4.3 Gruel Production**

253 Millet has been used for gruel production or as breakfast meals which are in turn produced into  
254 pap, “ogi”, “akamu” etc and are taken with any other desired snacks for adequate nourishment  
255 and some the gruels are been enriched with other food products like soybeans, ginger and host of  
256 other [67].

257 In the production of the gruels, the raw millet is graded, washed and soaked for 72 hours and the  
258 water decanted, some producers change the water daily that is after 24 hours while others leave it  
259 for that period of 72 hours fermentation. The fermented grains are then washed and wet milled in  
260 a clean grinding machine. The grain slurry is then filtered with muslin cloths and the filtrate is  
261 allowed to sediment and the water decanted. Then the slurry or gruel can be cooked and made  
262 into pap, ogi, akamu, kwokwo and host of others dependant on the choice of the producers [64].

### 263 **4.4 Local Alcoholic Beverage Production**

264 In many times past barley has been the sole cereal grains that are used in the production of  
265 alcoholic beverages in Western part of Africa and Nigeria inclusive, this practice has left us in  
266 the dependant stage of life, adding no dividend to the economy of the country; rather it takes  
267 from it to expand and enrich others and growing us in the rank of a dependent Nation. Recent  
268 research works has strive to break that barrier of over dependency by introducing other means of  
269 using home grown grains in the production or manufacture of some of this alcoholic beers and  
270 beverages, of which millet is part of this innovations. Alcoholic beverages are divided into three  
271 (3) general classes: beers, wines and spirits. Alcoholic beverages that have lower alcohol content  
272 (beer and wine) are produced by fermentation of sugar- or starch containing plant materials.

273 Beverages of higher alcohol content (spirits) are produced by fermentation followed by  
274 distillation. The major local alcoholic beverages produced in Nigeria are Burukutu, palmwine,  
275 pito, and Ogogoro. Burukutu beer is a traditional cereal-based fermented beverage. Cereals are  
276 important in many parts of the world as food sources, and starches from them differ in  
277 physicochemical properties and molecular structures [68]. Millet is an important cereal crops  
278 grown in Nigeria with starch as its main chemical component. The basic characteristics of  
279 Burukutu include a sour taste due to the presence of lactic acid, a pH of 3.3 to 3.5 and an opaque  
280 colour because of suspended solids and yeast. It contains vitamins, iron, manganese, magnesium,  
281 potassium and calcium and also contains about 26.7g of starch and 5.9g of protein per liter [69].  
282 The local beverage is known as Techoukoutou in Benin or Togo, Dolo in Burkina-Faso, Pito in  
283 Ghana, Burukutu or Otika in Nigeria, Bilibili in Tchad, Mtama in Tanzania, Kigage in Rwanda  
284 [70, 71]. The manufacturing processes are very variable and dependent on the geographical  
285 location. Generally the production process of cereals involves, malting, steeping, germination,  
286 milling, mashing, boiling, fermentation and maturation.

#### 287 **4.5 Local Non-Alcoholic Beverage Production**

288 Millet drink (Kunu or Kunun-zaki) is a non-alcoholic, non-carbonated and refreshing cereal  
289 beverage popular in Northern Nigeria and is becoming widely consumed in the South [72]. It  
290 serves as breakfast drink, appetizer, weaning food and is also medicinal [73]. Kunu, is a  
291 nutritious non-alcoholic drink that is produced from various cereal grains such as millet,  
292 sorghum, maize and rice. Kunu is a drink that has found great appeal in the northern part of  
293 Nigeria, its consumption is spread over every class of personality and it is consumed either as a  
294 food supplement or thirst quencher. Kunu is cheaply available and serve as an alternative to  
295 carbonated drinks products which have little or no nutritional benefits to its consumers. Kunu is

296 one of the complex mixtures which contain macromolecules such as protein, carbohydrates and  
297 lipids [74]. The major important cereals which are used in the preparation of kunu are millet,  
298 maize, guinea corn and rice. During the preparation of kunu, the ingredients needed are ginger  
299 (*Zingiber officinals*), alligator pepper (*Afromonium melegueta*), red pepper (*Capsicum species*),  
300 black pepper (*Piper guineense*) and kakandoru or eru. All these ingredients perform one function  
301 or the other in the course of the preparation. The most abundant constituents of kunu is water and  
302 it acts as the medium in which all other constituents are dissolved and contain only traces amount  
303 of in-organic substances. The high nutritive value of kunu is attributed to the presence of protein,  
304 carbohydrates and particularly, vitamin B [75]. Kunu is widely accepted as food drink in some  
305 urban centres especially in the Hausa land. The quality and quantity of the products depend  
306 largely on the quality of the ingredients and handling technique in the course of production by  
307 the producer. The product could be obtained quantitatively after 2 days and it could be stored for  
308 another 3 days when refrigerated [75]. It has however been reported that, if kunu is kept  
309 overnight in hot season without being refrigerated, its quality begins to deteriorate and this may  
310 lead to the spoilage which when consume could constitute danger to health [76]. Spoilage of this  
311 product from observation occurs from improper handling, constant fermentation of the  
312 ingredients especially the carbohydrates and enzymatic action on the substrates [75]. Hence there  
313 is need for proper formulation and carbonation of the product. Carbonated drinks are desired and  
314 preferred because of it sharp, unique and refreshing taste. Carbonated drinks are non-alcoholic  
315 beverages that consist of CO<sub>2</sub>, water, flavoring and some other types of sweet syrup [77]. The  
316 CO<sub>2</sub> when introduced increases the acidity level of the drink, thereby keeping some micro-  
317 organisms from growing. Thrive of microbes in a drink is what usually reduces the shelf-life of  
318 the product [78]. Carbonated millet beverages (kunu) are expected to make a lot of difference

319 when compared to other available carbonated drinks because of its nutritive values among many  
320 other properties which include:

- 321 i. Its ability to aid digestion and absorption of components into the body system.
- 322 ii. Boosting the immune system of the body against microbial attack.

323 Currently in Nigeria, carbonated drinks are very expensive to buy. A bottle of 50Cl costs an  
324 average price of N100.00 and these have little or no nutritive value because they contained high  
325 concentration of sugar and artificial concentrates. Kunu however seem to be highly nutritious  
326 with relatively low cost of production and consumption. It is being prepared from our local  
327 cereals which are very common and are part of our staple food substances. The problem facing  
328 the satisfaction derived from kunu comes from its fast deterioration due to microbial activities  
329 causing its spoilage.

330 To produce kunu, millet grain will be cleaned and steeped in twice its volume of water for 24 h.  
331 Thereafter the steeped grains will be washed and spices added. The spices added will include  
332 ginger, red pepper, cloves and black pepper. The steeped millet grains and spices will then be  
333 wet milled in a grinding machine and sieved to remove the shafts after which the supernatant will  
334 be decanted from the slurry. The slurry will be divided into two equal halves with one half added  
335 to boiling water while stirring for 5 minutes, cooled to a temperature of 35<sup>o</sup>C and subsequently  
336 added to the remaining half slurry. Adequate amount of water will be added to the mixture,  
337 stirred and left to settle. After which, the mixture will be sieved using a muslin cloth and the  
338 filtrate sweetened with granulated sugar and mixed properly to obtain the freshly processed  
339 millet beverage. The product will be bottled in plastic bottles.

340

341

342 **5.0 Conclusion**

343 In this study the composition, functional properties and food application of millet was  
344 extensively examined. The review work showed high nutritional composition of this cereal  
345 grains (millet). The behavioral pattern was also discussed when used industrially which portrays  
346 their functional capabilities. The review showed that the product processes wide food  
347 applications as also serve as good functional abilities that could help to promote human health.  
348 However, research needs to focus on improving its shelf-life for industrial production.

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363



364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386

## References

1. Shahidi F, Chandrasekara A. Bioaccessibility and antioxidant potential of millet grain phenolics as affected by simulated in vitro digestion and microbial fermentation. *Journal of Functional Foods*. 2012; 4:226–37.
2. Sireesha Y, Kasetti RB, Swapna SAS, Apparao C. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology*. 2011; 18(2):159–64.
3. Pradeep SR, Guha G. Effect of processing methods on the nutraceutical and antioxidant properties of little millet (*Panicum sumatrense*) extracts. *Food Chemistry*. 2011; 126:1643–7.
4. Devi PB, Vijayabharathi RS, Sathyabama NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *Journal of Food Science and Technology*. 2011; 11:584-9.
5. Kumar KK, Parameshwaran PK. Characterization of storage protein from selected varieties of foxtail millet (*Setaria italica* (L) Beauv). *Journal of Agricultural Science*. 2016; 77:535–42.
6. Siwela M, Taylor JRN, de-Milliano WAJ, Duodu KG. Influence of phenolics in finger millet on grain and malt fungal load, and malt quality. *Food Chemistry*. 2010; 121(2):443–9.
7. Songr'e-Ouattara LT, Mouquet-Rivier C, Icard-Vernière C, Humblot C, Diawara B, Guyot JP. Enzyme activities of lactic acid bacteria from a pearl millet fermented gruel (ben-saalga) of functional interest in nutrition. *International Journal of Food Microbiology*. 2008; 128(2):395-400.

- 387 8. Jideani VA, Nkama I, Agbo EB, Jideani IA. Fura production in some northern states of  
388 Nigeria—A survey. *Plant Foods for Human Nutrition*. 2012; 56:23.
- 389 9. Ramashia SE, Anyasi TA, Gwata ET, Meddows-Taylor S, Jideani AIO. Processing,  
390 nutritional composition and health benefits of finger millet in sub-saharan Africa. *Food*  
391 *Science and Technology*. 2019; 39(2): 253-266.
- 392 10. FAO (Food and Agricultural Organization). 2012. Economic and Social Department: The  
393 Statistical Division. Statistics Division 2012.
- 394 11. Chandrasekara A, Naczk M, Shahidi F. Effect of processing on the antioxidant activity of  
395 millet grains. *Food Chemistry*. 2012; 133:1–9.
- 396 12. Jideani AI, Jideani VA. Traditional and Possible Technology uses of *Digitaria exillis*  
397 (*Acha*) and *Digitaria ibrua*: a review. *Plant Foods for Human Nutrition*. 2011; 54:362-  
398 373.
- 399 13. Kaur KD, Jha A, Sabikhi L, Singh AK. Significance of coarse cereals in health and  
400 nutrition: a review. *Journal of Food Science and Technology*. 2014; 11:612-9.
- 401 14. Krishnan RM, Dharmaraj U, Malleshi NG. Influence of decortication, popping and  
402 malting on bioaccessibility of calcium, iron and zinc in finger millet. *Food Science and*  
403 *Technology*. 2012; 48:169–74.
- 404 15. Truswell AS. Cereal grain and coronary heart disease. *European Journal of Clinical*  
405 *Nutrition*. 2002; 56(1):1–4.
- 406 16. Gupta N, Srivastava AK, Pandey VN. Biodiversity and nutraceutical quality of some  
407 indian millets. *Proceedings of the National Academy of Sciences, India Section B:*  
408 *International Journal of Advanced Research in Biological Sciences*. 2012; 20(4): 1555-  
409 1561.

- 410 17. Bouis HE. Enrichment of food staples through plant breeding: a new strategy for fighting  
411 micronutrient malnutrition. *Nutrition*. 2000; 16:701–4.
- 412 18. Pradhan A, Nag SK, Patil SK. Dietary management of finger millet (*Eleusine coracana*  
413 *L. Gaerth*) controls diabetes. *Current Science*. 2010; 98(6):763–5.
- 414 19. Verma S, Srivastava S, Tiwari N. Comparative study on nutritional and sensory quality  
415 of barnyard and foxtail millet food products with traditional rice products. *Journal of*  
416 *Food Science and Technology*. 2015; 52(8):5147–5155.
- 417 20. Puranik S, Kam J, Sahu PP, Yadav R, Srivastava RK, Ojulong H, Yadav R. Harnessing  
418 Finger Millet to Combat Calcium Deficiency in Humans: Challenges and Prospects.  
419 *Frontiers in Plant Science*. 2017; 8:1311.
- 420 21. Saha S, Gupta A, Singh, SRK, Bharti N, Singh KP, Mahajan V, Gupta HS.  
421 Compositional and varietal influence of finger millet flour on rheological properties of  
422 dough and quality of biscuit. *Food Science Technology*. 2011; 44(3):616–21.
- 423 22. Singh P, Raghuvanshi RS. Finger millet for food and nutritional security. *African Journal*  
424 *of Food Science*. 2012; 6(4):77–84.
- 425 23. Ahmed SM, Saleh QZ, Jing C, Qun S. Millet Grains: Nutritional Quality, Processing, and  
426 Potential Health Benefits. *Comprehensive Reviews in Food Science and Food Safety*.  
427 2013; 12:280-292.
- 428 24. Mal B, Padulosi S, Ravi SB. Minor millets in South Asia: learnings from IFAD-NUS  
429 Project in India and Nepal. Maccaresse, Rome, Italy: Bioversity Intl and Chennai, India:  
430 M.S. Swaminathan Research Foundation. 2010; 1–185.

- 431 25. Singh KP, Mishra A, Mishra HN. Fuzzy analysis of sensory attributes of bread prepared  
432 from millet-based composite flours. *LWT—Food Science and Technology*. 2012;  
433 48:276–82.
- 434 26. Sreeramulu D, Vijaya CKR, Raghunath M. Antioxidant activity of commonly consumed  
435 cereals, millets, pulses and legumes in India. *Indian Journal of Biochemistry and*  
436 *Biophysics*. 2009; 46:112-115.
- 437 27. Gyan-Chand K, Morya V, Mishra HS, Shakya S, Raj B, Yadav KN. *International Journal*  
438 *of Advanced Ayurveda, Yoga, Unani, Siddha and Homeopathy*. 2017; 6(1): 447-452.
- 439 28. Mohamed TK, Zhu K, Issoufou A, Fatmata T, Zhou H. Functionality, *in vitro*  
440 digestibility and physicochemical properties of two varieties of defatted foxtail millet  
441 protein concentrates. *International Journal of Molecular Science*. 2009; 10:522–438.
- 442 29. Azad MOK, Jeong DI, Adnan M, Salitxay T, Heo JW, Naznin MT, Lim JD, Cho DH,  
443 Park BJ, Park CH. Effect of Different Processing Methods on the Accumulation of the  
444 Phenolic Compounds and Antioxidant Profile of Broomcorn Millet (*Panicum miliaceum*  
445 L.) Flour. *Foods*. 2019; 8:230-243.
- 446 30. Rathore T, Singh R, Kamble DB, Upadhyay A, Thangalakshmi S. Review on finger  
447 millet: Processing and value addition. *The Pharma Innovation Journal*. 2019; 8(4): 283-  
448 291.
- 449 31. Gupta R. RAGI, a boon to nutritional security. Indian Agriculture Research Institute Pusa  
450 Campus, New Delhi-110012. 2014.
- 451 32. Glew RS, Chuang LT, Roberts JL, Glew RH. Amino acid, fatty acid and mineral content  
452 of black finger millet (*Eleusine coracana*) cultivated on the Jos Plateau of Nigeria.  
453 *Foods*. 2008; 2(2):115–8.

- 454 33. Hegde PS, Chandra TS. ESR spectroscopic study reveals higher free radical quenching  
455 potential in kodo millet (*Paspalum scrobiculatum*) compared to other millets. Food  
456 Chemistry. 2005; 92:177–82.
- 457 34. Kalinova J, Moudry J. Content and quality of protein in proso millet (*Panicum miliaceum*  
458 *L.*) varieties. Plant Foods for Human Nutrition. 2006; 61:45–9.
- 459 35. Subramanian S, Viswanathan R. Bulk density and friction coefficients of selected minor  
460 millet grains and flours. Journal of Food Engineering. 2007; 81(1):118–26.
- 461 36. Liu J, Tang X, Zhang Y, Zhao W. Determination of the volatile composition in brown  
462 millet, milled millet and millet bran by gas chromatography/ mass spectrometry.  
463 Molecules. 2012; 17:2271–82.
- 464 37. Khader V. Textbook of Food Science and Technology. ICAR, New Delhi. Khatkar BS  
465 (2005). Trends in cereal processing. Bev. Food World. 2001; 32(11):65-67.
- 466 38. Guo XX, Sha XH, Rahman E, Wang Y, Ji B, Wu W, Zhou F. Antioxidant capacity and  
467 amino acid profile of millet bran wine and the synergistic interaction between major  
468 polyphenols. Journal of Food Science and Technology. 2018; 55(3):1010–1020.
- 469 39. Kaur M, Singh N. Relationships between selected properties of seeds, flours, and starches  
470 from different chickpea cultivars. International Journal of Food Properties. 2006; 9:597-  
471 608.
- 472 40. Siddiq M, Nasir M, Ravi R, Dolan KD, Butt MS. Effect of defatted maize germ addition  
473 on the functional and textural properties of wheat flour. International Journal of Food  
474 Properties. 2009; 12:860-870.

- 475 41. Thilagavathi T, Kanchana S, Banumathi P, Hemalatha G, Vanniarajani C, Sundar M,  
476 Ilamaran M. Physico-chemical and functional characteristics of selected millets and  
477 pulses. Indian Journal of Science and Technology. 2015; 8(S7), 147–155.
- 478 42. Butt MS, Batool R. Nutritional and functional properties of some promising legumes  
479 proteins isolates. Pakistan Journal of Nutrition. 2010; 9(4):373–379.
- 480 43. John EK, Nicholas M. Functional properties of proteins in foods: A survey. CRC critical  
481 reviews in food science and nutrition. 2009; 7(3):201-206.
- 482 44. Amir G, Kamlesh P, Pradyuman K. Physico-chemical, Functional and Antioxidant  
483 Properties of Millet Flours, Conceptual Frame Work and Innovations in Agroecology  
484 and Food Sciences. 2016; 585-589.
- 485 45. Yadahally N, Vadakkoot B, Vishwas M. Nutritional implication and flour functionality of  
486 popped/ expanded horse gram. Journal of Food Chemistry. 2008; 108:891- 899.
- 487 46. Krishnan R, Dharmaraj U, Sai MR, Malleshi NG. Quality characteristics of biscuits  
488 prepared from finger millet seed coat based composite flour. Food Chemistry. 2011;  
489 129:499–506.
- 490 47. Appiah F, Asibuo JY, Kumah P. Physicochemical and functional properties of bean  
491 flours of three cowpea (*Vigna unguiculata* L. Walp) varieties in Ghana. African Journal  
492 of Food Science. 2011; 5(2): 100 – 104.
- 493 48. Yagoub A, Abdalla AA. Effect of domestic processing methods on chemical  
494 composition, *in vitro* digestibility of protein and starch and functional properties of millet  
495 seed. Research Journal of Agriculture and Biological Science. 2007; 3:24–34.

- 496 49. Akubor PI. Effect of Processing Methods on the Chemical Composition and Functional  
497 Properties of Pigeon Pea Seed. *Asian Journal of Advances in Agricultural Research*.  
498 2017; 2(2): 1-8.
- 499 50. Jayathilake C, Visvanathan R, Deen A, Bangamuwage R, Jayawardana BC, Nammic S,  
500 Liyanage R. Cowpea: an overview on its nutritional facts and health benefits. *Journal of*  
501 *the Science of Food and Agriculture*. 2018; 20(1):8-15.
- 502 51. Udensi E, Okoronkwo K. Effects of fermentation and germination on the  
503 physicochemical properties of *Mucuna cochinchinensis* protein isolate. *African Journal of*  
504 *Biotechnology*. 2006; 5:896–900.
- 505 52. Iwe MO, Michael N, Madu NE, Obasi NE, Onwuka GI. Physicochemical and Pasting  
506 Properties High Quality Cassava Flour (HQCF) and Wheat Flour Blends.  
507 *Agrotechnology*. 2017; 6: 167.
- 508 53. ARSO (2012). African Organization for Standardization. High Quality Cassava Flour  
509 Specification CD-ARS 840 First Edition 2012 Reference No. ARS 840:2012(E) ICS  
510 67.080.20.
- 511 54. Eleazu OC, Eleazu KC, Kolawole S. Use of indigenous technology for the production of  
512 high quality cassava flour with similar food qualities as wheat flour. *Acta Scientiarum*  
513 *Polonorum Technologia Alimentaria*. 2014; 13: 249-256.
- 514 55. Chandra S, Samsher V. Assessment of functional properties of different flours. *African*  
515 *Journal of Agricultural Research*. 2013; 8(38): 4849-4852.
- 516 56. Sanni LO, Ikuomola DP, Sanni SA. Effect of length of fermentation and varieties on  
517 the quality of sweet potato gari. *Proceedings of the 8th Triennial Symposium of the*  
518 *International Society for Tropical Root Crops Africa Branch*. 2001; 3:208-211.

- 519 57. Bhupender SK, Rajneesh B, Baljeet SY. Physicochemical, functional, thermal and  
520 pasting properties of starches isolated from pearl millet cultivars. African Journal of  
521 Biotechnology. 2013; 5:96–103.
- 522 58. Zhang X, Tong Q, Zhu W, Ren F. Pasting, rheological properties and gelatinization  
523 kinetics of tapioca starch with sucrose or glucose. Journal of Food Engineering. 2013;  
524 114:255–261.
- 525 59. Yadav BS, Yadav RB, Kumar M. Suitability of pigeon pea and rice starches and their  
526 blends for noodle making. Food Science and Technology. 2011; 44: 1415-1421.
- 527 60. Alsuhaibani AMA. Rheological and Nutritional Properties and Sensory Evaluation of  
528 Bread Fortified with Natural Sources of Calcium. Journal of Food Quality. 2018;  
529 20(1):1-8.
- 530 61. Karim AA, Norziah MH, Seow CC. Methods for study of starch retrogradation. Food  
531 Chemistry. 2000; 71: 9-36.
- 532 62. Shifeng Y, Ying M, Lucile M, Da-Wen S. Physicochemical properties of starch and flour  
533 from different Rice cultivars. Food and Bioprocess Technology. 2010; 5(2):626-637.
- 534 63. Shobana S, Malleshi NG. Preparation and functional properties of decorticated finger  
535 millet (*Eleusine coracana*). Journal of Food Engineering. 2007; 79(2):529–38.
- 536 64. Odusola KB, Ilesanmi FF, Akinloye OA, Ilesanmi OS, Zaka KO. Nutrient composition  
537 and acceptability of cereal gruel (Ogi) produced from sorghum (*Sorghum bicolor*)  
538 supplemented with soybeans (*Glycine max*) and Ginger (*Zingiber officinale*), A  
539 publication of the School of Agriculture Technology, The Federal University of  
540 Technology, Akure, Nigeria. 2015.



- 541 65. Jideani VA, Wedzicha BL. Shelf life and predominating micro-flora of a non fermented  
542 pearl millet dough (fura). Journal of Food Science. 2001; 11:83.
- 543 66. Agu HO, Anosike AN, Jideani IA.. Physicochemical and microbial qualities of  
544 assessment of the isolates for industrial potential. Bioresources Technology. 2007;  
545 98:1380.
- 546 67. Khetarpaul N, Goyal R, Garg R. Nutritional and sensory evaluation of nutritious porridge  
547 prepared using combinations of soy and sorghum grits. Nutrition and Health. 2004; 17(4):  
548 309-315.
- 549 68. Togo CA, Feresu SB, Mutukumira AN. Identification of Lactic Acid Bacteria from  
550 opaque beer for potential use as a starter culture. Journal of Food Technology. 2002;  
551 7:93-97.
- 552 69. Mbajuka D, Omeh YS, Ezeja MI. Fermentation of sorghum using yeast (*Saccharomyces*  
553 *cerevisiae*) as a starter culture for Burukutu production. Continental Journal of Biological  
554 Science. 2010; 3: 63-74.
- 555 70. Maoura N, Mbaiguinam M, Nguyen HV, Gaillardin C, Pourquoi J. Identify and typing of  
556 yeast strains isolated from *billi billi*, a traditional sorghum beer of Chad. African Journal  
557 of Biotechnology. 2005; 4: 646-656.
- 558 71. Lyumugabe F, Kamaliza G, Thonart B. Microbiological and physico-chemical  
559 characteristics of Rwandese traditional beer *Ikage*. African Journal of Biotechnology.  
560 2010;9 (27): 4241 – 4246.
- 561 72. Sowonola OA, Tunde-Akintunde TY, Adedeji F. Nutritional and sensory qualities of  
562 soymilk *kunun* blends. African Journal of Food and Nutrition Science. 2005; 5(2): 2-5.

- 563 73. Akoma O, Jiya EA, Akumka DD, Mshelia E. Influence of malting on the nutritional  
564 characteristics of *kunu-zaki*. African Journal of Biotechnology. 2006; 5(10): 996-1000.
- 565 74. Gaffa T, Jideani IA, Nkama I. Traditional Production, Consumption and Storage of  
566 Kunu- a non-alcoholic cereal beverage. Plant Foods for Human Nutrition. 2002; 57(1):  
567 73-81.
- 568 75. Wakil S. Isolation and Screening of Antimicrobial Producing Lactic Acid. Indian Journal  
569 of Science and Technology, 2004; 8(S7):147–155.
- 570 76. Adebayo GB, Otunola GA, Ajao TA. Physicochemical, Microbiological and Sensory  
571 Characteristics of Kunu Prepared from Millet, Maize and Guinea Corn and Stored at  
572 Selected Temperatures. Advance Journal of Food Science and Technology. 2010;  
573 2(1):41-46.
- 574 77. Abdulkareem AS, Uthman H, Jimoh A. Development and Characterization of a  
575 Carbonated Ginger Drink. Department of Chemical Engineering. Federal University of  
576 Technology, Minna. 2011.
- 577 78. Julio A. Fermented Foods and Beverages of the North-East India. Food Processing, Dried  
578 Food, Frozen Food, Juice and Beverage. 2011.