

# The Effects of Chicken Manure Application Rates on Growth and Yield of Swiss Chard

(*Beta vulgaris* var. *cicla* L.)

## ABSTRACT

Swiss chard (*Beta vulgaris* var. *cicla*) is a leafy vegetable that belongs to the *Chenopodiaceae* family. The leaves are cooked, if still tender they are used in salads. Over the years, Swazis have adopted the use of inorganic fertilisers as they are easy to apply and come with recommended application rates. However, their main drawback is that they are environmentally unfriendly especially when washed into rivers, streams and other water bodies. For this cause, the use of animal manures has been promoted. Four week old Swiss chard seedlings were transplanted on the 4<sup>th</sup> of February, 2016 in 1.5 x 1.5 m plots with an inter and intra row spacing of 45 cm and they were irrigated twice a day during the first week and every second day from the second week until the end of the experiment. The experiment was conducted at the Horticulture Department Farm, Faculty of Agriculture and Consumer Sciences, Luyengo Campus of the University of Swaziland to determine the effects of chicken manure application rates on growth, yield and quality of Swiss chard. Four chicken manure application rates (10, 20, 40 and 80 t/ha) and a recommended 900kg/ha, inorganic basal fertiliser with a 125kg/ha LAN top dressing fertiliser used as a control. A Randomised Complete Block Design (RCBD) with four replicates was used. The study showed that the application of 80 t/ha of chicken manure improved the growth and yield of Swiss chard. It is recommended that farmers may use 80 t/ha of chicken manure because it gave the best results compared to the other treatments.

23

24

25

## 26 1.0. INTRODUCTION

27 Swiss chard is leafy vegetable that belongs to the *Chenopodiaceae* family and is scientifically  
28 known as *Beta vulgaris var. cicla*. It belongs to the same family as beetroot and mangel-wurzel.  
29 Unlike beetroot and mangel-wurzel, Swiss chard lacks the large bulbous tap root. It is one of the  
30 most nutritious vegetable crops in the world. Swiss chard may be grown in Eswatini/ Swaziland  
31 all year round, in all the ecological zones.

32 Swiss chard forms part of the several leafy green vegetables that are known as 'greens.' It is a  
33 biennial plant with large dark leaves. The leaves are large, glossy and crispy and can grow up to  
34 37 cm long and 25 cm wide [1]. Stalks of Swiss chard come in a variety of colours depending on  
35 the cultivar, they are usually white, yellow, orange or red [2]. The first records of cultivation  
36 place the origin of Swiss chard in the Mediterranean region particularly Italy and was first  
37 written about by the Greek philosopher Aristotle in 4 B.C [1]. Swiss chard is a short day (SD)  
38 plant with critical day length of 12 hours. It grows best at temperatures ranging from 7 to 24 °C.  
39 Swiss chard can withstand light frosts but an extended exposure to temperatures less than 5°C  
40 induces bolting. In hot weather, the leaves remain small and are of inferior quality [3]. Leaves of  
41 Swiss chard are harvested usually within eight weeks from sowing and once they are in good size  
42 [2]. Harvesting is done continuously so that the leaves do not stay long and lose their colour or  
43 become tough.

44

45 The use of inorganic fertilisers has resulted in residual toxicities and degradation of the soil  
46 structure. These inorganic fertilisers become an environmental threat to aqua life when washed  
47 into rivers, streams and other water bodies. They are relatively expensive such that not all farmers  
48 afford them. As a result some farmers produce Swiss chard below the expected optimum level.

49 The main objective of this study was to improve the production of Swiss chard and to contribute  
50 towards food security and income generation in Eswatini/Swaziland. The specific objective was  
51 to determine the optimum level of chicken manure application on growth, yield and quality of  
52 Swiss chard.

53

54

## 55 2.0 MATERIAL AND METHODS

### 56 2.1 Experimental site

57 The experiment was conducted at the Horticulture Department Farm, Faculty of Agriculture and  
58 Consumer Sciences, Luyengo Campus of the University of Swaziland. The farm is located at  
59 Luyengo, Manzini region, in the Middleveld agro-ecological zone. Luyengo is located at latitude  
60  $26^{\circ}4''$  S and longitude  $31^{\circ}4''$  E. The average altitude of this area is 750 m above sea level. The  
61 mean annual precipitation is 980 mm with most of rain falling between October and April.  
62 Drought hazard is about 40%. The average summer temperature is  $27^{\circ}\text{C}$  and winter temperature  
63 is about  $15^{\circ}\text{C}$ . The soils of Luyengo are classified under Malkerns series. They are ferrasolic or  
64 merely a ferralitic soil integrated to fersialitic soils or typical ultisols. The soil in the  
65 experimental area was a sandy loam [4].

## 66 2.2 Plant Materials

67 Four **week** old Swiss chard seedlings were obtained from Greenhouse Seedlings, Ezulwini. They  
 68 were transplanted on the 4<sup>th</sup> of February, 2016 in 1.5 x1.5 m plots with an inter and intra row  
 69 spacing of 45 cm and they were irrigated twice a day during the first week and every second day  
 70 from the second week until the end of the experiment.

## 71 2.3 Experimental Design

72 Four chicken manure application rates (10, 20, 40 and 80 t/ha) and a recommended 900kg/ha,  
 73 inorganic basal fertiliser with a 125kg/ha LAN top dressing fertiliser was used as a control  
 74 (Table 1). A Randomised Complete Block Design (RCBD) with four replicates was used. Each  
 75 plot had four rows and there were four plants in each row which gave a total of 320 plants used  
 76 **in** the experiment.

77 Table 1: Treatment descriptions.

Treatment code	Treatment
1	80 t/ha
2	40 t/ha
3	20 t/ha
4	10 t/ha
5	900 kg of 2:3:2 (22) and 125 kg of LAN (28)

78

## 79 2.4. Soil analysis

80 Soil chemical properties were analysed at the Soil Chemistry laboratory of the University of  
 81 Swaziland, Luyengo Campus.

## 82 **2.5. Manure analysis**

83 **Chicken manure** chemical properties were analysed at the soil Chemistry laboratory of the  
84 University of Swaziland, Luyengo Campus.

## 85 **2.6. Data collection**

86 Data was collected weekly, from the second week after transplanting. Five plants were randomly  
87 selected in each plot for data recording. Data was collected on the following growth parameters:  
88 plant height, number of leaves and leaf area while leaf area index was calculated. The fresh mass  
89 and dry mass of the Swiss chard were measured after harvesting.

## 90 **2.7. Growth parameters**

### 91 **2.7.1. Plant height**

92 Five plants were randomly selected per plot and plant height was measured from the base of the  
93 plant to the leaf apex (tip) **using a 30 cm ruler.**

### 94 **2.7.2. Number of leaves**

95 The number of leaves per plant was determined by physically counting all the leaves on each  
96 selected plant. Five plants were selected per plot and it was done on a weekly basis, which was at  
97 week 3, 4, and 5 after transplanting.

### 98 **2.7.3. Leaf area**

99 The leaf area of the Swiss chard was determined by multiplying the leaf width and leaf length  
100 and then multiplying the product by 0.75 (correction factor) [5]. **(Edje and Ossom,** It was  
101 expressed in  $\text{cm}^2$ .

### 102 **2.7.4. Leaf area index**

103 The leaf area index was determined by dividing the leaf area in  $\text{cm}^2$  by the area occupied by a  
104 single plant in  $\text{cm}^2$ . **[5].Edje and Ossom**

### 105 2.7.5. Fresh and dry mass

106 This was determined at the end of the cropping season by weighing the harvested leaves per plot.  
 107 Five plants per plot were used to determine the fresh and dry mass in this experiment. The plants  
 108 were randomly selected per plot and their shoot fresh mass was measured using a digital scale  
 109 balance. They were then oven dried at a temperature of 72°C for 72 hours to determine their  
 110 shoot dry mass[5].

### 111 2.8. Data analysis

112 The data collected was subjected to analysis of variance (ANOVA) using MSTAT-C statistical  
 113 package, Version 1.4[6]. Where significant differences were detected mean separation were  
 114 performed using Duncan's New Multiple Range Test (DNMRT) at 5 % probability level [7].

## 115 3.0 RESULTS

### 116 3.1. Soil analysis

117 Soil chemical properties were analysed at the Chemistry Laboratory of the University of  
 118 Swaziland, Luyengo Campus. The results of the soil chemical properties are shown in Table 2.

119 Table 2: Soil analysis

Soil parameter	Value
Soil pH	5.8
Phosphorus (mgP/kg)	39.56
Potassium (cmolc/kg)	1.54

120

121 **3.2. Manure analysis**


122 Chemical properties of the chicken manure were analysed at the Chemistry Laboratory of the  
 123 University of Swaziland, Luyengo Campus. The results of the chemical properties of chicken  
 124 manure are shown in Table 3.

125 Table 3: Chicken manure analysis

Manure parameter	Value
pH	7.2
Phosphorus(mgP/kg)	17
Potassium(cmolc/kg)	1 895
Magnesium	Not determined-

126 **3.3. Plant height**

127 Plant height of spinach was significantly ( $P < 0.05$ ) different among the different treatments. The  
 128 highest plant height (34.6 cm) was obtained in spinach treated with 80 t/ha of chicken manure  
 129 while the lowest plant height (26.0 cm) was obtained in Swiss chard plants treated with 10 t/ha  
 130 of chicken manure (Figure 1). The plant height of Swiss chard plants treated with inorganic  
 131 fertilizers was higher (28.1 cm) but not significantly ( $P > 0.05$ ) different from those treated with  
 132 10 t/ha of chicken manure (26.0 cm) (Figure 1).




133

134 Figure 1: Effects of chicken manure on **Swiss chard** plant height. Vertical bars are standard  
 135 error (se) below and above the mean.

#### 136 3.4. Number of leaves

137 The number of leaves per plant were not significantly ( $P>0.05$ ) different among the **Swiss chard**  
 138 plants. The highest number of leaves (7.6) was obtained in plants treated with 80 t/ha of chicken  
 139 manure while the lowest number of leaves (6.4) was obtained in plants treated with 40 t/ha of  
 140 chicken manure (Figure 2).



141


142 Figure 2: Effects of chicken manure on number of leaves of **Swiss chard**. Vertical bars are  
 143 standard error (se) below and above the mean.

144

### 145 3.5. Leaf area

146 The leaf area of **Swiss chard plants** was significantly ( $P < 0.05$ ) different among treatments. The  
 147 highest leaf area ( $291.9 \text{ cm}^2$ ) was obtained in plants treated with 80 t/ha of chicken manure while  
 148 the lowest leaf area ( $169.8 \text{ cm}^2$ ) was obtained in plants treated with 10 t/ha of chicken manure at

149 5 WAT (Figure 3). The leaf area of **Swiss chard** increased with increasing application rates of  
 150 chicken manure.



151

152 Figure 3: Effects of chicken manure on the leaf area per plant of **Swiss chard**. Vertical bars are  
 153 standard error (se) below and above the mean.


### 154 3.6. Leaf area index

155 The leaf area index(LAI)was significantly ( $P<0.05$ ) different among treatments. The highest  
 156 LAI (1.1) was obtained in plants treated with 80 t/ha of chicken manure while the lowestLAI

157

158

159 (0.5) was obtained in plants treated with 10 t/ha of chicken manure at 5 WAT (Figure 4). The  
 160 leaf area of **Swiss chard** increased with increasing application rates of chicken manure.



161

162 Figure 4: Effects of chicken manure on the LAI per plant of **Swiss chard**. Vertical bars are  
 163 standard error (se) below and above the mean.


164

### 165 3.7. Fresh mass and dry mass

166 There was a significant ( $P < 0.05$ ) difference in the fresh shoot mass of **Swiss chard** plants among  
 167 treatments (Figure 5). The highest fresh shoot mass (237.5 g) was obtained in plants treated with

168 80 t/ha of chicken manure while the lowest fresh shoot mass (100.0 g) was obtained in plants  
 169 fertilised with inorganic fertilisers. **There was no significant difference in Swiss chard fresh maas**  
 170 **amended with 40t/ha or 80 t/ha chicken manure.**

171 There was a significant ( $P < 0.05$ ) difference in the dry shoot mass of **Swiss chard** plants **among**  
 172 the different **treatments** (Figure 5). The highest dry shoot mass (20.4 g) was achieved at 80 t/ha  
 173 of chicken manure while the lowest dry shoot mass (10.1 g) was obtained in **Swiss chard** plants  
 174 treated with inorganic fertilisers.



175  
 176 Figure 5: Effects of chicken manure on fresh and dry shoot mass of **Swiss chard** at week 5 after  
 177 transplanting. Bars followed by the same alphabet **are** not significantly different from  
 178 one another at  $P = 0.05$ . Mean separation by Duncan's New Multiple Range Test.

179

## 180 4.0. DISCUSSION

181 Different application rates of chicken manure had varying effects on growth, yield and quality of  
182 **Swiss chard**. Plants treated with 80 t/ha of chicken manure performed better in terms of growth  
183 in comparison with the other treatments. These **Swiss chard** plants had the highest plant height,  
184 number of leaves, fresh shoot mass, dry shoot mass, leaf area and leaf area index compared to  
185 spinach treated with 10, 20, 40 t/ha of **chicken** manure and application of inorganic fertilisers  
186 recommended for **Swiss chard** production. **Swiss chard** plants treated with 10 t/ha had the lowest  
187 plant height, leaf area and leaf area index. The highest number of leaves of **Swiss chard** plants  
188 from the highest application rate of chicken manure must have been as a result of relatively high  
189 amounts of nitrogen [8]. It was also noted that plant height, number of leaves, leaf area, leaf area  
190 index, fresh and dry shoot mass increased with increasing levels of chicken manure. These  
191 results are in agreement with those of [8]. who studied the effects of an organic fertilizer (cattle  
192 manure) on *Zea mays*. As chicken manure application rate was increased, the availability of plant  
193 nutrients in the soil also increased. This resulted in the increase of growth and yield.

194 Chicken manure at 80 t/ha performed better in comparison with inorganic fertilizers. These  
195 findings do not deviate much from those obtained by [9] Owen (2008) who reported that  
196 synthetic fertilisers do not have good characteristics in aggregating soil particles. The plants  
197 treated with inorganic fertilisers gave a lower yield than those treated with 80 t/ha of chicken  
198 manure. Animal manures have beneficial effects on physical and chemical properties of soil and  
199 therefore have the ability to retain water, supply macro- and trace elements absent in inorganic  
200 fertilisers. Increased vegetable yield with the use of manure have been previously reported for  
201 okra [10]. The benefits of organic fertilizer use in vegetable production has previously been  
202 reported [11,12,13] and very recently reported [14,15,16] in the Kingdom.

## 203 5.0. CONCLUSION AND RECOMMENDATION

204

205 The study showed that the application of 80 t/ha of chicken manure improved the growth and  
206 yield of **Swiss chard**. From these findings, it can therefore be concluded that 80 t/ha was best for  
207 **[8]**. under the conditions of this study. **If fresh mass at the end is the most important parameter**  
208 **farmers could as well use 40 t/ha because there was no significant difference in plants amended**  
209 **with 80 t/ha.**

210 It is recommended that farmers may use 80 t/ha of chicken manure because it gave the best  
211 results compared to the other treatments.

212

## 213 REFERENCES

214 1.Pierce LC. Vegetables: Characteristics, Production, and Marketing. John Wiley and Sons.  
215 New York, New York, USA. . 1987.

216 2.Hadfield J. Vegetable Gardening in Central Africa. Purnell and Sons. Johannesburg, South  
217 Africa.1960.

218 3. Gilber, Z, Hadfield J. Down-to- Earth Fruit and Vegetable Gardening in South Africa. Struik  
219 Publishers. Cape Town, South Africa.1987.

220 4.Murdoch G. Soils and Land Capacity in Swaziland. Swaziland Ministry of Agriculture.  
221 Mbabane, Swaziland. 1970.

222 5.Edje OT, Ossom EM. **Crop Science Handbook**. Blue Moon Publishers. Manzini,  
223 Swaziland.2009.

- 224 6.Nissen,O. MSTAT-C. A micro Computer Program Design for Management and  
 225 Analysis of Agronomic Research Experiments.Michigan State University. East  
 226 Lansing, Michigan, USA. 1989.
- 227 7.Gomez KA, Gomez A A. Statistical Procedures for Agricultural Research. John Wiley  
 228 and Sons 2<sup>nd</sup> Edition. Singapore.(1984).
- 229 8.Lungu O I, Chirwa B. Effect of lime and farmyard manure on soil acidity from and  
 230 maize growth on alfisols from Zambia. Tropical Agric(Trinidad)1993; 70; 309-315.
- 231 9.Owen P. Origin and distribution of lettuce.  
 232 (2008).<http://www.calettuceresearchboard.org/Origin.html>. 10/04/2016
- 233 10.Ogunlela VB, Masarirambi MT, Makuza SM. Effects of cattle manure application on  
 234 pod yield and yield indices of Okra (*Abelmoschus esculentus* L. Moench) semi-arid and  
 235 subtropical environment. J Food, Agric Environ. 2005;3 (1):125-129
- 236 11. Masarirambi MT, Hlawe MM, Oseni TO, Sibiyi TE. Effects of organic fertilizers on growth,  
 237 yield, quality and sensory evaluation of red lettuce (*Lactuca sativa* L.) 'Veneza Roxa'. Agric and  
 238 Biol J North America2010;1(6):1319-1324
- 239 12. Masarirambi M T, Dlamini P., Wahome PK, Oseni TO. Effects of chicken manure on  
 240 growth, yield and quality of lettuce (*Lactuca sativa* L.) 'Taina' under a lath house in a semi-arid  
 241 sub-tropical environment. American Eurasian J Agric Environ Sci,2012;12 (3): 399-406
- 242 13. Msibi, B.M., Mukabwe WO, Manyatsi AM, Mhazo N Masarirambi MT. Effects of liquid  
 243 manure on growth and yield of spinach (*Beta vulgaris* var cicla) in a sub-tropical environment in  
 244 Swaziland.Asian J Agric Sci,2014;6(2): 40-47

245 14. Kunene TR, Masarirambi MT, Wahome PK, Oseni TO. Influence of kraal manure, chicken  
246 manure and inorganic fertilizer on growth, yields and post-harvest quality of pepper (*Capsicum*  
247 *annuum* L.) in a sub-tropical environment. *Asian J Advances Agric Res*, 2019;11(1):1-11

248 15. Mabuza TZ, Masarirambi M.T. Nxumalo KA, Wahome PK. Effects of different rates of  
249 cattle manure on growth, yield and quality of pepper (*Capsicum annum* L.) in a sub-tropical  
250 environment of Eswatini (Swaziland). *Asian J Advances Agric Res*, 2019;11(1):1-7

251 16. Dlamini SN, Masarirambi MT, Wahome P K, Oseni. T.O. The effects of organic fertilizers  
252 on the growth and yield of amaranth (*Amaranthus hybridus* L.) grown in a lath house. *Asian J*  
253 *Advances Agric Res*, 2020;12(1):1-10

254

255

256

257

258

UNDER PEER REVIEW