

**STUDIES ON THE PHYTOCHEMICAL COMPOUNDS IN THE ETHANOLIC LEAF EXTRACT (ELE), ETHANOLIC BARK EXTRACT (EBE) AND ETHANOLIC ROOT EXTRACT (ERE) OF *Bridelia ferruginea* BENTH (EUPHORBIACEAE).**

**ABSTRACT**

*The phytochemical compounds of Bridelia ferruginea plant parts was carried out using qualitative method to determine the bioactive compounds present in the plant leave, stem bark and root extracts. The samples was weighed, of which 100g each of the powder were extracted in solvents (ethanol) 1000 ml macerated and stand for 72 hours. The solvents contained in the maceration bottle was decanted and filtered using a filter paper, the filtration was aided using a suction pump. The filtrate was concentrated using a rotary evaporator and then transferred into thermostatic water cabinet (temperature was set at 45°C), allowed to dry completely. The plant parts extracts were separately kept in a screw capped bottle for further research. The bioactive compound in the plant parts were detected. The result revealed that Carbohydrates, Saponins, Flavonoids, Tannins, Cardiac Glycosides, fats and oils were present. Alkaloid present in Dragendoff's test in all plant parts extract but absent in Mayer's test in only leaf extract. Terpenoids/Steroids present in Liebermann-Burchard's test in all plant parts extract but absent in Salkowski's test in only leaf extract. Anthraquinones were absent in all plant parts extracts using Bontrager's test. Therefore, the presence of these phyto-pharmacological compounds is an indicative that the plant is medicinal and it can be used for the treatment of bacterial and other microbial infections. Further study can be done to separate the individual metabolites to test their antimicrobial activity against some pathogenic bacteria like bacterial meningitis, tuberculosis and syphilis to determine their potency.*

**Keywords:** *Bridelia ferruginea*, ethanolic extracts, medicinal plants, phyto-pharmacological compound, qualitative.

## INTRODUCTION

30 *Bridelia ferruginea* is a common savannah, deciduous tree of genus *Bridelia*. It is usually a  
31 gnarled shrub which sometimes reaches the sizes of tree in suitable condition. *Bridelia*  
32 *ferruginea* Benth (Guinea Fula-Pulaar) in English and their local common names are kizni, kirni  
33 (Hausa), Mirehi (Fulani), Iroladan or “*Epo Ira*” (Yoruba), Ola (Igbo), Awuya (Ebira), (Akuodor  
34 *et al.*, 2012). Its habitat is the savannah, especially in the moister region extending from Guinea  
35 to Zaire and Angola. *Bridelia ferruginea* grows up to 3-4 m high and may be 27.5 cm in width  
36 (Olatunji *et al.*, 2010; Ezike *et al.*, 2011). The stem is often crooked with branches occurring at  
37 the lower regions. The bark is gray, rough and often scaly (Abubakar *et al.*, 2017). The plant of  
38 ten bears spines and may be crimson coloured. The leaves may be small to medium sized,  
39 *Bridelia* species belong to the family Euphorbiaceae and comprise approximately 60-70 species  
40 found in Asia, Africa and Australia ((Abubakar *et al.*, 2017); Olatunji *et al.*, 2010; Kumar and  
41 Pandey, 2013). *Bridelia ferruginea* is utilized in traditional African Medicine in treating disease  
42 conditions such as arthritis, bruises, boils, dislocation, burns, fever, headaches, stiffness,  
43 rheumatic pains and oedema (Olumayokun *et al.*, 2012). Other uses include intestinal disorders,  
44 diabetes, thrush, epilepsy, cough, gonorrhoea, infectious diseases including sexually transmitted  
45 diseases, skin diseases and eruption, skin cancers, roundworm (Chikezie *et al.*, 2015). It is also  
46 an antidote for arrow poison (Kumar and Pandey, 2013), and used as anti-inflammatory (Anulika  
47 *et al.*, 2016) and antitumor agent ((Abubakar *et al.*, 2017)).

48 Reports on the plant have shown that aqueous leaf and root extracts of the plant possesses  
49 hypoglycemic activities. Ethnopharmacological reports have shown that the stem bark extract  
50 possesses antiulcerative properties (Ezike *et al.*, 2011) and anti-inflammatory and antibacterial  
51 (Anulika *et al.*, 2016) properties. Research further indicates that extract of the stem bark  
52 possesses antioxidant properties (Adetutu *et al.*, 2011), antipyretic and analgesic activities  
53 (Akuodor *et al.*, 2011). The aqueous stem bark extract possesses antihypertensive, diuretic and  
54 sedative actions (Nene-Bi *et al.*, 2010; Nene-Bi *et al.*, 2012). The stem bark extracts possess  
55 antioxidative and neuroprotective activities (Omotade, 2012). Furthermore, studies have shown  
56 that the aqueous extract of *B. ferruginea* stem bark reduces vascular permeability in both  
57 cyclophosphamide-induced hemorrhagic cystitis and acetic acid induced vascular permeability in  
58 rats and mice (Anulika *et al.*, 2016). *B. ferruginea* have shown that the stem bark extracts exhibit  
59 anti-inflammatory properties, which were attributed to the suppression of up-regulation of

60 tumour necrosis factor alpha (TNF $\alpha$ ) (Anulika *et al.*, 2016). *B. ferruginea* stem bark extract  
61 inhibits xanthine oxidase and possesses superoxide scavenging activity due to the presence of 3-  
62 O-methylquercetin, myricetin, ferrugin and quercetin 3-O-glucoside (Chikezie *et al.*, 2015). The  
63 stem bark and leaf extracts have contractile effects on the smooth muscle of the bladder (Shakya,  
64 2016). The extracts of *B. ferruginea* possess anti-thrombotic effects Chemical and  
65 pharmacological studies of *Bridelia species* have shown the presence of flavonoids,  
66 sesquiterpenes, triterpenoids, and phenolic compounds (Kumar and Pandey, 2013). *Bridelia*  
67 *species* possess variety of biological activities including antiamebic, antianemic, antibacterial,  
68 anticonvulsant, anti-diabetic, antidiarrhoeal, antihelminthic, anti-inflammatory, antineuro-  
69 inflammatory, antimalarial, antinociceptive, antiviral, and hypoglycemic (Kumar and Pandey,  
70 2013).  
71 Traditional medicine constitutes an important source of drugs for ethnopharmacological  
72 relevance and investigation. Various medicinal food-plants and animal products-supplements are  
73 available for use in certain immune deficiency disease conditions related to malnutrition such as  
74 infectious disease and hemorrhagic sepsis (Kokori *et al.*, 2019)



77  
78 Plate 1: *Bridellia ferruginea* Stem Bark

79



80

81

82

Plate 2: *Bridellia ferruginea* Root



83

84

85

Plate 3: *Bridellia ferruginea* Leaf

86

87

## METHODOLOGY

### 88 **Collection and Identification:**

89 The fresh leave, bark and root samples of *Bridelia ferruginea* were collected at Baba Wali Street,  
90 NTA Community, Behind Kogi State University, Anyigba, Kogi State, Nigeria, April, 2019. The  
91 plant parts were identified and authenticated by U.S Gallah at the Department of Biological  
92 Sciences (Botany option), Kaduna State University. A voucher samples of the plant deposited in  
93 the herbarium unit and the voucher number KASU/BS/1323 was deposited in the herbarium.

94

### 95 **Processing of the plant samples:**

96 The plant parts were washed thoroughly with distilled water, shade dried for 3-4 weeks at room  
97 temperature in the Histology laboratory of the Anatomy Department, Kogi State University,  
98 Anyigba. The plant parts were pulverized in a mortar and pestle and was grounded into fine  
99 powder of 40mm mesh size. The samples were stored in an air-tight container for further use.  
100 The samples was weighed, of which 100g each of the powder were extracted in solvents  
101 (ethanol) 1000 ml macerated and stand for 72 hours. The solvents contained in the maceration  
102 bottle was decanted and filtered using a filter paper, the filtration was aided using a suction  
103 pump. The filtrate was concentrated using a rotary evaporator and then transferred into  
104 thermostatic water cabinet (temperature was set at 45°C), allowed to dry completely. The  
105 extracts obtained was scrapped using a clean spatula and grounded using small laboratory mortar  
106 and pestle, the extracts were weighed using weighing balance.

107 The percentage yield were calculated as follows:

$$108 \quad \% \text{ Yield} = \frac{\text{Weight of extract}}{\text{Weight of plant material}} \times 100$$

109

110 The extract was stored air-tight in a refrigerator prior to use.

111

### 112 **Determination of Phytochemical Compounds in *Bridelia Ferruginea* Parts Extracts**

113 The methods of Chukwuma and Chigozie (2016) and Aziz *et al.*, (2015) were used, the  
114 phytochemical compound was carried out in the Ethanolic Leaf Extract (ELE), Ethanolic Bark

115 Extract (EBE) and Ethanolic Root Extract (ERE) of *B. ferruginea* according to standard  
116 procedures as follows:

### 117 **Test for Carbohydrate**

#### 118 **Molisch's test**

119 **0.5g** of the ELE, EBE AND ERE in a test tube, 3 drops of molisch's reagent was added followed  
120 by concentrated sulfuric acid. The formation of a reddish colored ring at the interface indicates  
121 the presence of carbohydrates (Chukwuma and Chigozie 2016; Aziz *et al.*, 2015).

#### 122 **Test for Saponins**

##### 123 **Frothing test**

124 About 10ml of distilled water was added to **(0.5g)** of the leave, bark and root extract and was  
125 shaken vigorously for 30seconds. The solution was allowed to stand for 5 minutes, the formation  
126 of a persistent froth indicates the presence of saponins (Chukwuma and Chigozie 2016; Aziz *et*  
127 *al.*, 2015).

#### 128 **Test for Flavonoids**

##### 129 **Shinoda test**

130 The extracts **(0.5g)** was dissolved in 2ml of methanol and pieces of metallic magnesium chips  
131 were added followed by few drops of concentrated hydrochloric acid, the formation of a pink,  
132 orange or red to purple coloration indicates the presence of flavonoids (Chukwuma and Chigozie  
133 2016; Aziz *et al.*, 2015).

##### 134 **Sodium hydroxide test**

135 Two drops of 10% Sodium hydroxide was added to the solution of the extracts **(0.5g)**, yellow  
136 coloration indicates the presence of flavonoids (Chukwuma and Chigozie 2016; Aziz *et al.*,  
137 2015).

##### 138 **Ferric chloride test**

139 An amount of 2 to 3 drops of ferric chloride solution were added to the solution of the extracts  
140 **(0.5g)**. Green-blue colour was observed (Chukwuma and Chigozie 2016; Aziz *et al.*, 2015).

141

142

143 **Test for tannins**

144 **Lead sub-acetate test**

145 0.5g of the extracts, 4 drops of lead sub-acetate solution was added, the formation of a cream  
146 coloured precipitate indicates the presence of tannins (Chukwuma and Chigozie 2016; Aziz *et*  
147 *al.*, 2015).

148 **Test for Terpenoids/Steroids**

149 **Salkowski's test**

150 0.5g of the extracts was dissolved in 2ml of chloroform, 3 drops of concentrated sulphuric acid  
151 was added at the side of the test tube. A red brown coloration at the interface indicates the  
152 presence of terpenoids.

153 **Liebermann-Burchard's test**

154 0.5g of the extracts equal volume of acetic anhydride was added and mixed gently. 1ml of  
155 concentrated sulphuric acid was added down the test tube. This was observed for instant colour  
156 changes and over a period of one hour. Blue to blue-green colour in the upper layer and reddish,  
157 pink or purple colour at the junction of the two layers indicates the presence of triterpene  
158 (Chukwuma and Chigozie 2016; Aziz *et al.*, 2015).

159 **Test for Alkaloids**

160 **Dragendoff's test**

161 The extracts (0.5g) was dissolved in 2ml of 5% H<sub>2</sub>SO<sub>4</sub> in 50% ethanol with continuous stirring  
162 in water bath. The mixture was filtered and few drops of Dragendoff's reagent was added, rose  
163 red precipitate indicates the presence of alkaloids (Chukwuma and Chigozie 2016; Aziz *et al.*,  
164 2015).

165 **Mayer's test**

166 To 2ml acidic solution of the extracts (0.5g) in a test tube, few drops of Mayer's reagent were  
167 added, a cream precipitate indicate the presence of alkaloids (Chukwuma and Chigozie 2016;  
168 Aziz *et al.*, 2015).

169

## 170 **Test of Anthraquinones**

### 171 **Bontrager's test**

172 0.5g of the extract was dissolved in 5ml chloroform, shaken and filtered. To the filtrate, an equal  
173 volume of 10% ammonium solution was added with continuous shaking, bright pink colour in  
174 the aqueous upper layer indicates the presence of anthraquinones (Chukwuma and Chigozie  
175 2016; Aziz *et al.*, 2015).

### 176 **Test for Cardiac Glycosides**

#### 177 **Keller-Kiliani test**

178 0.5g of the extracts was dissolved in 1ml glacial acetic acid containing traces of ferric chloride  
179 solution. The solution was then transferred into a dry test tube to which an equal volume of  
180 sulphuric acid was added, a brown ring obtained at the interface will indicate the presence of  
181 deoxy sugar (Chukwuma and Chigozie 2016; Aziz *et al.*, 2015).

### 182 **Test for Fats and Oils**

183 Filter paper soaked in the extracts (0.5g) solution or impregnated with extracts was allowed to  
184 dry and checked for translucence film, which indicates the presence of fats and oils.

185

186

187

188

189

190



191

192

193

194

195 **Table 1. Biologic Activity of Main Groups of Natural Compounds.**

<b>Compound Type</b>	<b>Pharmacological Properties</b>
Terpenoid/steroid	Antimicrobial, antiviral, anthelmintic, antibacterial, anticancer, antimalarial, anti-inflammatory, antineuroinflammatory (Shakya, 2016).
Phenolics acids	Anticarcinogenic and antimutagenic, anti-inflammation and anti-allergic (Anulika <i>et al.</i> , 2016).
Saponins	Antitumor, antiviral, antifungal, anti-inflammatory, immunostimulant, antihypoglycemic, antihepatotoxic and hepatoprotective, anticoagulant, neuroprotective, antioxidant (Negi <i>et al.</i> , 2013).
Flavonoids	Antioxidant activity, cardiovascular protective, anti-inflammatory, hepatoprotective, antiviral, antibacterial (Kumar and Pandey, 2013).
Alkaloids	Antispasmodic, antimalarial, analgesic, diuretic activities, local anesthetic, antihypertensive, antiasthma, antimalarials, diuretic, bactericidal (Chikezie <i>et al.</i> , 2015).
Tannins	Antioxidant, anti-carcinogenic, diuretics, hemostatic, anti-mutagenic, metal ion-chelators, antiseptic, (Saxena <i>et al.</i> , 2013).

196

197

198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220

## RESULT

The results obtained during the course of this experiment/ projects are presented below.

### Percentage Yield of Ethanolic Leave Extract, Ethanolic Bark Extract and Ethanolic Root Extract of *Bridelia ferruginea*

The percentage yield of the extracts (plant parts of *Bridelia ferruginea*) samples is calculated below:

$$\% \text{ yield} = \frac{\text{Weight of extracts}}{\text{Weight of plant}} \times 100\%$$

Weight of ELE= 24.91g, EBE=24.25 and ERE=16.37.

Weight of plant= 100g.

$$\text{Therefore, \% yield} = \frac{24.91\text{g}}{100\text{g}} \times 100\%, \quad \frac{24.25\text{g}}{100\text{g}} \times 100\% \quad \text{and} \quad \frac{16.37\text{g}}{100\text{g}} \times 100\%$$

= 24.91%, 24.25% and 16.37%.

The percentage yield of the extracts obtained is calculated above.

221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240

**Table 2: Percentage Yield of the Plant Parts of *Bridelia ferruginea* Samples**

<b>Samples</b>	<b>Weight of Plant Parts (g)</b>	<b>% Yield Extract (g)</b>	<b>Observed Coloration</b>
Leave extract	100	24.91	Light green
Stem bark extract	100	24.25	Reddish
Root extract	100	16.37	Brownish

**The Results of Phytochemical Compounds.**

The table below is a summary of the phytochemical compounds or secondary metabolites of the Ethanolic Leaf Extract (ELE), Ethanolic Bark Extract (EBE) and Ethanolic Root Extract (ERE) of *B. ferruginea* were tabulated below.

241

242

243

244

245

246 **Table 3: Phytochemical Compounds of ELE, EBE and ERE of *Bridelia ferruginea* Parts**

Phytochemicals	Test	Interference		
		ELE	EBE	ERE
Carbohydrates	Molisch's test	+	+	+
Saponins	Frothing's test	+	+	+
Flavonoids	Shinoda's test	+	+	+
	Sodium Hydroxide's test	+	+	-
	Ferric Chloride's test	+	+	+
Tannins	Lead Sub-acetate's test	+	+	+
Terpenoids/Steroids	Salkowski's test	-	+	+
	Liebermann-Burchard's test	+	+	+
Alkaloids	Dragendoff's test	+	+	+
	Mayer's test	-	+	+
Anthraquinones	Bontrager's test	-	-	-
Cardiac Glycosides	Keller-kiliani's test	+	+	+
Fats and Oils		+	+	+

247 Key: + = Present

248 - = Absent

249

250

251

252

253  
254  
255  
256  
257

## 258 **DISCUSSION**

259 The table 1, shown Biologic activity of main groups of natural compounds. Table 2, percentage  
260 yield of the plant parts of *Bridelia ferruginea* Samples and table 3, shown the Phytochemical  
261 Compounds of ELE, EBE and ERE of *Bridelia ferruginea* which revealed that Carbohydrates,  
262 Saponins, Flavonoids, Tannins, Cardiac Glycosides, fats and oils were present. Alkaloid present  
263 in Dragendoff's test in all plant parts extract but absent in Mayer's test in only leaf extract.  
264 Terpenoids/Steroids present in Liebermann-Burchard's test in all plant parts extract but absent in  
265 Salkowski's test in only leaf extract. Anthraquinones were absent in all plant parts extracts using  
266 Bontrager's test. Temitayo *et al.*, (2017) also carried out the same research and detected the  
267 presence of alkaloids, tannins, flavonoid, cardiac glycosides, saponins and using ethanol. This  
268 result indicates that the parts of the plants have active ingredients responsible for the  
269 antimicrobial activity. The presence of these secondary compounds makes the plants fits or good  
270 for the treatment of bacterial and other microbial infections because most therapeutic effects of  
271 medicinal plants are traced to the plant constituents and the medicinal actions of these plant parts  
272 extract are unique to particular species or family (Yunana *et al.*, 2018). This plant may have high  
273 antimicrobial activity due to the presence of these metabolites. Further study can be done to  
274 separate the individual metabolites to test their antimicrobial activity against some pathogenic  
275 bacteria like bacterial meningitis, tuberculosis and syphilis to determine their potency.

## 276 **Conclusion and Recommendation**

277 The phytochemical composition of the leaf, stem bark and root extracts of the *Bridelia*  
278 *ferruginea* indicate the presence of eight active constituents. The presence of these phyto-  
279 pharmacological compounds is an indicative that the plant has antibacterial property and it can  
280 be used for the treatment of bacterial and other microbial infections. Further investigation,

281 purification and determination of these promising constituents can be done to assay their  
282 antimicrobial activity as alternative medicine.

### 283 **Competing Interests**

284 Authors have declared that no competing interests exist.

285

### 286 **REFERENCES**

287 Abubakar, L., Bagna, E.A. and Dogarai, B.B.S. (2017). Effects of Column Fractions of the  
288 Leaves Extract of *Bridelia ferruginea* on Bacteria. *Bayero Journal of Pure and Applied*  
289 *Sciences*: 10(1): 137 - 141, ISSN 2006 – 6996

290 Adetutu, A., Morgan, W.A. and Corcoran, O. (2011): Antibacterial, antioxidant and fibroblast  
291 growth stimulation activity of crude extract of *Bridelia ferruginea* leaf, a wound-healing  
292 plant of Nigeria. *Journal of Ethnopharmacology* **133**(1): 116-119.

293 Akudor, G.C, Mbah, C.C., Anyalewechi, N.A., Idris-Usman, M., Iwuanyanwu, T.C. and  
294 Osunkwo, U.A. (2011): Pharmacological profile of aqueous extract of *Bridelia ferruginea*  
295 stem bark in the relief of pain and fever. *Journal of Medical and Plants Research*. **5**(22):  
296 5366- 5369.

297 Akuodor GC, Mbah CC, Essien AD, Akpan JL, Ezeokpo BC, Iwuanyanwu TC, Osunkwo UA  
298 (2012). Ulcer-protective and Antidiarrhoeal Effects of the Aqueous Stem Bark Extract of  
299 *Bridelia ferruginea* in Rodents. *Pharmacology* 3: 591–597

300 Anulika NP, Ignatius EO, Raymond ES, Osasere OI, Abiola AH. The chemistry of natural  
301 product: Plant secondary metabolites. *International Journal of Technology Enhancements*  
302 *and Emerging Engineering Research*. 2016; 4(8):1–8. ISSN 2347-4289

303 Aziz M. A., Bashar K., Chowdhury M. M. H. and Faruque A. (2015). Antibacterial, Toxicity and  
304 Larvicidal Potentiality of *Microcos Paniculata* Barks. *Pharmacology Online. Archives*.  
305 *2015. Volume 1. 109-120.*

306 Chikezie PC, Ibegbulem CO, Mbagwu FN. Bioactive principles from medicinal plants. *Research*  
307 *Journal of Phytochemistry*. 2015; 9(3):88–115

308 Chukwuma S. Ezeonu and Chigozie M. Ejikeme (2016). Qualitative and Quantitative  
309 Determination of Phytochemical Content of Indigenous Nigerian Softwoods. *New*  
310 *Journal of Science*. Volume 2016, Article ID 5601327, 9 Pages.

311 Ezike, A. C., Akah, P. A., Nnamani, M. E., Okoli, C. O., Ojike, F. U., Eze, F. S., Chime, C. E.  
312 and Azosiri, I. J. (2011): Studies on the antiulcer and gastrointestinal effects of stem bark  
313 extract of *Bridelia ferruginea*. *Journal of Complementary and Integrated Medicine* **8** (1):  
314 Article 27. DOI. 10.2202/1553-3840.1507.

315 Kokori B. T., Abdullahi A. A. and Abdullahi A. S. (2019). Studies on Phytochemical,  
316 Nutraceutical Profiles and Potential Medicinal Values of *Allium sativum* Linn (Lilliaceae)  
317 on Bacterial Meningitis. *International Neuropsychiatric Disease Journal* 13(2): 1-15;  
318 Article no.INDJ.51609, ISSN: 2321-7235, NLM ID: 101632319

319 Kumar S, Pandey AK. (2013). Chemistry and biological activities of flavonoids: An overview.  
320 *The Scientific World Journal*. 2013: Article ID 162750

321 Negi JS, Negi PS, Pant GJ, Rawat SM, Negi SK (2013). Naturally Occurring Saponins:  
322 Chemistry and Biology. *Journal of Poisonous and Medicinal Plant Research*. 2013;  
323 1(1):001–006

324 Nene-Bi, S. A., Zahoui, O. S., Soro, T. Y. and Traore, F. (2012): Diuretic and natriuretic activity  
325 of an aqueous extract of *Bridelia ferruginea* Benth in normal rats. *J. Physio.Pharmacol.*  
326 **2**(10): 349-353.

327 Nene-Bi, S.A., Zahoui, O.S., Soro, T.Y. and Traore, F. (2010): Evaluation de leffet dun extrait  
328 aqueux de *Bridelia ferruginea* Benth (Euphorbiaceae) sur les activites spontanee et  
329 locomotrice chez le rat. *Intl. J. Biol. Chem. Sci.* **04**(1): 34-41.

330 Olatunji, M. K. and Taofik, O. S. (2010): Effect of Waste Water Treated with Methanolic Bark  
331 Extract of *Bridelia ferruginea* Benth on Rat Kidney and Liver. *Journal of Applied Science*  
332 *and Environmental Sanitation* **7**:14-23.

333 Olumayokun, A.O., Mutalib, A.A., Uchechukwu, P. O. and Bernd, L. F. (2012):  
334 *Bridelia ferruginea* produces antineuroinflammatory activity through inhibition of nuclear  
335 factor- KappaB and P38 MAPK signaling. *Evidence and Complementary Alternative*  
336 *Medicine* **(10)**:1155-1163.

337 Omotade, I. O and Seun, O. B. (2012): *In vitro* Antioxidant Activity of Ethanolic Extract of *B.*  
338 *ferruginea* (stem bark). *Academic Research International* **2**: (2) 246-251.

- 339 Saxena M, Saxena J, Nema R, Singh D. and Gupta A. (2013). Phytochemistry of medicinal  
340 plants. *Journal of Pharmacognosy and Phytochemistry*. 2013; 1(6):168–182
- 341 Shakya A. K. Medicinal plants: Future source of new drugs. *International Journal of Herbal*  
342 *Medicine*. 2016; 4(4):59-64
- 343 Temitayo Omotola Akinsete, Bukola Christianah Adebayo-Tayo and Aimbola Olumide  
344 Adekanmbi (2017). The phytochemical and Antimicrobial Potentials of the Crude Extract  
345 of *Bridelia ferruginea* and the Extracellular Biosynthesized Silver Nanoparticles. *Journal*  
346 *of Advances in Medical and Pharmaceutical Sciences*. 14(3): 1-13.
- 347 Yunana B. T., Guiyi J. C. and Bukar B. B (2018). In vitro and in vivo evaluation of antibacterial  
348 activity of *Bridelia ferruginea* extracts on some clinical isolates. *The Journal of*  
349 *Phytopharmacology*. 2018; 7(4): 392-398 Online at: [www.phytopharmajournal.com](http://www.phytopharmajournal.com)
- 350