

**Nutritional Composition of ‘AjuMbaise’ Herbal Cocktail**

**ABSTRACT**

This study was carried out to evaluate the nutritional content of ‘AjuMbaise’ herbal mixture. The experiment involves the collection of fresh plant samples that make up ‘AjuMbaise’ cocktail. The study was done in the Research Laboratory of the Department of Biochemistry, Faculty of Science, University of Port Harcourt, Nigeria. The plants samples were collected from Mbaise locality in Imo State, Nigeria, and were identified as *Cnestis ferruginea*, *Xylopii aethiopica*, *Uvariachamae*, *Palisotahirsuta*, *Scleria sp.*, *Napoleonaimperialis*, *Dialium guineense*, *Combretum racemosum*, and *Heterotis rotundifolia* respectively. The dietary, mineral, and vitamin compositions were determined accordingly. The proximate, vitamin, and mineral composition of the herbal cocktail were investigated in line with their standard methods of analysis. The proximate analysis revealed that the cocktail extract contains carbohydrate (69.51%), crude protein (10.05%), moisture (8.89%), crude fat (5.17%), fibre (3.745), and ash content (2.65%). The vitamin analysis revealed reasonable concentration of vitamins A, B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>6</sub>, B<sub>12</sub>, C, D, and K. The mineral evaluation showed high concentration of calcium, magnesium, potassium, sodium, zinc, and iron. This study revealed high nutritional value of ‘AjuMbaise’ herbal cocktail which is responsible for the plant’s usefulness in the management, treatment, as well as maintenance of good health.

**Keywords:** AjuMbaise, Minerals, Extraction, Vitamins, and Cocktail.

**1. Introduction**

‘AjuMbaise’ is a herbal therapy composed of combination of leaves, roots and trunk of different medicinal plants wrapped together. This herbal cocktail has its origin from ‘Mbaise’, a sub-tribe of Igbo in Imo State, Nigeria. According to [1], the cocktail effectively detoxifies, and sanitizes the womb, as well as reduces stomach to its original size and shape in good time when administered to women after childbirth. According to [2], the cocktail is made up of *Sphenocentrum jollynum*, *Cnestis ferruginea*, *Xylopii aethiopica*, *Uvariachamae*, *Palisotahirsuta*, *Scleria sp.*, *Napoleona sp.*, *Dialium guineense*, *Combretum racemosum*,

and *Heterotisrotundifolia*. These plants have shown their individual therapeutic effects against so many diseases. According to [3], the root extract of *S. jollynumis* generally used by men as aphrodisiac (Spanish fly). Its capability in stimulating the central nervous system (CNS), management of mental and inflammatory disorders, pain and depression was reported by [4]. Other restorative activities of *S. jollyanum* includes anti-diabetic [5], antioxidant [6], anti-inflammatory [7], anti-allergic [8], anti-bacterial [9], and anti-malaria [6]. According to [5], *S. jollyanum* extract stimulates haematopoietic stem cells. In this manner, improves haematological indices. *C. ferruginea*, according to [10], showed in vivo hypoglycaemic activity in STZ-induced diabetic rodents. Hypoglycemic and hypocholesterolemic activities of *C. ferruginea* leaves have also been reported by [11]. The antimicrobial activities of water and organic extracts of *C. racemosum* leaves have been reported by [12]. Other therapeutic benefits such as, anti-inflammatory, vasorelaxant, and trypanocidal properties of *C. racemosum* extract has been confirmed by [13] and [14]. According to [15], *D. guineense* bark is used for oral hygiene and treatment of stomach ache, fever, prenatal pains and edema. [16], also reported the use of the fruits to improve lactation and check genital infection in women. [17], reported the use of the leaves and stem bark for the treatment of diarrhea, severe cough, bronchitis, wound, stomach ache, malaria fever, jaundice, anti-ulcer and hemorrhoids. According to [18], the fruit pulp contains over 13% of dietary fiber which increases its bulk and augments bowel movement, thereby prevent constipation. [19], reported the utilization of *H. rotundifolia* for treating rheumatism and diarrhea. [20], also reported the antimicrobial effects of *H. rotundifolia* leaves and stems extracts. The leaves extract of *N. imperialis* has shown antibacterial and wound healing properties in rats [21]. [22], demonstrated that a herbal ointment made of methanolic extract of *N. imperialis* showed a better wound healing property. According to [23], the leaves extracts of *P. hirsuta* possess anxiolytic and antidepressant effect. The anti-inflammatory, antipyretic, and anti-oxidant effects of this plant have been reported by [24]. The use of this plant alone or as combination therapy with orthodox medicine in the treatment of various painful inflammatory conditions has also been reported by [25]. According to [26], ethanolic extract of *P. hirsuta* leaves showed to be a potent anti-arthritis therapy in rat adjuvant-induced arthritis when administered prophylactically, curatively or in combination with standard anti-rheumatic drugs. The use of leaf juice of *U. chamae* in the treatment of injuries, swellings, ulcers, iritis and conjunctivitis has been reported by [27]. According to [28], its root bark is a powerful anti-inflammatory, astringent, febrifuge, galactagogue and styptic agent. Its root was also prescribed to be a "lady's prescription" by [28], as it effectively averted amenorrhoea, miscarriage; and quelled labour torments. Also [29], reported that the plant extracts are powerful

solutions for gastroenteritis, intestinal sickness, fever, vomiting, diarrhea, wounds, sore throat and inflamed gums. [30], further expressed the hypoglycemic, antifungal, bacteriostatic and antimalaria properties of *U. chamae*. According to [31], *X. aethiopica* is a potential ethno-medication against rheumatism, arthritis, cough, stomachache, bronchitis, biliousness and dysentery. [32], reported an increased steroid hormone by ethanolic extract of *X. aethiopica*, while its aqueous extract showed diminished amylase and lipase activity [33]. According to [34], *X. aethiopica* can be ingested as remedy for diarrhoea, bronchitis, ulceration, skin disease and female sterility. Other medicinal activities of *X. aethiopica* include antibacterial [35], anti-parasitic [36] and anti-plasmodial [37]. The medicinal abilities of these plants are as a result of the bioactive compounds present in them. Previous study by [2], revealed that the herbal cocktail of 'AjuMbaise' contains various phytochemicals such as alkaloids, flavonoids, glycosides, hydrogen cyanide, phenols, saponins, steroids, tannins, and terpenoids. And so, there is need to further ascertain the nutritive value of the herbal cocktail. Therefore, the present study was designed to investigate the nutritional composition, vitamins and mineral content of 'AjuMbaise' herbal cocktail as dietary supplement for preservation of good health and promoting better life.

## **2. Methodology**

### **2.1 Collection of Plant materials**

Fresh samples of the plants that make up AjuMbaise were collected at Obodo Ujichi, Ahiazu and Amuzi, Ahiaza Towns, both in Ahiazu Mbaise L.G.A, of Imo State, Nigeria. The plants were identified as *Cnestis ferruginea*, *Xylopiia aethiopica*, *Uvariachamae*, *Palisotahirsuta*, *Scleria sp.*, *Napoleonaimperialis*, *Dialium guineense*, *Combretum racemosum*, and *Heterotis rotundifolia*, respectively by Dr. Chimezie Ekeke of the Department of Plant Science and Biotechnology, University of Port Harcourt, Nigeria. The fresh plants after collection were dried. The dried samples were pulverized with a mechanical grinder. The powdered sample was used to determine the proximate analysis, vitamins and mineral content.

### **2.2 Proximate Analysis**

The proximate composition of 'AjuMbaise' herbal cocktail was investigated with the standard methodology according to Association of Analytical Chemist [38].

#### **2.2.1 Estimation of ash content**

Sample weighing 2g was placed in an empty dried platinum crucible of weight ( $W_1$ ). The weight of the sample and the weight of the platinum crucible were noted ( $W_2$ ). The crucible containing the sample was subjected to ashing for 3 hours in a muffle furnace at a temperature of 500°C. The crucible was removed and cooled in a desiccator and the final weight ( $W_3$ ) was determined as follows;

$$\text{Ash content (\%)} = \frac{W_3 - W_1}{W_2 - W_1} \times 100$$

### **2.2.2 Estimation of Carbohydrate content**

Differential method of AOAC [38] was adopted for the estimation of carbohydrate content. The carbohydrate content was calculated thus;

$$\% \text{ Carbohydrate content} = 100 - (\% \text{ Protein} + \% \text{ Moisture} + \% \text{ Ash} + \% \text{ Fat} + \% \text{ Fibre})$$

### **2.2.3 Estimation of Crude fibre content**

A sample of weight 2g was defatted with petroleum ether, boiled under reflux for 30 mins with 200ml of 0.25M sulphuric acid ( $H_2SO_4$ ) solution. It was filtered through a linen fabric and the insoluble matter was washed severally with hot water until it was free of acid. The residue was then transferred to a beaker and boiled for 30 mins with 100ml of hot 0.31M sodium hydroxide (NaOH) solution. This process done repeatedly and the final residue filtered through a thin but close pad of washed and ignited asbestos in a Gooch crucible. The sample was then dried in an electric oven, weighed, incinerated, cooled and reweighed. The crude fibre content was determined as;

$$\% \text{ Crude fibre} = \frac{\text{weight of fibre} \times 100}{\text{weight of sample}}$$

### **2.2.4 Estimation of Crude Protein content**

Sample of weight 0.5g was gently placed in a 30ml Kjeldahl flask. The content of the flask was digested by heating cautiously in the presence of sulphuric acid, anhydrous sodium sulphate and 0.5g of the Kjeldahl catalyst (copper) until a clear solution (ammonia) appears. The clear solution was allowed to stand and cool. After cooling, it was made up to 100ml with distilled water which was added to avoid caking and then 5ml was transferred to the Kjeldahl distillation apparatus, followed by 5ml of 40% NaOH. A receiving flask containing 5ml of 2% boric acid

and mixture of 5 drops of bromocresol blue and 1 drop of methylene blue indicators was subjected to the distillation apparatus such that the tap was about 20cm inside the solution. The distillation was initiated immediately, and was titrated with 0.01M HCl to an end product. The percentage protein content was calculated as;

$$\% \text{ Protein content} = \% \text{ Nitrogen} \times 6.25$$

$$\text{While } \% \text{ Nitrogen} = \text{titre value} \times 0.01 \times 14 \times 4$$

### **2.2.5 Estimation of Crude fat**

A clean dried 500ml round bottom flask containing anti-knocking granules, was loaded up with 300ml petroleum ether of boiling point 40 - 60°C, and stopped neatly with cotton wool. The Soxhlet extraction apparatus was set up and allowed to reflux for about 6 hours. It was gently expelled and the oil ether gathered in the top compartment of the set up was condensed into a flask for re-use. At the point when the compartment was practically free of oil ether, the oil was expelled and dried. It was allowed to cool and afterward measured as;

$$\% \text{ Crude fat} = \frac{(\text{weight of flask} + \text{oil}) - (\text{weight of flask})}{\text{weight of sample}} \times 100$$

### **2.2.6 Estimation of Moisture content**

With a dried clean crucible of weight ( $W_1$ ), a sample weighing 2g was placed in it to make up  $W_2$  (weight of crucible + weight of sample). The crucible containing the sample was placed in an oven for drying at temperature of 105°C for about 24 hours. After the drying process, the weight ( $W_3$ ), of the crucible containing the dried sample was noted. Thus, moisture content was determined as;

$$\text{Moisture content (\%)} = \frac{W_2 - W_3}{W_2 - W_1} \times 100$$

### **2.3 Determination of Vitamin composition**

The amount of vitamin A, C, D, B<sub>6</sub>, and B<sub>12</sub> in the sample was determined according to the method described by [38; 39]. Vitamins B<sub>1</sub> and B<sub>3</sub> were determined according to the method described by [40] while Vitamins B<sub>2</sub> and K were determined using the method described by [41].

### **2.4 Estimation of mineral composition**

The AOAC Official Method 985.01 [42] with little modification described by [43] were used for the analyses of the following metals: Ag, Ar, Al, Co, Mg, Mn, Mo, Hg, Zn, Fe, Cd, Cu, Pb, Ca, Ni, and Se, while the flame photometer was used in the analyses of K and Na. Using Atomic Absorption Spectrophotometer (Buck Scientific model 210 VGP), the ash solution of the plant sample was prepared by weighing 5g of the powdered herbal sample, this was washed at 550°C in muffle furnace for 5 hours, and the residues dissolved in 100ml of deionized water. Suitable salts of the metals were used to make their standards, and suitable lamps were fixed. The standard mineral solutions were injected to calibrate the AAS using acetylene gas as the flame source. An aliquot of ash solutions were injected and the concentrations were extrapolated from the standard plots/curves of each metal.

### 2.5 Statistical Analysis

Data were analysed using SPSS Version 20.0. Results were expressed as Mean  $\pm$  SD of three replicates determinations.

### 3. Results and Discussion

The results of the analysis (proximate, vitamin, and mineral composition) carried out on the herbal cocktail were represented in Tables 1-3 respectively.

**Table 1: Proximate composition of 'AjuMbaise' herbal cocktail**

Parameter	Concentration (g/100g)
Ash	2.65 $\pm$ 0.10
Carbohydrate	69.51 $\pm$ 1.66
Crude Fibre	3.74 $\pm$ 0.12
Lipids	5.17 $\pm$ 0.35
Moisture	8.89 $\pm$ 1.01
Crude Protein	10.05 $\pm$ 1.80

Values represent Mean  $\pm$  Standard deviation, and n = 3.

**Table 2: Vitamins composition of 'AjuMbaise' herbal cocktail**

Parameter	Concentration (mg/100g)
Vitamin A	0.27 $\pm$ 0.02
Vitamin B <sub>1</sub>	0.54 $\pm$ 0.06
Vitamin B <sub>2</sub>	0.34 $\pm$ 0.02
Vitamin B <sub>3</sub>	0.13 $\pm$ 0.03
Vitamin B <sub>6</sub>	0.32 $\pm$ 0.02
Vitamin B <sub>12</sub>	0.23 $\pm$ 0.05
Vitamin C	0.55 $\pm$ 0.06

Vitamin D	0.27±0.04
Vitamin K	0.21±0.11

Values represent Mean ± Standard deviation, and n = 3.

**Table 3: Mineral composition of ‘AjuMbaise’ herbal cocktail**

Mineral	Concentration (mg/kg)
Arsenic	0.10±0.00
Aluminium	0.11±0.01
Cadmium	0.39±0.02
Calcium	3580.00±98.00
Cobalt	0.13±0.01
Copper	0.25±0.01
Iron	27.84±0.96
Lead	0.12±0.00
Magnesium	6750.00±12.00
Manganese	0.24±0.00
Mercury	0.09±0.00
Molybdenum	0.06±0.00
Nickel	0.13±0.00
Potassium	2640.00±8.00
Selenium	2.61±0.06
Silver	0.11±0.02
Sodium	4270.00±13.00
Zinc	28.83±1.07

Values represent Mean ± Standard deviation, and n = 3.

### Discussion

The utilization of natural products (herbs and vegetables) for medicinal purpose is a global practice aimed at achieving good health [44]. These plants are believed to contain many bioactive compounds which include phytochemicals, proximate nutrients, vitamins and minerals. Our research plant ‘AjuMbaise’ is not an exemption, as its constituent plants possess such compounds. From the results presented above (see Table 1), the herbal extract possesses high amount of carbohydrate, protein, moisture and fat, while the fibre and ash content are in little amount. Nutrients such as carbohydrate, proteins, lipids, fibre, minerals and vitamins are known

to be required by the body in suitable proportion to improve and maintain good health. The proximate analysis of the herbal cocktail (AjuMbaise) revealed carbohydrate, crude protein, lipids, crude fibre, moisture, and ash. According to [45], leaves extract of *S. jollyanum*, which is a constituent plant of 'AjuMbaise' is rich in fat, moisture, protein, carbohydrate, ash, and fibre substances respectively, with a great energy value. *Dialium guineense*, another constituent plant of 'AjuMbaise' contains high concentration of carbohydrate, proteins, and moisture [46]. Thus, gave the herbal cocktail its high nutrient content. Minerals and vitamins are essential nutrients, which the body needs its various forms in appropriate proportion to remain healthy. According to [47], they serve as fundamental cofactors and coenzymes for some physiological and metabolic actions and furthermore help essential reactions at cell level (which includes glycolysis, tricarboxylic acid cycle and lipid and amino acid digestion), important for energy generation, and maintenance of life. As indicated by [48], minerals remain an important daily necessity for appropriate tissue functioning. The minerals and vitamins analysis of the 'AjuMbaise' herbal cocktail revealed significant amount of minerals and vitamins such as, vitamin A, vitamin B<sub>1</sub>, vitamin B<sub>2</sub>, vitamin B<sub>3</sub>, vitamin B<sub>6</sub>, vitamin B<sub>12</sub>, vitamin C, vitamin D, vitamin K, arsenic, aluminum, cadmium, calcium, cobalt, copper, iron, lead, magnesium, manganese, mercury, molybdenum, nickel, potassium, selenium, silver, sodium, and zinc (see Tables 2 and 3). This is consonant with the report of [49], that 'AjuMbaise' contained appreciable concentration of minerals (potassium, calcium, magnesium, sodium, iron, zinc, phosphorus, copper, manganese and chromium). [50], revealed the presence of considerable amount of macro- and micro-elements such as calcium, magnesium, potassium, iron, manganese, zinc, and sodium in seeds and leaves of *S. jollyanum* (a component plant of the herbal cocktail), making the plant an important component in building solid bones, production of energy, and carrying out of some metabolic reactions in the body. According to [46], *Dialium guineense*, another constituent plant of 'AjuMbaise' contains appreciable concentration of iron, magnesium, sodium, calcium, potassium, and reasonable amount of ascorbic acid. It can be scientifically expressed that utilization of this herbal cocktail can help fight against mineral and vitamin deficiencies since it contains a wide range of mineral elements and vitamins which are equivalent to those found in different foods necessary to improve health and growth in people. Minerals are crucial for normal growth, muscles activities and skeletal development, cellular activities and oxygen transport (iron and copper), chemical reaction in the body and intestinal absorption (magnesium), fluid balance and nerve transmission (sodium and potassium). As indicated by [51], supplements are imperative to both grown-up and youngsters, as its insufficiency can be adverse to human health. The herbal cocktail extract contained



reasonable amount of vitamin C, a powerful antioxidant that aids the transport and uptake of non-heme iron at the mucosa, and stimulates cortisol synthesis. [52], expressed its adequacy in diminishing sorbitol secondary diabetes and lipid peroxides generated by oxidative stress and free radicals. Its deficiency leads to fragility of blood vessels, gum rot and scurvy. According to [53], pyridoxine (Vit. B<sub>6</sub>) is essential for protein digestion, growth, synthesis of phospholipids and sphingolipids, while cobalamin (Vit. B<sub>12</sub>) is essential for blood formation, maintenance of healthy body, and normal functioning of the brain and nervous system. In another study by [2], it was observed that 'AjuMbaise' herbal cocktail is rich in phytochemicals such as phenols, flavonoids, tannins, alkaloids, saponins, steroids, glycoside, and terpenoids. This is in consonance with the report of [49], that 'AjuMbaise' herbal mixture contains appreciable amount of alkaloids, tannins, flavonoids, cyanogenic glycoside, and saponin.

### **Conclusion**

This study has shown that 'AjuMbaise' herbal cocktail contains numerous essential nutrients that play very important biological roles in the human system that aids in the maintenance of good health and promotion of quality healthy living.

### **Acknowledgement**

The authors are grateful to Messrs Ekeocha Akujobi, Cajethan Ugorji and Franklin Nwuda, for their assistance in the sample collection that led to the success of this work.

### **Competing Interests**

Authors have declared that no competing interests exist.

### **Authors' Contributions**

**Nnadiukwu, T. A., and Monago-Ighorodje, C. C.**, designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript.

**Nnadiukwu, T. A.**, managed the analyses of the study.

**Nnadiukwu, T. A., and Chuku, L. C.**, managed the literature searches.

All authors read and approved the final manuscript.

### **Ethical Approval**

The Research work was approved by the Research Ethics Committee of the University of Port Harcourt with authorization number: UPH/CEREMAD/ REC/MM64/003.

## References

1. Ogueke CC, Owuamanam CI, Onyedinma C, Iroanya A, Bede EN, Nwachukwu IN. Antibacterial activity, phytochemical properties and mineral Content of “AjuMbaise” decoction: A liquid extract administered to nursing mothers. *Niger. J. Nutr. Sci.* 2016;37(1):114-21.
2. Nnadiukwu AT, Monago-Ighorodje CC, Chuku LC. Phytochemical Composition of Ethanol Extract of a Cocktail Herbal Mixture (AjuMbaise). *J. Adv. Biol. Biotechnol.* 2019 Jul 16:1-1.
3. Soelberg J, Asase A, Akwetey G, Jäger AK. Historical versus contemporary medicinal plant uses in Ghana. *J ETHNOPHARMACOL.* 2015;160:109-32.
4. Oke JM, Hamburger MO. Screening of some Nigerian medicinal plants for antioxidant activity using 2, 2, diphenyl-picryl-hydrazyl radical. *Afr J Biomed Res.* 2002;5(1-2).
5. Mbaka GO, Adeyemi OO, Adesina SA. Anti-diabetic activity of the seed extract of *Sphenocentrumjollyanum* and morphological changes on pancreatic beta cells in alloxan-induced diabetic rabbits. *J. Med. Med. Sci.* 2010;1:550-6.
6. Olorunnisola OS, Afolayan AJ. In vivo antioxidant and biochemical evaluation of *Sphenocentrumjollyanum* leaf extract in *P. berghei* infected mice. *Pak. J. Pharm. Sci.* 2013;26(3):445-50.
7. Alese MO, Adewole OS, Ijomone OM, Ajayi SA, Alese OO. Hypoglycemic and hypolipidemic activities of methanolic extract of *Sphenocentrumjollyanum* on streptozotocin-induced diabetic wistar rats. *Eur. J. Med. Chem.* 2014; 12:353-64.
8. Olorunnisola OS, Adetutu A, Popoola RB, Owoade AO, Adegbola P, Adesina BT. Nephroprotective effect of ethanolic leaf extract of *Thaumatococcusdanielli* (benth.) in streptozotocin induced diabetic Rats. *Funct. Food Health Dis.* 2017;7(12):923-35.
9. Koleosho AT, Jose AR, Oyibo PG, Roland-Ayodele MA, Uloko ME. Antimicrobial Activity of *Sphenocentrumjollyanum* and *MangiferaindicaLinn* on *Salmonella Typhi*. *IOSR J. Pharm.* 2013;5(3):50-4.
10. Rahmat AA, Dar FA, Choudhary IM. Protection of CCl4-induced liver and kidney damage by phenolic compounds in leaf extracts of *Cnestisferruginea* (de Candolle). *Pharmacogn. Res.* 2014;6(1):19.
11. Adisa R, Choudhary M, Adewoye E, Olorunsogo O. Hypoglycaemic and biochemical properties of *Cnestisferruginea*. *AFR J TRADIT COMPLEM.* 2010;7(3).
12. Kola KA, Benjamin AE. Comparative antimicrobial activities of the leaves of *Combretummicranthum* and *C. racemosum*. *Glob. J. Med. Sci.* 2002;1(1):13-7.
13. Eze JI, Anosa GN, Ozota CA. In vitro and in vivo trypanocidal activity of *Combretumracemosum* leaves. *Afr. J. Biotechnol.* 2012;11(46):10611-6.
14. Manga FN, El Khattabi C, Fontaine J, Berkenboom G, Duez P, Nzunzu JL, Pochet S. Vascular effects and antioxidant activity of two Combretum species from Democratic Republic of Congo. *J ETHNOPHARMACOL.* 2012;142(1):194-200.
15. Idu M, Umweni AA, Odaro T, Ojelede L. Ethnobotanical Plants Used for Oral Healthcare Among the Esan Tribe of Edo State, Nigeria. *Ethnobot Leaflets.* 2009; (4):15.
16. Nwosu MO. Plant resources used by traditional women as herbal medicine and cosmetics in Southwest Nigeria. *Arzte fur NaturFahr.* 2000;41(11).

17. Bero J, Ganfon H, Jonville MC, Frédéric M, Gbaguidi F, DeMol P, Moudachirou M, Quetin-Leclercq J. In vitro antiplasmodial activity of plants used in Benin in traditional medicine to treat malaria. J ETHNOPHARMACOL. 2009;122(3):439-44.
18. Besong EE, Balogun ME, Djobissie SF, Obu DC, Obimma JN. Medicinal and economic value of *Dialiumguineense*. Afr J Biomed Res. 2016;19(3):163-70.
19. Dougnon VT, Klotoé JR, Sènou M, Roko GO, Dougnon G, Fabiyi K, Amadou A, Aniambossou A, Assogba P, Bankolé H, Dougnon J. Chemical composition, cytotoxicity and antibacterial activity of selected extracts of *Euphorbia hirta*, *Citrus aurantifolia* and *Heterotisrotundifolia* on enteropathogenic bacteria. EC Microbiol. 2017; 12(4): 180-195.
20. Abere TA, Okoto PE, Agoreyo FO. Antidiarrhoea and toxicological evaluation of the leaf extract of *Dissotisrotundifoliatriana* (Melastomataceae). BMC Complement Altern. Med. 2010;10(1):71.
21. Chah KF, Eze CA, Emuelosi CE, Esimone CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. J ETHNOPHARMACOL. 2006;104(1-2):164-7.
22. Esimone CO, Ibezim EC, Chah KF. The wound healing effect of herbal ointments formulated with *Napoleonaimperialis*. J med Pharm Allied Sci. 2006;3(1):294-9.
23. Woode E, Boakye—Gyasi E, Danquah CA, Anisah C, Duwiewua M. in Freund's Adjuvant-Induced Arthritis in Rats. Int. J. Pharmacol. 2009;5(3):181-90.
24. Boakye-Gyasi E, Ainooson GK, Abotsi WK. Anti-inflammatory, antipyretic and antioxidant properties of a hydroalcoholic leaf extract of *Palisotahirsuta* K. Schum.(Commelinaceae). West African J Pharm. 2011;22(1).
25. Bello OM, Fasinu PS, Ogbesejana AB, Bunmi AJ, Abdulrahman B, Osibemhe M, Stephen OO. Anti-inflammatory activity of extracts of selected West African ethnomedicinal plants. Eurasia J Biosci. 2019;13:1-9.
26. Woode E, Boakye-Gyasi E, Amidu N, Anisah C, Duwiewua M. Anxiolytic and antidepressant effects of a leaf extract of *Palisotahirsuta* K. Schum.(Comelinaceae) in mice. Int. J. Pharmacol. 2010;6(1):1-7.
27. Omajali JB, Hussaini JS, Omale J. Cytotoxicity and anti-inflammatory studies on *Uvariachamae*. J. Pharmacol. Toxicol. 2011;2(7):1-9.
28. Achigan-Dako EG, Pasquini MW, AssogbaKomlan F, N'danikou S, Yédomonhan H, Dansi A, Ambrose-Oji B. Traditional vegetables in Benin. Institut National des RecherchesAgricoles du Bénin, Imprimeries du CENAP, Cotonou. 2010.
29. OKWU DE. The potentials of *Ocimumgratissimum*, *Penrgulariaextensa* and *Tetrapleuratetraptera* as spice and flavouring agents. Niger. Agric. J. 2003;34:143-8.
30. Okwuosa OM, Chukwura EI, Chukwuma GO, Okwuosa CN, Enweani IB, Agbakoba NR, Chukwuma CM, Manafa PO, Umedum CU. Phytochemical and antifungal activities of *Uvariachamae* leaves and roots, *Spondiasmombin* leaves and bark and *Combretumracemosum*leaves. Afr. J. Med. Med. Sci. 2012;41:99-103.
31. Fleischer TC, Mensah ML, Mensah AY, Komlaga G, Gbedema SY, Skaltsa H. Antimicrobial activity of essential oils of *Xylopiiathiopica*. AFR J TRADIT COMPLEM. 2008;5(4):391-3.
32. Woode E, Alhassan A, Abaidoo CS. Effect of ethanolic fruit extract of *Xylopiiathiopica* on reproductive function of male rats. Int J Pharm Biomed Res. 2011;2(3):161-5.
33. Etoundi CB, Kuaté D, Ngondi JL, Oben J. Anti-amylase, anti-lipase and antioxidant effects of aqueous extracts of some Cameroonian spices. J. Nat. Prod. 2010;3(2010):165-71.
34. Mshana NR. Abbiw. K, Addae-Mensah, I., Adjanohoun, E., Ahyi, MRA, Ekpere, JA, Enow-Orock, EG, Gbile, Z. O., Noamesi, GK, Odei, MA, Odunlami, H., Oteng-Yeboah, AA, Sarpong, K., Soforowa, A. and Tackie, AN. Traditional medicine and pharmacopoeia,

- contribution to the revision of ethnobotanical and floristic studies in Ghana. *OAU/STRC Technical Report*. 2000:676-7.
35. Kuete V. Potential of Cameroonian plants and derived products against microbial infections: a review. *Planta Med*. 2010;76(14):1479-91.
  36. Tatsadjieu LN, Ngang JE, Ngassoum MB, Etoa FX. Antibacterial and antifungal activity of *Xylopiiaaethiopica*, *Monodoramyristica*, *Zanthoxylumxanthoxyloides* and *Zanthoxylumleprieuriifrom* Cameroon. *Fitoterapia*. 2003;74(5):469-72.
  37. Boyom FF, Ngouana V, Zollo PH, Menut C, Bessiere JM, Gut J, Rosenthal PJ. Composition and anti-plasmodial activities of essential oils from some Cameroonian medicinal plants. *Phytochemistry*. 2003;64(7):1269-75.
  38. Association of Official Agricultural Chemists, Horwitz W. *Official methods of analysis*. Washington, DC: Association of Official Analytical Chemists; 1975.
  39. Achikanu CE, Eze-Stephen PE, Ude CM, Ugwuokolie OC. Determination of the vitamin and mineral composition of common leafy vegetables in South Eastern Nigeria. *Int J CurrMicrobiolAppl Sci*. 2013;2(11):347-53.
  40. Okwu DE. Phytochemicals, vitamins and mineral contents of two Nigerian medicinal plants. *Int. J. Mol. Med. Adv. Sci*. 2005;1(4):375-81.
  41. Okwu DE, Josiah C. Evaluation of the chemical composition of two Nigerian medicinal plants. *Afr. J. Biotechnol*.2006;5(4):357-61.
  42. AOAC Int. *Official methods of analysis*. Assoc. Offic. Anal. Chem. 2006.
  43. Chinedu KE, Onyeike EN, Ikewuchi CC. Nutrient Composition of *Brillantaisiaguianensis P. Beuv* Leaves. *Int J Biochem Res Rev*. 2019; 27(3): 1-12.
  44. Ali-Shtayeh MS, Jamous RM, Jamous RM. Plants used during pregnancy, childbirth, postpartum and infant healthcare in Palestine. *Complement. Ther. Clin. Pract*. 2015;21(2):84-93.
  45. Ajayi IA, Ojelere OO, Ghaidaa M, Yanchang W, Abdallah H, Chang H, Zhao JJ, Ye SD, Ekise IE, Nahayo A. Phytochemical analysis and mineral element composition of ten medicinal plant seeds from South-west Nigeria. *N Y Sci J*. 2013;6(9):1-7.
  46. Ayessou NC, Ndiaye C, Cisse M, Gueye M, Sakho M, Dornier M. Nutrient composition and nutritional potential of wild fruit *Dialiumguineense*. *J Food Compost Anal*. 2014;34(2):186-91.
  47. Balogun IO, Olatidoye OP. Chemical composition and nutritional evaluation of velvet bean seeds (*Mucunautilis*) for domestic consumption and industrial utilization in Nigeria. *Pak J Nutr*. 2012;11(2):116.
  48. Iniaghe OM, Malomo SO, Adebayo JO. Proximate composition and phytochemical constituents of leaves of some Acalypha species. *Pak J Nutr*. 2009;8(3):256-8.
  49. Ezejindu CN, Iro OK. Antibacterial activity, phytochemical properties and mineral content of “AjuMbaise” decoction administered to nursing mothers. *Direct Res. J. Health Pharm*. 2017; 5(3): 33-38.
  50. Ibiro AA, Olusola OO. Phytochemical analysis and mineral element composition of ten medicinal plant seeds from South-west Nigeria. *N Y Sci J*. 2013; 6:2–28.
  51. Beers MH. *The Merck manual of health & aging*. Random House Digital, Inc.; 2005.
  52. Riccioni G, Bucciarelli T, Mancin IB, Corradi F, Dillio C, Mattei PA, D’Orazio N. Antioxidant vitamin supplementation in cardiovascular diseases. *Ann. Clin. Lab. Sci*. 2007; 37: 89-95.
  53. Shils ME, Shike M, editors. *Modern nutrition in health and disease*. Lippincott Williams & Wilkins; 2006.