

## **Evaluation of some Valencia cvs. performance under new reclaimed soil conditions**

---

### **ABSTRACT**

**Aims:** The objective of this study was to evaluate the performance of three imported Valencia cultivars (*Citrus sinensis* (L.) Osb.) namely 'Campbell', 'Olinda' and 'Delta' all were budded on Volkamer lemon (*Citrus volkameriana*) and grown in sandy soil under drip irrigation system.

**Study design:** One way completely randomized design was used for the experiment.

**Place and Duration of Study:** The experimental was carried out during two successive seasons 2014- 2015 and 2015- 2016 at a private citrus orchard in South El Tahrir, El Beheira governorate, Egypt.

**Methodology:** The study covers some morphological, anatomical and physiological characteristics for the three investigated cultivars. Yield, fruit quality, total indoles, total phenols, total carbohydrates and minerals concentrations were determined.

**Results:** Anatomical study showed that the 'Campbell' leaves gave the higher values of most tissues measurements under study i.e., palisade thickness, mid vein thickness and main vascular bundle length and width comparing to 'Olinda' and 'Delta' cvs.. Moreover, fruitlets structure demonstrated that, the highest values of flavedo thickness were observed in 'Delta' and the highest value of segments width was obtained by 'Olinda'. Likewise, histological characters of the leaf of three cultivars and their fruitlets structure seemed to be reflected on their growth, yielding and fruit quality. Generally, the results revealed that, 'Campbell' produces the highest yield and best fruit quality parameters, whereas 'Olinda' fruits gave the highest fruit juice percentage which is an extremely important parameter for its industrial processing.

**Conclusion:** Thus, it could be concluded that 'Campbell' proved as reliable high yielding cultivar with good fruit characteristics followed by 'Olinda' with advantage for juice processing under the prevailing agro-climatic conditions of South El Tahrir district, Egypt.

*Keywords: Evaluation, Campbell, Olinda, Delta, Anatomical characteristics, Fruit quality*

### **1. INTRODUCTION**

Citrus is one of the leading fruit crops under tropical and sub-tropical conditions of the world with respect to its area and production. Among different citrus species, sweet orange (*Citrus sinensis* Osbeck) is one of the prominent groups with wide range of varieties and distribution. More than 60 percent global citrus production is contributed by the sweet orange [1].

The orange tree is small, spiny tree, typically growing to 7.5m, but occasionally reaching heights up to 15m. Leaves are leathery and evergreen, and range from elliptical to oblong to oval, 6.5-15cm length and 2.5-9.5cm wide, often with narrow wings on the petioles. The

fragrant white flowers, produced singly or in cluster of up to 6 are around 5cm wide, 5 petals and 20 to 25 yellow stamens. The fruit, which may be globose to oval, is typically 6.5 to 9.5 cm wide, and ripens yellow to orange. The fruit rind contains numerous small oil glands. The fruit pulp is typically juicy and sweet, divided into 10 to 14 segments (while, there are seedless varieties) and ranges in color from yellow to orange to red [2].

Valencia orange is considered as one of the best and most popular late-maturing citrus varieties, prized for its high productivity and good juice quality. For these reasons, Valencia orange is the most widely cultivated citrus variety in the world. The most well-known clonal selections of Valencia orange are 'Cutter', 'Delta', 'Frost', 'Lue Gim Gong', 'Olinda', and the vigorous clone 'Campbell' [3]. Valencia oranges are known for their high- quality juice, which has a deep orange color and high sugar content. However, the fruit is medium in size with few seeds (0-6) [4].

The different ecological conditions effects on citrus productions are apparent. Thus, it is valuable to know the favorable ecological conditions for the cultivars chosen and their interactions under these parameters by ecological conditions of the growing sites. Further, factors like cultivar characteristics, rootstocks, growing conditions along with cultural practices, type of flowers, and the fruit drops can affect yield and quality performance of citrus cultivars [5]. Since environmental conditions and cultural practices are unique and vary considerably from one area to another, thus this study was carried out to determine the horticultural adaptability and performance of 'Campbell', 'Olinda' and 'Delta' cultivars under conditions of South El Tahrir, El Beheira governorate, Egypt.

## 2. MATERIAL AND METHODS

The present study was carried out during two successive experimental seasons 2014 - 2015 and 2015 - 2016 in a private citrus orchard in El Beheira governorate, Egypt. Three Valencia orange cultivars (*Citrus sinensis* (L.) Osb.) namely 'Campbell', 'Olinda' and 'Delta' budded on Volkamer lemon (*Citrus volkameriana*), trees were eleven years old grown on sandy soil at 4 × 6 m under drip irrigation system, were used. The total number of trees in this experiment was forty five trees (3 cultivars x 5 replicate x 3 trees in each replicate).

Following parameters were investigated:

### 2.1 Tree canopy

Tree canopy volume was determinate at the end of February during two experimental seasons; tree canopy volume was estimated according to the formula of Obreza [6].

Tree canopy volume (m) = H × D × 0.5238

Whereas H= tree height (m) and D = diameter of tree periphery (m).

### 2.2 Anatomical studies

Leaves at the first week of March and fruitlets at the first week of May were collected from the three studied cultivars ('Campbell', 'Olinda' and 'Delta' trees) throughout the 2<sup>nd</sup> growing season of 2015/2016. Specimens were killed and fixed for 48 hours in F.A.A. (10 ml formalin, 5 ml glacial acetic acid, 50 ml ethyl alcohol 95%, and 35ml distilled water). Plant materials were washed in 50% ethyl alcohol and dehydrated in a normal butyl alcohol series before being embedded in paraffin wax (melting point 56 °C). Transverse sections, 20 µm thick, were cut using a rotary microtome, double stained with crystal violet/erythrosine, cleared in xylene and mounted in Canada balsam [7]. Examination and photomicrographs were taken at Botany Department, Faculty of Agriculture, Cairo University.

### **2.3 Fruit set**

Total number of flowers was counted at 75% of blooming and number of fruitlets was counted at the second week of June and then fruit set percentage (%) was calculated according to the equation:

$$\text{Fruit set\%} = (\text{number of fruitlets} / \text{number of flowers}) \times 100$$

### **2.4 Yield**

At harvest time (at the third week of March under these experimental conditions) fruits of each tree were harvested and the yield was estimated as number of fruits and weight in Kg.

### **2.5 Fruit quality**

At harvest stage, representative sample of 10 fruits was taken from each tree and the following characters were determined:

#### **2.5.1 Fruit physical properties**

Average fruit weight (g), average fruit size (cm<sup>3</sup>), fruit height and diameter (cm) were measured and fruit shape index (length/diameter) was calculated, peel thickness (cm), fruit firmness (l.b/ inches<sup>2</sup>) and fruit juice percent % (w/w) were measured.

#### **2.5.2 Fruit chemical properties**

TSS %, acidity % (as mg citric acid/100 cm<sup>3</sup> juice), TSS/ acid ratio and vitamin C (ascorbic acid as mg/100 ml juice) were determined according to A.O.A.C.[8].

### **2.6 Leaf chemical composition**

Both total indoles and total phenols were determined in fresh leaves three times (March, May and July) at the two experimental seasons. Total indoles were determined according to Larsen [9]. Total phenols were determined according to Swain and Hillis [10]. Total carbohydrates in dry shoots of spring cycle were determined in September at the two experimental seasons by using 3,5-dinitrosalicylic acid method according to Miller [11]. N, P, K, Fe, Mn and Zn concentrations in dried leaves were determined in September of the two experimental seasons. Total N% was determined by semi-micro Kjeldahl method described by Plummer [12]. Phosphorus was estimated colorimetrically by using the chlorostannous reduced molybdophosphoric blue colour method as described by King [13]. Potassium concentration was determined by using the flame photometer. Fe, Mn and Zn concentrations were determined by using atomic absorption spectrophotometer.

### **2.7 Statistical analysis**

One way completely randomized design was used for the experiment. The data statistical analysis carried out according to Snedecor and Cochran [14]. The multiple comparisons of means were performed according to Duncan's multiple test range [15] using COSTAT computer program.

## **3. RESULTS AND DISCUSSION**

### **3.1 Anatomical studies**


#### **3.1.1 Leaf studies**

Leaves are simple, leathery texture, deep green colour, ranging from elliptical to oblong to oval in shape, 5.6 cm long and 2.3 cm wide, the apex is acuminate, it have entire margin, bearing narrow wings on the petioles. Anatomically, data showed that the mesophyll in cross section of the leaf of the three cultivars is heterogeneous, consist of 2 rows of palisade and 8-9 rows of spongy tissues. The upper and lower epidermis is represented by one row of cell. Lamina thickness in 'Campbell' is higher in values (578 µm), while 'Olinda' (570 µm) and 'Delta' (510 µm). This is because the thickness of palisade and spongy tissues were increased in 'Campbell' recorded 170 and 400 µm, respectively. Whereas, in 'Olinda' recorded 165 and 365 µm and in 'Delta' are 150 and 300 µm, respectively. The mid vein

thickness in 'Campbell' is higher (1380  $\mu\text{m}$ ) followed by 'Olinda' (1115  $\mu\text{m}$ ), but in 'Delta' recorded 940  $\mu\text{m}$  as a result of increasing length and width of midvein bundle recorded 690 and 1060  $\mu\text{m}$  in 'Campbell', and 480 and 890  $\mu\text{m}$  in 'Olinda', whereas in 'Delta' recorded 445 and 830  $\mu\text{m}$ , respectively. These results are in harmony with Sedeeq *et al.* [16] on *Citrus maxima*. Microscopically measurements and microphotographs of histological characters at the leaf of three cultivars of *Citrus sinensis*, 'Campbell', 'Olinda' and 'Delta' are given in Table (1) and Figure (1).

**Table (1): Anatomical characters of citrus cultivars leaf during 2<sup>nd</sup> season**

Characters ( $\mu\text{m}$ )	Cultivars		
	Campbell	Olinda	Delta
Lamina thickness	578	570	510
Palisade thickness	170	165	150
Spongy thickness	400	365	300
Mid vein thickness	1380	1115	940
Main vascular bundle			
- Length	690	480	445
- Width	1060	890	830







Fig. (1): Transverse sections through the middle part of citrus leaf, (A) 'Campbell' (B) 'Olinda' (C) 'Delta' (X 40): Up. Epi: upper epidermis, Pal: palisade, Spo: spongy, Xyl: xylem, Phl: phloem, Lo. Epi: lower epidermis

### 3.1.2 Fruitlets structure

A transverse section of the three studied cultivars of *Citrus sinensis* fruitlets were taken and illustrated in Table (2) and illustrated in Fig. (2). The fruitlets, globose to oval in shape, it is 7.0 to 9.5 cm wide, and ripens to orange. Concerning anatomical structure of fruitlets, results showed that it is composed of an outer flavedo layer that contains the exterior fruitlets color and oil glands (epicarp). Flavedo made of parenchymatous cells covered with cuticle and embedded oil glands. The thickness of flavedo recorded the highest values in 'Delta' (290  $\mu\text{m}$ ) while in 'Campbell' showed 262.5  $\mu\text{m}$  and meanwhile 'Olinda' recorded the lowest (230  $\mu\text{m}$ ). Under the flavedo is a white spongy albedo layer (mesocarp). Albedo formed of polygonal parenchyma cells, showing air spaces and vascular bundles. The thickness of albedo ranged from 1150  $\mu\text{m}$  in 'Delta' to 1635  $\mu\text{m}$  in 'Olinda'. The endocarp is a membranous parenchymatous cells. It is separated into 10 to 14 segments; filled with juice vesicles that are elongated and attached to the center of the fruit. Segments width showed an increase value in 'Olinda' (477.5  $\mu\text{m}$ ) followed by 'Campbell' (420  $\mu\text{m}$ ) then 'Delta' recorded 400  $\mu\text{m}$ . The wall thickness between segments showed the most increase value in 'Campbell' (105  $\mu\text{m}$ ), while recorded 85 and 90  $\mu\text{m}$  in 'Delta' and 'Olinda', respectively. Similar results were obtained by Sedeek *et al.* [16] on *Citrus maxima* (Burm.) Merrill.

Table (2): Anatomical characters of citrus cultivars fruitlets during 2<sup>nd</sup> season

Characters ( $\mu\text{m}$ )	Cultivars		
	Campbell	Olinda	Delta
Flavedo thickness	262.5	230.0	290.0
Albedo thickness	1340.0	1635.0	1150.0
Segments width	420.0	477.5	400.0
Wall thickness	105.0	90.0	85.0



**Fig. 2: Transverse sections through the fruitlets of different citrus cultivars (A), 'Campbell' (B) 'Olinda' (C) 'Delta' (X 40)**

### **3.2 Tree canopy**

The data in Table (3) showed that, in the two experimental seasons the highest value of tree canopy was observed in 'Campbell' Valencia trees as compared with the other cultivars, which, may affected by the increment of palisade 'Campbell' leaves comparing to the other cultivars (Table1). Since, the leaf vein features responsible for water, nutrient, and sugar transport, and biomechanical support; thicker veins may have greater water and sugar transport capacity [17].

**Table (3): Tree canopy, fruit set, fruit number and yield of studied cultivars**

Cultivars	Tree canopy (m)		Fruit set (%)		Fruit number		Yield (Kg/ tree)	
	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
Campbell	12.58 a	14.10 a	25.43 a	28.92 a	263.33 a	254.33 a	69.17 a	64.49 a
Olinda	10.37 b	11.58 b	18.04 b	16.86 b	205.00 b	216.67 b	61.57 a	62.03 a
Delta	9.55 b	9.61 c	14.81 b	14.17 b	180.00 b	205.00 b	42.50 b	51.18 b

Means in each column followed by the same letter did not differ at  $p < 0.05$  according to Duncans multiple range tests

It clear from the results that, there are positive relationship between tree canopy and fruit yield. Whereas, 'Campbell' trees had the highest tree canopy and fruit yield as compared with other cultivars.

These results are in agreement with those were obtained with Hostler, *et al.* [18] who found that, there was a positive correlation between tree canopy and fruit yield of citrus as well as Zaman *et al.*[19] who noticed that, tree age, size and yield maps produced similar spatial patterns with the grove, as high yielding areas were associated with large tree canopies.

### **3.3 Fruit set**

The data in Table (3) revealed that, in the two experimental seasons the highest value of fruit set was recorded by 'Campbell' trees followed by 'Olinda'.

Flowering induction and flower number are main factors for yield and fruit setting production in citrus crops [20]. Fruit set rather than flowering is the step that limits yield in most *Citrus* cultivars [21].

In this respect, there are correlations between accumulation of carbohydrates and flower formation, but carbohydrate levels are not the sole factor regulating citrus flowering [22]. Although the evidence is still mostly indirect, it may be concluded that the level of carbohydrates is often a major factor limiting fruit set [23]. Developing fruits serve as competitive sinks for available metabolites [24]. Also, during the period of fruit abscission, in which competition for carbohydrates is considered to be a limiting factor for fruit retention, fruit nutrition is supported by current photosynthesis and stored reserves [25].

### **3.4 Fruit yield**

It is clear from the data presented in Tables (3) that, the 'Campbell' trees significant higher fruit yield comparing to 'Olinda' trees at the two experimental seasons, followed by 'Delta' trees. This lake of significant in fruit yield between the two cultivars results from the superiority of 'Campbell' in producing more fruits while; 'Olinda' had more weighty fruits.

This results are in harmony with those were obtained Qureshi *et al.* [26] who found that, maximum fruit number and fruit yield were recorded by 'Campbell' Valencia followed by 'Hinkely' and 'Olinda' Valencia orange.

In this respect, 'Campbell' Valencia was found to be heavy yield as compare with 'Olinda' Valencia, with higher juice volume but had high acidity level [26]. The highest yield was

achieved when the 'Olinda Valencia' orange was grafted on Macrophylla and 'Volkamer' lemon rootstocks while those grafted on Cleopatra mandarin produced the lowest yield [27].

### **3.5 Fruit quality**

#### *3.5.1 Physical Properties*

The data in Table (4) showed that, in the two experimental seasons, the highest values of fruit weight and fruit size were recorded in 'Olinda' trees as compared with the other cultivars.

These results are in agreement with those obtained with Singh and Gill [28] who found that, 'Olinda' recorded maximum values for fruit weight, peel weight and percent juice content.

**Table (4): Fruit physical properties of studied cultivars**

Cultivars	Fruit weight (g)		Fruit size (cm <sup>3</sup> )		Fruit shape index		Fruit firmness (l.b/inches <sup>2</sup> )		Peel thickness (cm)	
	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
Campbell	262.67 b	253.58b	306.67 b	289.67a	1.08 a	1.07 a	15.51 b	15.50 b	0.59 a	0.56 a
Olinda	300.33 a	286.27a	347.67 a	305.00 a	1.04 a	1.06 a	17.45 b	16.00 b	0.58 a	0.53 a
Delta	235.33 c	249.67b	269.33c	270.00 b	1.10 a	1.09 a	18.05 a	17.20 a	0.58 a	0.57 a

Means in each column followed by the same letter did not differ at  $p < 0.05$  according to Duncans multiple range tests

Fruit size is the main factor affecting the market price of Valencia fruit [3]. In general, fruit size is correlated with fruit number per tree. The fewer fruit on the tree, the larger and heavier are the fruit. Moreover, in a particular year beside fruit load, the ultimate size a citrus fruit achieves is the result of many complex factors including nutrition and irrigation programs, pruning, and the rootstock-scion combination. Large fruit size is most often preferred in the fresh fruit market and brings higher prices early in the season [29].

In this respect, citrus fruit quality may be indicated by external fruit features, such as peel colour, size, rind texture, and physical as well as biochemical characters of its internal features, like seediness, juice and vitamin C contents, total soluble solids, titratable acidity and TSS\acid ratio [30].

The data in Table (4) demonstrated that, there was no significant difference in peel thickness of the studied cultivars. It is obvious from data in Table (4) that, in the two experimental seasons the highest values of fruit firmness was observed in 'Delta' cv. fruit. This may explained as the thickness of 'Delta' flavedo recorded the highest values in as showed previously in its anatomical parameters (Table 2). As Sirisomboon and Lapchareonsuk [31] cleared that, the average diameter was found to correlate with the initial firmness and toughness of the flavedo. While, the highest value of fruit juice percent was recorded by 'Olinda' cv. fruit. This may refer to the augmentation of the 'Olinda' fruit segments width (Table 2). As, segment length was positively correlated with fruit juice weight and fruit juice percentage [32].

These results are in agreement in with those obtained by Singh and Gill [28] who found that, highest juice percentage was recorded in fruits harvested from 'Olinda' cultivar.



In this respect, the juice percentage in the fresh citrus fruit is considered to be very important factor due to the increasing demand in fruit juice consumption [33]. Highest juice percentage in citrus fruits is an ultimate customer's demand [34].

### 3.5. 2 Chemical Properties

The data in Table (5) demonstrated that, in the two experimental seasons, no significant variation in vitamin C of the studied cultivars. However, no constant trend could be detected on total soluble solids in fruits of the studied cultivars at the two experimental seasons, whereas the lowest value of acidity at the second season and the highest value of T.S.S/ acid ratio were observed in 'Campbell' trees in the two experimental seasons.

In this respect, in the juice industry, fruits are sold based on the amount of soluble solids content and therefore the growers are interested to maximize the productivity of soluble solids [33]. TSS is an important measure of the sugar content of fruits, as sugars constitute approximately 85% of the soluble solids in citrus fruits [34]. Also, for purposes of fresh fruit sales, external appearance is more important, therefore the internal maturity factors are deemphasized, although in most areas a minimum T.SS/acid ratio is established to maintain acceptable quality. For processing, internal quality is the overriding factor, therefore juice percentage and higher T.SS/acid ratio is emphasized [35].

**Table (5): Fruit chemical properties of studied cultivars**

Cultivars	Fruit juice percent (w/w)		Vitamin C (mg/100ml)		T.S.S (%)		Acidity (%)		T.S.S/ acid ratio	
	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
Campbell	52.88 a	51.07 b	36.96 a	37.43 a	10.50 a	10.00 a	0.852 a	0.869 b	12.32 a	11.56 a
Olinda	56.73 a	57.33 a	35.36 a	35.13 a	10.00 a	10.67 a	0.855 a	1.010 a	11.70 b	10.60ab
Delta	43.89 b	44.91c	40.80 a	40.03 a	10.50 a	10.33 a	0.911 a	1.001 a	11.53 b	10.32 b


Means in each column followed by the same letter did not differ at  $p < 0.05$  according to Duncan's multiple range tests

## 3.6 Chemical composition

### 3.6.1 Total indoles

Regarding to the total indoles concentration of leaves in the studied cultivars (Fig.3), the highest values were recorded by 'Olinda' and 'Campbell' leaves at the three date of sampling at the two experimental seasons, with some exceptions. Moreover, the highest values of total indoles in March and May samples were obtained by 'Olinda' leaves. Also, it was observed that, July sample have the highest concentrations of total indoles in the studied cultivars.

In this respect, auxins promote cell enlargement rather than cell division. Also endogenous auxins increase in developing ovaries [36].



**Fig.(3): Total indoles concentration (mg/g.f.w) for studied cultivars at three date samples through two successive seasons**

### **3.6.2 Total phenols**

It was noticed that total phenols concentration was increased in March samples at the first season in the studied cultivars when compared with other sampling dates. This is may be due to phenolic compounds have been implicated in process of division, development and differentiation into new tissues [37].

Also, the highest value of total phenols concentration was recorded by 'Delta' leaves followed by 'Olinda' leaves at the three dates of sampling in both seasons when compared with 'Campbell' leaves, with some exceptions.

Furthermore, from the present results, it can be suggested that, the increasing of total indoles concentration especially in July and decreasing total phenolic compounds in 'Campbell' and 'Olinda' leaves affected positively the enhancing of tree canopy, fruit number, fruit yield and fruit quality of 'Campbell' cultivar as well as the increment of fruit size, fruit weight and fruit juice percent of 'Olinda' cultivar as compared with 'Delta' cultivar.

In this respect, polyphenolic compounds are essential for the growth of plants and affect various physiological events. They actively inhibited or stimulate some physiological process, such as defending system against pathogens and stress, growth as well as development and reproduction. Phenolic compounds have been shown to have both stimulatory and inhibitory effects on plant development [38]. Phenolic compounds are considered as bioactive non nutritional compounds, due to their antioxidant functions [39].




Fig.(4): Total phenols concentration (mg/g.fw) for studied cultivars at three date samples through two successive seasons

### 3.6.3 Total carbohydrates

The data presented in Table (6) revealed that, the highest values of total carbohydrates were recorded by 'Campbell' shoots followed by 'Olinda' shoots in the two experimental seasons.

In this respect, the availability of carbohydrates, flower intensity and the competition between them, the competition between fruitlets and fruit weight have been suggested as the most significant factors affecting the final citrus fruit size [40]. Also, a strong relationship between the carbohydrate amounts available for citrus fruitlets, especially soluble sugars, and their probability of abscission has been suggested [41]. Carbohydrates content may be a biochemical signal involved in the mechanisms controlling citrus fruit abscission [42]. Moreover, carbohydrate reserves are used in the formation and development of flowers and fruits of citrus trees [23].

Table (6): Total carbohydrates and minerals concentrations of studied cultivars

Cultivars	Total carbohydrates (mg/g. d.w.)		N (%)		P (%)		K(%)		Fe (ppm)		Mn(ppm)		Zn (ppm)	
	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
Campbell	1.820a	1.836 a	2.39a	2.40a	0.48a	0.47a	1.190a	1.280a	60.65b	58.03b	37.24b	35.60b	24.90b	23.66b
Olinda	1.426b	1.401 b	2.38a	2.67a	0.53a	0.54a	1.325a	1.310a	58.74b	55.57b	45.82a	43.71a	23.13c	23.42b
Delta	1.097c	1.106 c	2.49a	2.30a	0.49a	0.53a	1.490a	1.453a	75.93a	78.05a	33.87b	33.94b	27.02a	26.96a

Means in each column followed by the same letter did not differ at  $p < 0.05$  according to Duncans multiple range tests

### **3.6.4 Minerals**

The data in Table (6) showed that, there is no significant variation between the studied cultivars on nitrogen, phosphorus and potassium leaves concentrations in the two experimental seasons. Concerning to iron and zinc concentration, the highest values were obtained by 'Delta' leaves, while the highest value of manganese leaves concentration was recorded by 'Olinda' trees when compared with the other two cultivars in the two experimental seasons.

In this respect, availability of essential minerals during morphological and physiological process can play an important role in growth and fruit setting in Valencia orange trees [20]. Plant nutrition status has also been associated with citrus flowering [43]. The number of growing citrus fruitlets that survive after June drop is mainly determined by nutritional factors such as photo assimilates [44]. Also, fertilization play important role in the production of fruit for the fresh market and processing [45].

## **4. CONCLUSION**

In conclusion, it could be reported that, microscopically measurements and microphotographs showed distinct differences between the histological leaf and fruitlets characters of the studied cultivars signifying to affect their growth fruit characters and yielding. Whereas, 'Campbell' had the utmost values of most competent leaf tissues measurements under study i.e., palisade thickness, mid vein thickness and main vascular bundle length and width comparing to 'Olinda' and 'Delta' cvs. While, the highest values of flavedo thickness were recorded by 'Delta' fruitlets and segments width showed the highest value in 'Olinda' once. While, the thickness of the wall between segments showed the highest values in 'Campbell'. Hence, 'Campbell' trees achieved the highest values of tree canopy, fruits number, fruit yield, T.SS/ acid ratio and carbohydrates concentration. The highest values of fruit size, fruit weight as well as fruit juice percentage were recorded by 'Olinda' cvs fruits. The highest values of fruit firmness, fruit peel thickness and total phenols concentration were recorded by 'Delta' cultivar. In addition, 'Campbell' produced the highest yield and best fruit quality parameters, whereas 'Olinda' fruits gave the highest fruit juice percentage which is an extremely important parameter for its industrial processing, being also related to size were obtained by 'Olinda' Valencia trees. So, we may recommend that 'Campbell' proved as reliable high yielding cultivar with good fruit characteristics followed by 'Olinda' with advantage for juice processing under the prevailing agro-climatic conditions of South El Tahrir district, Egypt.

## **REFERENCES**

1. Khan AS, Shaheen T, Malik AU, Rajwana IA, Ahmad S, Ahmad I. Exogenous application of plant growth regulators influence the reproductive growth of *Citrus sinensis* Osbeck Cv. Blood Red Pak J Bot. (2014); 46(1):233–8.
2. Vogel S. Comparative Biomechanics: Life's Physical World. Princeton University Press. (2003);580p.
3. Valencia orange tree Varieties as affected by harvest date. Fruits. (2008); 63 (6): 327-334.
4. Maurer M. Low desert citrus varieties. The Univ. of Arizona Tucson Arizona. (1998); 1-6.
5. Demirkaser TH, Eti S, Kaplankiran M. The effects of GA<sub>3</sub> and BA treatments on fruit set and quality on Nova mandarin (in Turkish with an English summary). p. 181-184. IV

National Symposium on Horticulture (Tr), Antalya, Turkey. (2003); 8-12 September. Akdeniz University, Antalya, Turkey.

6. Obreza TA. Young Hamlin orange tree using nitrophenolate to increasing the size of Valencia fertilizer response in southwest Florida. Proc. Fla. orange fruit. State Hort. Soc. (1991);103: 12-16.
7. Nassar MA, El-Sahhar KF. Botani-cal Preparation and Microscopy (Microtech-nique), (In Arabic) Academic Bookshop, Dokki, Giza, Egypt. (1998); 219 pp.
8. A.O.A.C. Association of Official Analytical Chemists. 15<sup>th</sup>ED .Washington DC,USA. (1995); pp:490-510.
9. Larsen P. On the biogenesis of some indole compounds. Physiol. Plants. (1962); 15: 552-565.
10. Swain T, Hillis WF. The quantitative analysis of phenolic constituent. J.Sci., Food Agric. (1959); 10: 63-69.
11. Miller GL. Analytical Chemistry. (1959); 31, 426-428.
12. Plummer DT. An introduction to practical biochem. Published by Mc Graw Hill Book Company (U.K.) Limited. (1971).
13. King EJ. Micro-analysis in medical biochemistry. 2nd Ed., Churchil, London. (1951); 222 p.
14. Snedecor GW, Cochran WG. "Statistical Methods", 6<sup>th</sup> Ed. Iowa State Univ. Amess. Iowa. (1980).
15. Duncan DB. Multiple range and multiple F. Tests Biometrics. 1955;11:1-24.
16. Sedeek MS, Kirollos NF, Michel CG, Abdel-Kawy MA. Botanical and genetic characterization of *Citrus maxima* (BURM.) Merrill. F. Rutaceae. International Journal of Pharmacy and Pharmaceutical Sciences. (2017); 9(1): 260-272.
17. Sack L, Scoffoni C. Leaf venation: structure, function, development, evolution, ecology and applications in the past, present and future. New Phytologist. (2013); 198: 983–1000. [www.newphytologist.com](http://www.newphytologist.com).
18. Hostler K, Buchanon S, Zaman Q. Relating citrus canopy size and yield to precision fertilization. Proc. Fla. State Hort. Soc. (2006); 119: 148-154.
19. Zaman QU, Schumann AW, Hostler K. Estimation of citrus fruit yield using ultrasonically-sensed tree size. App. Eng. In Agric. (2006); 22(1): 39-44.
20. Chermahini SA, Moallemi N, Nabati, DA Shafieizargar AR. Winter application of foliar urea can promote some quantitative and qualitative characters of flower and fruit set of Valencia orange trees. Journal of Food, Agriculture & Environment.9 (2011); (1): 252 – 255.
21. Ruiz R, Garcíâ- Luis A, Monerri C, Guardiola J L. Carbohydrate availability in relation to fruitlet abscission in citrus. Annals of Botany. (2001); 87: 805- 812.

22. Garcia-Luis A, Guardiola JL. Influence of citrus tree internal factors and climatic effects on flowering. In: *Ninth Meeting of the International Citrus Congress*, Orlando, USA, (2000); 292-295.
23. Monerri C, Fortunato-Almeida A, Molina R V, Nebauer SG, Garcia-Luis A Guardiola, J L. Relation of carbohydrate reserves with the forthcoming crop, flower formation and photosynthetic rate, in the alternate bearing 'Salustiana' sweet orange (*Citrus sinensis* L.). *Scientia Horticulture*. (2011); 129(1): 71-78
24. Nzima MDS, Martin GC, Nishijima C. Seasonal changes in total nonstructural carbohydrate within branches and roots of naturally off and on 'Kerman' pistachio trees. *Journal of American Society for Horticultural Science*. (1997); 122(6): 856-862.
25. Goldschmidt EE. Carbohydrate supply as a critical factor for citrus fruit development and productivity. *Hort. Science*. (1999); 34(6): 1020-1024.
26. Qureshi KM, Hashim ML, Khokhar KM, Shah A H.(1993). Comparative evaluation of some sweet orange varieties at Islamabad. *Pak. J. Agri. Sci., Vol. 30, No.1*.
27. Al-Mutairi AKF. Performance of 'Olinda' Valencia orange trees grown on eight rootstocks in riyadh region. A thesis submitted in partial fulfilment of the requirements for the degree of Master of Science in Horticulture Plant Production Department, College of Food and Agricultural Sciences, King Saud University. Saudi Arabia. M. Sc Thesis.(2008).
28. Singh TC, Gill PPS. Performance of exotic Sweet orange (*Citrus sinensis* Osbeck) cultivars on different rootstocks under North Western India. *Indian Journal of Science and Technology*.(2015); *Vol 8(16)*, 59391.
29. Al-Jaleel A, Zekri M. Yield and fruit quality of " Olinda Valencia" trees grown on nine rootstocks in Saudia arabia. *Proc. Fla. State. Hort. Soc.* (2002); 115:17-22.
30. Ahmed W. Biophysical studies of combinations in "Kinnow" mandarin (*Citrus reticulata* Blanco ). Ph.D. Thesis, Ins. of Hort. Sci., Faisalabad, Pakistan. (2005).
31. Sirisomboon P, Lapchareonsuk R. Evaluation of the physicochemical and textural properties of pomelo fruit following storage. *Fruits*. (2012);67(6):399-413. <https://doi.org/10.1051/fruits/2012034>
32. Gurteg S. Correlation Studies on Fruit Traits of Some Mandarin Genotypes Grown Under Sub-Tropical Conditions of India. *J Krishi Vigyan*. (2017);6 (1):40-44.
33. Shafieizargar A, Awang Y, Juraimi A, Othman R. Yield and fruit quality of 'Queen' orange [*Citrus sinensis* (L) Osb.] grafted on different rootstocks in Iran. *Australian Journal of Crop Science*. (2012);6 (5): 777- 783.
34. Azher MN, Ahmed W, Maqbool M, Ali BS, Hussain Z, Aziz M, Shafique A. Characteristics of some potential cultivars for diversification of citrus industry of pakistan *Int. J. Agric. Appl. Sci.*(2012); (4): 1, 58.

35. Eskin NAM. Quality and Preservation of Fruits CRC Press is an imprint of Taylor & Francis Group, an In forma business. 6000 Broken Sound Parkway NW, Suite 300 Boca Raton, FL (2018);33487-2742.
36. Iglesias DJ, Cercós M, Colmenero-Flores JM, Naranjo MA, Ríos G, Carrera E, Ruiz-Rivero O, Lliso I, Morillon R, Tadeo FR, Talon M. Physiology of citrus fruiting. Braz. J. Plant Physiol. (2007); 19(4):333-362.
37. Misirli A. Bazı sert çekirdekli meyve türlerinde eşeyssel uyumsuzluk ile fenolik madde içeriği arasındaki ilişkiler. J Ege Univer Agri. Faculty. (2000); 37 (1):161-168.
38. Sulusoglu M. Phenolic compounds and uses in fruit growing. Turkish Journal of Agricultural and Natural Sciences. (2014); 947-956.
39. Reis Gaida ML. Food phenolic compounds: Main classes, sources and their antioxidant power, oxidative stress and chronic degenerative diseases-A Role for antioxidants, (ed: Dr. Jose Antonio Morales- Gonzalez, ISBN: 978-953-51-1123-8, In Tech) (2013); p: 87-112.
40. El-Otmani M, Coggins CW, Agustí Jr M, Lovatt C. Plant growth regulators in citriculture: World current uses. Critical Reviews in Plant Sciences. (2000);19:395-447.
41. Iglesias DJ, Tadeo FR, Primo-Millo E, Talon M. Fruit set dependence on carbohydrate availability in citrus trees. Tree Physiology.(2003); 23: 199-204.
42. Iglesias DJ, Tadeo FR, Primo-Millo E, Talon M. Carbohydrate and ethylene levels regulate citrus fruitlet drop through the abscission zone A during early development. Trees. (2006); 20: 348-355.
43. Albrigo LG. Effects of foliar applications of urea or nutriphite on flowering and yields of Valencia orange trees. Proc. Fla. Sta. Hort. Soc. (1999); 112:1-4.
44. Mehouchi J, Serna D, Zaragoza S, Agusti M, Talon M, Primo-Millo E. Defoliation increases fruit abscission and reduces carbohydrate levels in developing fruits and woody tissues of *Citrus unshiu*. Plant Science. (1995);107: 189-197.
45. Koo RCJ. Irrigation and fertilization effects on fruit quality, in factors affecting fruit quality, *Proc.1988 Citrus Short Course*, Lake Alfred, FL. Ferguson, J. J., and Wardowski, W. F., Eds. (1988); 35.